

Sudden confusion

Sudden confusion means the brain's thinking surface (the cerebral cortex) or its deep wake-up switch (the reticular activating system in the brainstem) just lost oxygen, fuel, or normal chemistry—find the cause and fix it in minutes, because the clock is ticking.

THE STORY IN ONE SENTENCE

Sudden confusion means the brain's thinking surface (the cerebral cortex) or its deep wake-up switch (the reticular activating system in the brainstem) just lost oxygen, fuel, or normal chemistry—find the cause and fix it in minutes, because the clock is ticking.

“Sudden confusion tells you the brain is choking on bad chemistry, starving for sugar or oxygen, or a clot or bleed just hit—check blood sugar, push oxygen, reverse opioids with naloxone if breathing is slow, then hunt the upstream break.”

Normal anatomy — the body's job

The brain stays awake and thinking when the cerebral cortex (the outer thinking layer) and the reticular activating system (a web of nerve cells in the brainstem that keeps you alert) both work together and get steady oxygen, glucose (sugar fuel), and normal blood chemistry every second.

BODY'S JOB

The cerebral cortex runs all higher thinking—language, memory, planning, controlling your muscles. The reticular activating system flips the consciousness switch on; without it, you slip into a coma even if the cortex is healthy. Both need a constant stream of oxygen (carried by red blood cells in the bloodstream) and glucose (the brain's only fuel) every second, plus stable levels of sodium, calcium, and pH (the acid-base balance in your blood). Any break in this supply—a clot, a bleed, low sugar, wrong salt levels, a toxin, or an infection—shuts the system down instantly because brain cells cannot store energy or survive even brief interruptions.

1. Your heart pumps oxygen-rich blood through arteries to the brain every second—like a delivery truck bringing groceries to a store that never closes.

2. That blood carries glucose (sugar) to every brain cell—brain cells burn glucose to make energy the way a car burns gas to run the engine.
3. The cerebral cortex (the outer layer of the brain) uses that energy to think, remember, talk, and control movement—it is the command center where all conscious activity happens.
4. The reticular activating system sits deep in the brainstem (the stalk connecting brain to spinal cord) and uses the same oxygen and glucose to stay awake—it is the on-off switch for consciousness.
5. Blood chemistry—sodium, calcium, pH—must stay in a narrow range or the electrical signals brain cells use to talk to each other go haywire, like static crackling on a radio.
6. When oxygen drops, glucose crashes, chemistry tilts, or blood flow stops, brain cells stop firing within seconds and confusion or coma appears instantly—there is no backup battery.

What breaks

One root cause—structural (a piece of the brain dies or gets squeezed), metabolic (body chemistry goes wrong), infectious (germs or inflammation hit the brain), or toxic (a drug or poison floods in)—explains all the confusion and odd behavior you see.

CLINICAL TERM

Structural brain injury: blood-starved (ischemic) stroke (a clot blocks an artery feeding the brain), bleeding-driven (hemorrhagic) stroke (a blood vessel bursts and bleeds into brain tissue or the space around it), subdural a pooled collection of blood (hematoma) (blood pools between the brain and the skull after head injury), epidural a pooled collection of blood (blood pools between the skull and the outer brain lining, usually from a skull fracture), tumor (a mass presses on brain tissue), or cerebral swelling (edema) (the brain swells from trauma, infection, or high altitude). Metabolic breaks: low blood sugar (hypoglycemia) (blood sugar below 55 mg/dL—brain cells starve), low sodium (hyponatremia) (sodium below 120 mEq/L) or high sodium (hypernatremia) (sodium above 160 mEq/L—electrical signals in the brain short-circuit), low oxygen in the blood (hypoxemia) (oxygen saturation below 90 percent—brain suffocates), hypercapnia (carbon dioxide above 50 mmHg from lung failure—acidifies the blood and brain), hepatic encephalopathy (ammonia builds up from liver failure and poisons neurons), severe waste buildup in the blood (uremia) (waste products pile up from kidney failure), thyroid storm (too much thyroid hormone speeds metabolism into chaos), or adrenal crisis (cortisol—the body's main stress hormone—drops dangerously low and blood pressure collapses). Infections: sepsis (body-wide infection triggers inflammatory chemicals that cross into the brain), bacterial meningitis (bacteria infect the meninges—the protective lining around the brain and spinal cord—most often *Streptococcus pneumoniae*, *Neisseria meningitidis*, or *Listeria monocytogenes*), or viral or autoimmune encephalitis (a virus like herpes simplex or the immune system attacks brain tissue itself). Toxins and drugs: opioid overdose (heroin, fentanyl, oxycodone slow breathing and starve the brain of oxygen), benzodiazepine overdose (Valium, Xanax, Ativan shut down the reticular activating system by boosting GABA—the brain's main calming messenger), alcohol intoxication (poisons brain cells directly by flooding GABA receptors and blocking glutamate—the brain's main wake-up messenger) or withdrawal (delirium tremens—the brain rebounds into overactivity and seizures when alcohol is stopped suddenly because GABA drops and glutamate rebounds), anticholinergic poisoning (Benadryl, many psych meds, plant toxins like jimsonweed block acetylcholine—a brain messenger needed for attention and memory—causing confusion, agitation, wide pupils, dry skin, and fast heart rate), carbon monoxide (binds to hemoglobin in red blood cells and blocks oxygen delivery), or serotonin syndrome (too much serotonin—a brain messenger that steadies mood and sleep—from mixing SSRIs with tramadol, MAOIs, or other serotonin-boosting drugs—causes confusion, agitation, fast heart, high blood pressure, muscle stiffness, and fever). Seizure-related: non-convulsive status epilepticus (a seizure that will not stop) (ongoing seizure without visible shaking, only seen on EEG brain-wave test—exhausts brain cells) or post-ictal confusion (the foggy recovery phase after a seizure ends, when the brain is drained of energy and recharging—lasts minutes to hours). The break does not stay local—downstream you see confusion,

agitation, lethargy (hard to wake), or coma (cannot wake at all). Read every sign backward to the one upstream cause and fix it.

How the body compensates

Vital signs—blood pressure, heart rate, oxygen saturation—can look stable on the monitor for minutes to hours while the brain is dying inside, because the body fights to protect the brain as long as it can. Your job is to spot the trajectory toward crash before it happens, not wait for the numbers to fall off a cliff.

1. The body tries to save the brain by shunting blood away from the skin and gut toward the brain and heart, releasing stress hormones (epinephrine and norepinephrine from the adrenal glands, which sit on top of the kidneys), and breathing faster to blow off carbon dioxide and push oxygen in.
2. This compensation phase is dangerous because the patient looks stable on the monitor—blood pressure 120/80, heart rate 90, oxygen 95 percent—while the brain is starving, swelling, or seizing inside. The numbers lie.
3. The crash point arrives suddenly: the patient loses airway protection (cannot cough or swallow, so saliva or vomit slides into the lungs—aspiration), the brain herniates (swelling or bleeding pushes brain tissue down through the foramen magnum—the opening at the skull base—and crushes the brainstem, stopping breathing and heartbeat), an unrecognized non-convulsive seizure keeps firing and exhausts the brain, or you lock onto the wrong diagnosis and miss the fixable cause (like reversible low blood sugar (hypoglycemia) or opioid overdose).
4. Watch for the crash before it tips—once herniation happens or the brain goes without oxygen for more than 4 to 6 minutes, the window to reverse damage slams shut. Dead brain cells do not come back.
5. Trend the data over time, not one snapshot: worsening Glasgow Coma Scale score (alertness drops—eyes stay closed, no words, no movement to pain), rising lactate (cells are starving and burning fuel without oxygen—anaerobic metabolism, like a car engine running without air), dropping blood sugar, or new one-sided weakness, vision loss, or pupil changes (one pupil big, one small, or one not shrinking when you shine a light) mean decompensation is underway right now.
6. A patient with stable vital signs but worsening confusion, sleepiness, or declining alertness is not stable—they are pre-crash. Do not wait for the blood pressure to drop or the oxygen to fall. By then, the brain may already be dying.

Domino effect

1. Confusion, agitation, lethargy (hard to wake), or coma (cannot wake at all) appears suddenly. — The cerebral cortex (the brain's thinking surface) and the reticular activating system (the brainstem's wake-up switch) need oxygen and glucose every second with no breaks. If blood flow stops (stroke), glucose crashes below 55 mg/dL (hypoglycemia), a toxin floods in (opioid or benzodiazepine overdose), or infection inflames the brain (meningitis or encephalitis), confusion or coma appears within seconds to minutes because neurons (brain cells) cannot fire their electrical signals anymore—like a power outage in a city.
2. One-sided weakness, vision loss, slurred speech, or unequal pupils point to structural brain injury—stroke, bleed, or tumor. — Stroke, bleed, or tumor destroys or squeezes a specific spot in the brain—left motor cortex (the strip of brain that controls movement on the right side of the body) dies and you see right-side weakness, occipital lobe (the back-of-brain region that processes vision) stroke kills vision and you see blindness or half-vision loss, brainstem stroke hits cranial nerves III, IV, or VI (the nerves that control eye movement) and you see double vision or crossed eyes. You are seeing a map of the injury printed on the body—not whole-brain sleepiness. This is a local break, not a global poison.
3. Whole-brain sleepiness and confusion without one-sided signs point to metabolic, toxic, or infectious cause poisoning the entire brain evenly. — Low blood sugar (Hypoglycemia), wrong sodium (hyponatremia or high sodium (hypernatremia)), low oxygen (hypoxemia), high carbon dioxide (hypercapnia), liver failure (hepatic encephalopathy), kidney failure (uremia), or drug overdose (opioids, benzodiazepines, alcohol) poison every neuron in the brain at once—no single spot dies more than another, so you see only confusion, agitation, and sleepiness, not one-sided weakness, vision loss, or slurred speech. The toxin or chemistry problem floods everywhere, not just one pipe.
4. Fever plus confusion raises a red alarm for infection of the brain or its lining—meningitis or encephalitis. — Bacterial meningitis (infection of the meninges—the protective layers around the brain and spinal cord) or viral encephalitis (infection of brain tissue itself, most often herpes simplex virus) triggers the hypothalamus (the brain's thermostat, sitting deep in the middle of the brain) to release pyrogens (fever-causing chemicals like interleukin-1 and prostaglandin E2) while the infection inflames neurons and blood vessels directly. Fever alone is common and often not dangerous. Fever plus confusion is a brain or brain-lining infection until proven otherwise—start antibiotics (ceftriaxone plus vancomycin) immediately, draw blood cultures, then do a lumbar puncture (spinal tap—a needle into the lower back to sample cerebrospinal fluid, the clear liquid around the brain and spinal cord) to confirm the diagnosis. Do not wait for test results to start treatment—minutes matter.

5. Asterixis (a flapping tremor when you hold your hands out like you are stopping traffic), tremor, muscle jerks (myoclonus), or abnormal posturing (arms and legs stiffen in weird positions) narrow the list of causes. — Asterixis signals liver failure (hepatic encephalopathy—ammonia poisons the basal ganglia, the brain's movement coordination center deep inside) or kidney failure (uremia—waste products disrupt the same circuits). Tremor points to alcohol withdrawal (the brain rebounds from chronic GABA boost and glutamate suppression), overactive thyroid (thyroid storm floods the body with thyroid hormone and speeds everything up), or severe electrolyte chaos (low magnesium, low calcium). Myoclonus (sudden, shock-like muscle jerks) suggests the brain just survived low oxygen (post-cardiac-arrest myoclonus), severe waste buildup in the blood (uremia) (waste products irritate the motor cortex), or ongoing seizure. Decorticate posturing (arms bend inward, legs straighten—means the cerebral cortex is damaged but the brainstem still works) or decerebrate posturing (arms and legs both straighten rigidly—means the brainstem itself is failing) are late, terrible signs of severe brain injury or herniation. These are not random—each ties back to a specific break upstream.
6. The body fights to protect the brain—agitation, fast heart rate, high blood pressure, fast breathing are survival attempts, not the disease itself. — Agitation and confusion are the brain's distress alarm that oxygen, glucose, or blood flow is failing—the brain is screaming for help. Fast heart rate (tachycardia) and high blood pressure (hypertension) are the heart and blood vessels trying to push more blood to the brain—like opening the throttle on a car struggling uphill. Fast breathing (tachypnea) blows off carbon dioxide to fix acid buildup (respiratory compensation for metabolic acidosis (acid building up in the blood)). These compensations buy you time to find and fix the cause—recognize them as red flags waving, not reassurance that the patient is stable.

7. Trace every finding backward to the one broken step, name it precisely, then pick the treatment that fixes that exact break. — Match asterixis plus confusion to liver failure (hepatic encephalopathy), treat with lactulose (a syrup that traps ammonia in the gut so it leaves in stool instead of crossing into the brain) and rifaximin (an antibiotic that kills ammonia-producing gut bacteria). Match confusion plus blood sugar 40 mg/dL to low blood sugar (hypoglycemia), give IV dextrose (sugar water) or intramuscular glucagon (a hormone that tells the liver to dump stored glucose into the blood) instantly and recheck sugar in 15 minutes. Match pinpoint pupils (pupils shrink to dots) plus slow breathing to opioid overdose, push IV or intranasal naloxone (Narcan—an opioid blocker that kicks opioids off mu-opioid receptors and reverses the overdose in 2 to 5 minutes). Match fever, stiff neck (cannot touch chin to chest—a stiff neck (nuchal rigidity)), confusion to bacterial meningitis, draw blood cultures, start IV ceftriaxone (a third-generation cephalosporin antibiotic that crosses into cerebrospinal fluid and kills *Streptococcus pneumoniae* and *Neisseria meningitidis*) plus vancomycin (covers resistant *Strep pneumoniae*) immediately, add ampicillin if the patient is over 50 or immunocompromised (covers *Listeria monocytogenes*), then do lumbar puncture to confirm and guide therapy. The pattern is always the same: find the broken upstream job, name the mechanism in plain words and clinical terms, then pick the drug, procedure, or intervention that fixes that exact break. Do not guess or treat everything at once—match the clue to the cause, then act.
8. Time is brain—every minute of confusion without treatment means more brain cells die or the window to reverse the cause closes. — Neurons (brain cells) cannot store energy and die within 4 to 6 minutes without oxygen. Low blood sugar (Hypoglycemia) below 40 mg/dL for more than 30 minutes causes permanent brain damage. Bacterial meningitis kills or disables 1 in 5 people even with treatment, and every hour you delay antibiotics raises the death rate. Blood-starved (Ischemic) stroke (clot-caused) has a 4.5-hour window for clot-busting medicine (tPA—tissue plasminogen activator, an enzyme that dissolves clots) and up to 24 hours for some mechanical clot removal (thrombectomy—threading a wire and device through an artery to pull the clot out), but the earlier you treat, the more brain you save ('time is brain'—every minute of untreated stroke kills 1.9 million neurons). Herniation (brain swelling pushes tissue down through the foramen magnum—the skull opening—and crushes the brainstem) stops breathing and heartbeat—once it happens, reversal is nearly impossible. Your job is to move fast: check blood sugar at the bedside (fingerstick glucose) in under 1 minute, push oxygen, give naloxone if you suspect opioids (no harm if you are wrong, life-saving if you are right), get a CT scan of the head without contrast (shows bleeds, strokes, tumors, swelling in 5 minutes), draw labs (complete metabolic panel for sodium, glucose, kidney and liver function; complete blood count; arterial blood gas for oxygen, carbon dioxide, and pH; lactate; blood cultures if fever), start antibiotics if you suspect meningitis (do not wait for lumbar puncture), and call neurology or the ICU early. The faster you find and fix the cause, the more brain you save. Every minute of confusion is a brain emergency until proven otherwise.

Symptoms · what the patient feels, and why

- **Disorientation to time, place, or person:** The thinking layer of the brain (the cortex) needs steady oxygen, steady sugar, and balanced chemistry to stitch together memory and what your senses see. When any of those is off—low oxygen, low blood sugar, wrong salt or calcium levels, poisons in the blood—the brain's map-making circuits fail. The person loses track of where they are, what day it is, or even who they are because the brain cannot weave the pieces together.
- **Agitation or combativeness:** When the brain senses low oxygen, low sugar, or poison, it flips into survival mode—fight or flight. The amygdala (the brain's alarm center that flags danger) fires hard and the person becomes restless, fights caregivers, or tries to run away. This is not a behavior problem; it is the brain screaming for help because it thinks it is in danger.
- **Lethargy or drowsiness (hard to keep awake):** Deep in the brainstem sits the reticular activating system (the wake-up center that keeps you alert). Poisons in the blood—like too much carbon dioxide, waste from failing kidneys (uremia), waste from failing liver (ammonia), or sedative drugs (opioids, benzodiazepines)—turn down the wake-up center's activity. The person drifts toward sleep or becomes impossible to rouse because the brain cannot stay alert.
- **Slurred speech or trouble finding words:** Language centers—usually Broca's area (the speech-making center) and Wernicke's area (the meaning-making center) on the left side of the brain—need steady blood flow and clear connections to work. When a stroke blocks blood or when whole-brain confusion scrambles the wiring, those language areas fail. The person cannot form words, cannot understand what you say, or their speech sounds garbled—like a tangled wire that cannot carry a clear signal.
- **One-sided weakness (arm, leg, or face droop):** A stroke kills brain tissue on one side—either the motor cortex (the control center for movement) or the internal capsule (the cable that carries movement signals down to the spinal cord). Dead brain tissue cannot send signals to muscles, so the opposite side of the body becomes weak or paralyzed instantly. If the left brain dies, the right arm and leg stop working; if the right brain dies, the left side stops.
- **Seizure or confusion after seizure (post-ictal confusion):** A seizure is an electrical storm—abnormal firing of neurons spreads across the cortex like lightning. It happens because of stroke, infection, wrong sodium or calcium, low glucose, or poison. After the storm stops, the exhausted brain stays confused for minutes to hours while neurons recharge—like a computer rebooting after a crash.

- **Asterixis (flapping tremor when hands held out):** When the liver or kidneys fail, poisons—especially ammonia from the liver and uremic toxins from the kidneys—build up in the blood. Those poisons disrupt the motor control centers in the basal ganglia (the brain's movement-smoothing hubs) and thalamus (the brain's relay station). The person cannot hold their hands steady when they stretch them out; instead, the hands flap or jerk—like trying to hold something steady with a weak battery.
- **Fever with confusion:** Bacterial meningitis or viral encephalitis means infection is attacking the brain or the meninges (the protective coverings around the brain). The immune system releases cytokines (alarm chemicals) that cause fever while the infection inflames and swells brain tissue. Fever plus confusion is a brain infection until proven otherwise and demands urgent spinal tap and antibiotics—minutes matter.
- **Pinpoint pupils and slow or shallow breathing:** Opioid drugs—heroin, fentanyl, prescription pain pills—lock onto mu-opioid receptors (docking sites) in the brainstem that control pupil size and the drive to breathe. The brainstem is poisoned, so the pupils shrink to tiny dots (like pencil points) and breathing slows or stops. The person can die from not breathing unless you reverse the drug with naloxone (an opioid blocker) immediately.
- **Decorticate or decerebrate posturing (arms bend or straighten rigidly):** Severe brain injury, herniation (the brain is pushed through an opening in the skull by swelling or bleeding), or profound oxygen injury damages the cortex or brainstem so badly that the brain loses control of posture. Muscles contract in abnormal, rigid patterns—arms bend inward (decorticate) or straighten stiffly (decerebrate). This signals the brain is failing and the person is near death without immediate help.

Focused exam

- **Level of consciousness:** is the person alert, sleepy, hard to wake, or completely unresponsive?: How awake someone is tells you how badly the brain is hurt right now. Two parts of the brain control wakefulness: the cerebral cortex (the wrinkly outer layer you use to think and remember) and the reticular activating system (a network of nerve fibers deep in the brainstem that acts like the brain's power switch). If either one stops working, the person can't stay awake. When someone is very sleepy or you can't wake them up, the damage is deep and serious—not just surface confusion.

- **Orientation:** can they answer four simple questions—Who are you? Where are you? What is today's date? Why are you here?: When someone is suddenly confused, they give wrong answers to these four key questions. If they know their name but not the date, or know where they are but not why, you're watching the thinking brain fail—not just sleepiness. This is real brain dysfunction. The pattern of what they remember and what they forget helps you see which brain systems are broken. Problems with recent memory (can't remember the date or why they're here) point to the hippocampus (the brain's recorder for new memories, shaped like a seahorse, deep in the temporal lobe).
- **One-sided weakness, face droop on one side, slurred speech, or sudden vision loss:** These signs point to physical damage inside the brain itself—like an blood-starved (ischemic) stroke (a blocked artery so blood can't reach part of the brain), a bleeding-driven (hemorrhagic) stroke (a burst vessel flooding brain tissue with blood), or a mass (a tumor or pocket of pus—abscess—pressing on tissue like a lump taking up space). Problems with the body's chemistry—like low blood sugar or wrong sodium levels—make the whole brain foggy but don't cause weakness on just one side. One-sided symptoms mean one specific part of the motor cortex (the strip of brain that controls movement) or the brainstem (the stalk connecting brain to spinal cord) is damaged.
- **Strange movements:** flapping tremor when holding arms out (asterixis), sudden muscle jerks (myoclonus), or stiff abnormal posturing: These movements tell you the brain is confused because the whole body's chemistry is off—metabolic encephalopathy (toxic waste or wrong salts poisoning brain cells)—or the person was exposed to a poison. Asterixis happens when support cells in the brain called astrocytes (helper cells that feed and protect neurons) are poisoned by ammonia (waste from protein breakdown) or carbon dioxide (waste gas from breathing). These movements do NOT happen with strokes or bleeding—they signal a body-wide chemical problem, not damage to one spot.
- **Fever, stiff neck that hurts when you try to bend the chin to the chest (nuchal rigidity), or both together:** Sudden confusion plus fever and a stiff neck screams infection inside or around the brain: bacterial meningitis (bacteria infecting the meninges, the three protective layers—dura, arachnoid, pia—covering the brain and spinal cord like wrapping paper), viral encephalitis (a virus attacking neurons in brain tissue itself), or sepsis (infection spreading through the bloodstream) reaching the brain. Every hour you wait raises the chance of death or permanent brain damage. The neck stiffness happens because inflamed, swollen meninges pull tight when you try to bend the neck forward—like a rope that's too short.

- Pupils: are they the same size in both eyes, and do they get smaller when you shine a light in them?: If the pupils are different sizes (anisocoria), very wide and don't react to light (fixed and dilated), or very tiny (pinpoint), pressure is building inside the skull—increased intracranial pressure (like a balloon inflating inside a closed box). This happens with bleeding, swelling, or when brain tissue shifts downward and squeezes the brainstem (herniation—the part deep inside that controls breathing and heartbeat gets crushed). Pinpoint pupils can also mean opioid poisoning (painkillers like fentanyl or heroin shutting down the brainstem). The brainstem controls the pupils through the oculomotor nerve (cranial nerve III, the cable that shrinks the pupil). This can be a brain surgery emergency.

Diagnostics · every test answers a question

- Fingerstick blood sugar (point-of-care glucose) → Is low blood sugar—low blood sugar (hypoglycemia)—making the person confused because the brain has run out of its main fuel?
- Complete metabolic panel: sodium, potassium, chloride, bicarbonate, BUN (blood urea nitrogen), creatinine, glucose, calcium → Are the blood's salts and chemicals out of balance—like sodium too low or too high, calcium too high, kidneys not cleaning waste, or blood acid-base balance way off—causing the brain to shut down?
- Arterial blood gas: oxygen (PaO₂), carbon dioxide (PaCO₂), and pH measured from blood drawn from an artery in the wrist or groin → Is the blood not carrying enough oxygen, holding too much carbon dioxide waste gas, or too acidic or alkaline for the brain to work?
- Liver function tests: ALT, AST, bilirubin, INR (how long blood takes to clot), and serum ammonia level → Is the liver failing, letting ammonia and other poisons reach the brain and cause hepatic encephalopathy (liver failure makes the brain confused)?
- Toxicology screen: urine drug screen, serum levels of acetaminophen, aspirin (salicylate), alcohol (ethanol), and toxic alcohols like methanol and ethylene glycol → Did the person take or get exposed to something poisonous—alcohol, opioids (painkillers like fentanyl or oxycodone), benzodiazepines (sedatives like lorazepam or diazepam), stimulants (like cocaine or methamphetamine), or toxic substances—that's causing confusion?
- Blood cultures: two sets drawn from separate sites (like one from each arm) before you give any antibiotics → Are bacteria swimming in the bloodstream—bacteremia—causing body-wide infection (sepsis, infection spreads everywhere with organ damage) or an infection inside the brain or the fluid around the brain causing meningitis or encephalitis?

- Non-contrast head CT: a special X-ray scan of the brain without dye, takes 5 minutes → Is there an blood-starved (ischemic) stroke (blocked artery, blood can't reach part of the brain), a bleeding-driven (hemorrhagic) stroke (burst vessel, blood floods brain tissue), a subdural or epidural a pooled collection of blood (hematoma) (blood pooling between the skull and brain from trauma), a mass (tumor or abscess pressing on tissue), or hydrocephalus (fluid backup raising pressure inside the skull)?
- Lumbar puncture (spinal tap) with cerebrospinal fluid analysis: cell count with differential (types of white cells), glucose, protein, Gram stain (dye to see bacteria under microscope), bacterial culture, and PCR for herpes simplex virus → Is there bacterial meningitis (bacteria infecting the meninges, the layers around the brain and spinal cord), viral encephalitis (a virus like herpes simplex attacking neurons in brain tissue), or subarachnoid bleeding (hemorrhage) (bleeding in the space between the arachnoid layer and brain surface)?

Treatment logic

- Check fingerstick blood sugar immediately. If it's below 70 mg/dL, give thiamine 100 mg into the vein (IV) first, then give 50% dextrose (D50W) 50 mL IV push—that's 25 grams of sugar. → Low blood sugar (hypoglycemia) shuts down neurons instantly because they can't make ATP (the energy packets cells burn to do work) without glucose. Neurons have almost no stored fuel—they live minute to minute on what the blood delivers. Giving dextrose (sugar water) restores fuel so neurons start firing again within 2 to 5 minutes—confusion often lifts right before your eyes. You give thiamine (vitamin B1) first in people who drink heavily or are malnourished to prevent Wernicke encephalopathy—a sudden, permanent brain injury that destroys the mammillary bodies (two tiny bumps in the hypothalamus that help make memories) and causes confusion, eye muscle paralysis (can't move eyes side to side), and loss of balance (ataxia) if you give sugar without thiamine first. This is because neurons need thiamine to burn glucose—without it, flooding the brain with sugar can push the last bit of thiamine into making energy and starve other critical brain areas.

- Draw emergency blood tests (stat labs) immediately: complete metabolic panel (sodium, potassium, chloride, bicarbonate, BUN, creatinine, glucose, calcium), liver function tests with serum ammonia, arterial blood gas, complete blood count, toxicology screen including acetaminophen and aspirin levels, two sets of blood cultures from separate sites, and troponin (heart damage marker—confusion can be the only sign of a silent heart attack in older adults). → These tests find the hidden cause poisoning the brain: sodium below 120 mEq/L causes cerebral swelling (edema) (brain swelling), ammonia above 100 mcg/dL poisons astrocytes (support cells for neurons), PaO₂ below 60 mmHg starves neurons of oxygen, PaCO₂ above 50 mmHg sedates the reticular activating system (the wake-up switch in the brainstem), pH below 7.20 breaks enzyme function (the workers inside cells can't do their jobs), bacteria in blood means sepsis or meningitis, high troponin means heart attack (myocardial tissue death from blocked blood flow (infarction)) cutting blood flow to the brain. The results tell you exactly what chemical or infection is poisoning the brain so you can fix the RIGHT problem. Treating blindly without labs is guessing—you might fix one thing while missing the real killer.
- Get a non-contrast head CT scan emergently (right away, within 15 to 30 minutes) if the patient has focal weakness (one-sided weakness, face droop, or slurred speech), sudden severe headache (worst headache of life), recent head injury (fell and hit head in past 3 months), takes anticoagulants (blood thinners like warfarin, apixaban, rivaroxaban, dabigatran), has a history of cancer (can spread to brain), age over 60, or confusion started suddenly during or right after physical effort. → The scan finds physical damage in the brain that needs emergency treatment. Blood-starved (Ischemic) stroke from a blocked artery needs thrombolysis (clot-busting medicine—IV alteplase 0.9 mg/kg, max 90 mg, within 4.5 hours of symptom start per AHA/ASA 2019 guidelines) or mechanical thrombectomy (threading a wire device up through the groin artery to the brain to pull the clot out—within 6 to 24 hours if it's a large vessel like the middle cerebral artery per AHA/ASA 2018 guidelines). Bleeding-driven (Hemorrhagic) stroke needs immediate blood pressure control (lower systolic to 140 mmHg within 1 hour per AHA 2022 ICH guidelines) to stop the bleeding from getting bigger, and possibly surgery to drain the blood. Subdural a pooled collection of blood (hematoma) (blood between the dura and brain) may need surgical drainage if it's over 10 mm thick or causing midline shift. Brain tumor may need dexamethasone (a steroid, 10 mg IV, to shrink swelling around the mass) and neurosurgery consultation. Hydrocephalus may need an external ventricular drain (a tube placed through a small skull hole into the fluid spaces to let pressure out).

- If the patient has fever (over 38.0°C or 100.4°F), stiff neck (nuchal rigidity), pinpoint bleeding spots (petechiae) (tiny purple or red spots on skin from burst capillaries that don't fade when you press on them), immunosuppression (weak immune system from HIV with CD4 below 200, chemotherapy, chronic steroids, or transplant drugs), or if you even suspect meningitis or encephalitis, give antibiotics and antivirals immediately: ceftriaxone 2 grams IV PLUS vancomycin 15–20 mg/kg IV (dose depends on kidney function, usually 1 to 2 grams) PLUS acyclovir 10 mg/kg IV every 8 hours (given slowly over 1 hour to protect kidneys). Do NOT wait for a lumbar puncture (spinal tap) if it can't be done within 30 minutes—every hour of delay increases death and permanent brain damage. If the patient is over 50 years old, has cancer, takes immune-suppressing drugs, or abuses alcohol, add ampicillin 2 grams IV every 4 hours to cover *Listeria monocytogenes* (a germ from contaminated deli meat, soft cheese, or hot dogs that invades the brainstem). → Bacterial meningitis kills 15–30% of patients (per Cochrane 2015 systematic review and IDSA 2004 bacterial meningitis guidelines) and causes permanent brain damage—deafness (the infection destroys the cochlear nerve, cranial nerve VIII), seizures, stroke from inflamed vessels, intellectual disability—in 30–50% of survivors if you delay antibiotics even 1 hour. Ceftriaxone (a third-generation cephalosporin) kills the two most common bacteria—*Streptococcus pneumoniae* (pneumococcus, a round germ that lives in the throat) and *Neisseria meningitidis* (meningococcus, a bean-shaped germ spread by coughing). Vancomycin kills resistant pneumococcus strains (PRSP, penicillin-resistant *Streptococcus pneumoniae*) that don't die from ceftriaxone—now 30–40% of pneumococcus in the U.S. per CDC 2022 data. Acyclovir treats herpes simplex virus encephalitis (HSV, usually HSV-1, attacks the temporal lobes—the parts of the brain on the sides above your ears—and causes brain swelling and tissue death). Without acyclovir, HSV encephalitis kills 70%; with it, mortality drops to 15–20% (per IDSA 2017 guidelines). Ampicillin covers *Listeria monocytogenes*, which invades the brainstem (especially the pons and medulla) and doesn't respond to ceftriaxone. Every hour of delay increases the bacterial load (number of germs) and the inflammatory damage (cytokines and pus destroy brain tissue and blood vessels).

- Fix severe salt imbalances carefully and slowly: If sodium is below 120 mEq/L with seizures, coma, or severe symptoms (acute symptomatic low sodium (hyponatremia)—happened in less than 48 hours), give 3% hypertonic saline (very salty water) 100 mL IV over 10 minutes, then recheck sodium in 1 hour and repeat if needed. If sodium is 120–130 mEq/L without seizures or severe symptoms, give 0.9% normal saline IV at 75–125 mL/hour. Never raise sodium faster than 8 mEq/L in the first 24 hours or 18 mEq/L in the first 48 hours (per NEJM 2015 low sodium review and European guidelines 2014) to avoid osmotic demyelination syndrome. If potassium is below 3.0 mEq/L (severe low potassium (hypokalemia)), give potassium chloride 40 mEq IV diluted in saline over 4 hours (max rate 10 mEq/hour through a peripheral IV, 20 mEq/hour through a central line). If magnesium is below 1.5 mg/dL, give magnesium sulfate 2 grams (16 mEq) IV over 15 minutes. If ionized calcium is below 1.0 mmol/L (total calcium below 7.5 mg/dL—severe low calcium (hypocalcemia)), give calcium gluconate 1 to 2 grams (10–20 mL of 10% solution) IV over 10 minutes while watching the heart monitor for arrhythmias. → Severe low sodium (hyponatremia) (low sodium) causes cerebral swelling (edema) (brain swelling) because water floods into brain cells down the osmotic gradient (water moves to where there's more salt to try to balance things out)—neurons swell, can't fire, and if they swell too much, they press against the skull and cause herniation (brain gets squeezed and stops working). Correcting sodium slowly and carefully restores normal osmotic balance so water stays in the bloodstream where it belongs and leaves brain cells. Low potassium (below 3.0 mEq/L) and low magnesium (below 1.5 mg/dL) disrupt the sodium-potassium ATPase pump (the machine in every neuron membrane that uses ATP energy to pump sodium out and potassium in, keeping the cell ready to fire an electrical signal) and cause abnormal heart rhythms like ventricular a fast heart rate (tachycardia) and torsades de pointes (a twisting dangerous rhythm that can become cardiac arrest). Low calcium (below 7.5 mg/dL) makes neurons fire too easily—hyperexcitability—causing tetany (muscle spasms in hands and face), seizures, and prolonged QT interval on EKG (heart rhythm abnormality). But fixing sodium too fast—more than 8 mEq/L in 24 hours—causes osmotic demyelination syndrome (also called central pontine myelinolysis)—permanent brainstem and pontine damage that destroys myelin (the fatty insulation around nerve fibers, like the rubber coating on a wire) and causes locked-in syndrome (awake but can't move or speak), difficulty swallowing, and movement problems that never fully recover. This happens because brain cells that shrank to adapt to low sodium can't re-expand fast enough when sodium rises quickly—they tear apart.

- If liver failure with elevated ammonia is causing hepatic encephalopathy, give lactulose 30 mL (20 grams) by mouth or through a nasogastric tube (feeding tube through the nose into the stomach) every 1 to 2 hours until the patient has 2 to 3 loose, mushy bowel movements per day, then adjust dose to keep 2–3 stools daily (usually 30–45 mL three times daily). Also give rifaximin 550 mg by mouth twice daily (per AASLD 2014 hepatic encephalopathy practice guidelines and NEJM 2010 rifaximin trial). → Lactulose (a synthetic sugar that humans can't digest) traps ammonia in the intestines by two mechanisms: First, gut bacteria break lactulose down into acids that lower stool pH (making it acidic, around pH 5–6). In acidic conditions, ammonia (NH_3 , which can cross from gut into blood) turns into ammonium (NH_4^+ , a charged form that can't cross gut walls) and gets stuck in the stool. Second, lactulose acts as an osmotic laxative (pulls water into the intestines) and flushes ammonia-filled stool out before ammonia can be absorbed. Rifaximin (a gut-selective antibiotic that stays in the intestines and isn't absorbed into the bloodstream) kills ammonia-producing gut bacteria—especially *Clostridium* species, *Bacteroides*, and *Escherichia coli*—that break down protein from food into ammonia. Together, lactulose and rifaximin lower blood ammonia levels from over 100–200 mcg/dL down to 40–60 mcg/dL and reverse confusion in 24 to 72 hours by stopping ammonia from poisoning astrocytes (support cells in the brain that clear glutamate, a neurotransmitter, and maintain the blood-brain barrier). High ammonia causes astrocytes to swell—cytotoxic swelling (edema)—and disrupts neurotransmission. Rifaximin reduces hepatic encephalopathy episodes by 58% compared to placebo (per NEJM 2010 trial, 299 patients).

- If the patient is agitated, combative, trying to pull out IV lines or tubes, or at risk of hurting themselves or staff, give low-dose antipsychotics cautiously: haloperidol 0.5–2 mg IV or intramuscular (IM shot into muscle)—but first check the QTc interval on an EKG (must be below 500 milliseconds to avoid torsades de pointes, a deadly heart rhythm)—or give a short-acting benzodiazepine like lorazepam 0.5–1 mg IV slowly over 1–2 minutes. Avoid sedatives if possible until you know the cause—sedation can hide worsening brain injury. If the patient has a seizure or witnessed seizure before arrival, give lorazepam 4 mg IV over 2 minutes (0.1 mg/kg, max 4 mg per dose), then load with fosphenytoin 20 mg phenytoin equivalents per kilogram (PE/kg) IV at a rate no faster than 150 PE/minute (or phenytoin 20 mg/kg IV at max 50 mg/minute) to prevent another seizure. → Haloperidol (a typical antipsychotic) blocks dopamine D2 receptors (docking sites for dopamine, a brain chemical that controls movement, arousal, and thought) in the basal ganglia (deep movement-control centers) and limbic system (emotional brain centers including the amygdala and hippocampus) to calm agitation and stop hallucinations without slowing breathing—no respiratory depression (lungs still work normally, unlike opioids or high-dose benzodiazepines). Lorazepam (a benzodiazepine) boosts GABA-A receptors (docking sites for GABA—gamma-aminobutyric acid—the brain's main 'stop' chemical that calms neurons by letting chloride ions flow into cells and making them less excitable) to halt seizure activity by stopping neurons in the motor cortex (the strip of brain from ear to ear over the top that controls movement) from firing out of control. Fosphenytoin (a prodrug that turns into phenytoin in the body) blocks voltage-gated sodium channels in neuron membranes (the doors that let sodium rush in to start an electrical signal) to prevent another seizure for 12 to 24 hours—buys you time to find and treat the root cause. Controlling severe agitation protects the patient from falling, pulling out life-saving tubes (breathing tube, IV lines, bladder catheter), or hitting staff. Stopping seizures prevents status epilepticus (a seizure that will not stop) (continuous seizure activity lasting over 5 minutes that causes permanent neuron death in the hippocampus, cerebellum, and cortex—happens when neurons run out of ATP and glutamate excitotoxicity kills cells).

Medications

Dextrose (sugar water given through a vein) · Emergency glucose replacement

The brain runs on sugar (glucose) the way a car runs on gas. If blood sugar drops too low (hypoglycemia), the brain cannot think or stay awake. Dextrose is pure sugar that goes straight into the bloodstream and reaches the brain in seconds, bringing the fuel level back up so brain cells can work again.

1. Sugar enters the blood through the IV line.
2. Sugar travels to every brain cell.

3. Brain cells pull the sugar inside and burn it to make energy (ATP, the body's power molecule).
4. The person wakes up and thinks clearly again.

SAFETY WATCH

ALWAYS give thiamine (vitamin B1) BEFORE or WITH dextrose if the person drinks a lot of alcohol, is malnourished, or has not eaten for days — otherwise dextrose can trigger Wernicke encephalopathy (a serious brain injury from lack of B1). · Check blood sugar with a fingerstick test before giving dextrose — do NOT guess. · Watch for a burning feeling at the IV site; dextrose is thick and can hurt the vein.

Thiamine (vitamin B1) · Essential vitamin replacement

Thiamine is a helper molecule (coenzyme) that brain cells need to turn sugar into energy. Without enough thiamine, the brain cannot burn glucose properly — even if sugar is in the blood. People who drink heavily, eat very little, or have certain illnesses run out of thiamine. Giving thiamine restores the missing helper so brain cells can make energy again.

1. Thiamine enters the bloodstream (usually given IV or IM).
2. It travels to brain cells and goes inside.
3. It joins enzymes (like alpha-ketoglutarate dehydrogenase and transketolase) that break down glucose into energy.
4. Brain cells start making energy normally again.

SAFETY WATCH

Give thiamine BEFORE giving dextrose in anyone who drinks alcohol heavily or has poor nutrition — dextrose can worsen brain injury (Wernicke encephalopathy) if thiamine is low. · Allergic reactions are rare but can happen; watch for rash, itching, or trouble breathing. · Give 500 mg IV (or 250 to 500 mg IM) before or with the first dose of dextrose.

Naloxone (Narcan) · Opioid receptor blocker (opioid antagonist)

Opioid drugs (like fentanyl, heroin, oxycodone, morphine) lock onto mu opioid receptors in the brain and slow down breathing and thinking until the person becomes unconscious. Naloxone is shaped like an opioid but does NOT turn the receptor on — it just blocks the spot. It kicks the opioid off the receptor, and within 1 to 3 minutes the person wakes up and starts breathing normally again.

1. Naloxone enters the bloodstream (IV, IM, nasal spray, or under the tongue).
2. It travels to the brain and spinal cord.
3. It bumps opioid molecules off their receptors and blocks the spots.

4. The medulla wakes up and restarts normal breathing, and the cortex wakes up so the person becomes alert.

SAFETY WATCH

Naloxone triggers sudden opioid withdrawal (sweating, shaking, vomiting, pain, agitation) — the person may be very uncomfortable or angry when they wake up. · Watch for return of overdose symptoms after 30 to 60 minutes; be ready to give another dose. · Do NOT use naloxone as a test to see if opioids are the problem — it can cause serious withdrawal in people who depend on opioids.

Flumazenil (Romazicon) · Benzodiazepine receptor blocker (benzodiazepine antagonist)

Benzodiazepines (like midazolam, lorazepam, diazepam) attach to GABA-A receptors in the brain and boost GABA (the brain's main calming signal), slowing down electrical activity and making the person sleepy or unconscious. Flumazenil blocks those receptors and reverses the sedation. It works within 1 to 2 minutes.

1. Flumazenil enters the bloodstream (given IV).
2. It travels to the brain and blocks GABA-A receptors.
3. Benzodiazepine molecules are bumped off the receptors.
4. Brain electrical activity speeds back up, and the person wakes up.

SAFETY WATCH

DO NOT USE if the person took pills at home, on the street, or in an unknown setting — flumazenil can trigger a dangerous seizure if the person also took seizure-causing drugs (like cocaine, tricyclic antidepressants) or has epilepsy and the benzodiazepine was preventing seizures. · ONLY use in the hospital if a doctor gave the benzodiazepine for a procedure and you know exactly what and how much was given. · Watch for seizures, anxiety, and agitation after giving flumazenil.

Ceftriaxone · Third-generation cephalosporin antibiotic

Ceftriaxone kills bacteria by breaking down their cell walls. Bacteria need strong walls to survive — without them, they burst and die. Ceftriaxone crosses the blood-brain barrier into the brain and spinal fluid and kills most bacteria that cause meningitis (infection of the lining around the brain) and encephalitis (infection inside the brain itself). It works on *Streptococcus pneumoniae*, *Neisseria meningitidis*, *Haemophilus influenzae*, and many other common germs.

1. Ceftriaxone enters the bloodstream (given IV).
2. It crosses into the brain and spinal fluid.

3. It attaches to penicillin-binding proteins (enzymes that build bacterial cell walls).
4. Bacteria cannot finish building their walls, so they burst and die.

SAFETY WATCH

Start ceftriaxone immediately (within 30 minutes of arriving at the hospital) if meningitis or encephalitis is suspected — do NOT wait for test results. · Allergic reactions can happen (rash, itching, swelling, trouble breathing); ask about penicillin or cephalosporin allergies. · Can cause diarrhea (including *C. difficile* infection). · Do NOT mix with calcium-containing IV fluids in newborns — can form dangerous crystals.

Vancomycin · Glycopeptide antibiotic

Vancomycin kills bacteria that are resistant to other antibiotics, especially MRSA (methicillin-resistant *Staph aureus*) and resistant *Streptococcus pneumoniae*. It works by blocking the final step of building the bacterial cell wall, so the bacteria cannot finish their protective coat and they die. It crosses into the brain and spinal fluid, especially when the brain lining is inflamed.

1. Vancomycin enters the bloodstream (given IV).
2. It crosses into inflamed brain tissue and spinal fluid.
3. It binds to the building blocks of the bacterial cell wall and stops the final assembly step.
4. Bacteria cannot finish their walls, so they burst and die.

SAFETY WATCH

Give vancomycin WITH ceftriaxone when treating suspected bacterial meningitis to cover resistant bacteria. · Can damage the kidneys — check blood levels and kidney function regularly. · Can cause red man syndrome (flushing, itching, low blood pressure) if given too fast — infuse slowly over at least 60 minutes. · Can damage hearing (ototoxicity) — higher risk if combined with other drugs that hurt the ears or kidneys.

Ampicillin · Penicillin antibiotic

Ampicillin kills bacteria by punching holes in their cell walls. Bacteria need a strong wall to hold their shape and survive — without it, they burst and die. Ampicillin crosses the blood-brain barrier and is one of the few antibiotics that kills *Listeria monocytogenes*, a dangerous germ that causes meningitis (brain lining infection) in older adults, newborns, pregnant women, and people with weak immune systems. It also works on many other bacteria like *Streptococcus* and some *E. coli* strains.

1. Ampicillin enters the bloodstream (given IV).
2. It crosses into the brain and spinal fluid.
3. It attaches to penicillin-binding proteins (the enzymes that build bacterial cell walls).

4. Bacteria cannot finish building their walls, so they burst and die.

SAFETY WATCH

Give ampicillin WITH ceftriaxone and vancomycin when treating suspected bacterial meningitis in older adults (over 50), newborns, or people with weak immune systems to cover *Listeria*. · Allergic reactions can happen (rash, itching, swelling, trouble breathing); ask about penicillin allergies. · Can cause a flat, red rash (especially in people with mononucleosis or taking allopurinol) — this is NOT always a true allergy. · Can cause diarrhea (including *C. difficile* infection).

Acyclovir · Antiviral (nucleoside analog)

Acyclovir stops herpes simplex virus (HSV) from copying itself inside brain cells. HSV is the most common virus that causes encephalitis (brain infection). Acyclovir looks like a building block of viral DNA, so the virus grabs it and tries to use it to make more viruses — but acyclovir is a fake block that stops the copying machine cold. The virus cannot multiply, and the immune system has time to clear it out.

1. Acyclovir enters the bloodstream (given IV).
2. It crosses the blood-brain barrier into the brain.
3. Infected brain cells activate acyclovir (using viral thymidine kinase) and turn it into its active form (acyclovir triphosphate).
4. The virus tries to use acyclovir to build new DNA but gets stuck — copying stops.
5. The virus cannot make more copies, and the infection is controlled.

SAFETY WATCH

Start acyclovir immediately if viral encephalitis (especially herpes) is suspected — do NOT wait for test results; every hour of delay increases the risk of permanent brain damage or death. · Can damage the kidneys — give IV fluids and check kidney function. · Can cause confusion or seizures if it builds up in people with kidney disease. · Can cause vein irritation at the IV site.

Lorazepam (Ativan) · Benzodiazepine (GABA enhancer)

A seizure is an electrical storm in the brain — neurons fire out of control. Lorazepam boosts the effect of GABA (the brain's main calming signal), which tells neurons to slow down. It opens chloride channels on neurons, making them harder to fire. This calms the electrical storm and stops the seizure. Lorazepam works within 1 to 3 minutes when given IV and is the first-line treatment to stop an ongoing seizure.

1. Lorazepam enters the bloodstream (given IV or IM).

2. It crosses the blood-brain barrier into the brain.
3. It attaches to GABA-A receptors and makes them more sensitive to GABA.
4. Chloride flows into neurons, making them less excitable.
5. The seizure slows and stops.

SAFETY WATCH

Can slow breathing — watch closely, especially if the person also took opioids or alcohol. · Can drop blood pressure. · Can cause confusion or sleepiness after the seizure stops. · If the seizure does NOT stop after two doses of lorazepam (total 8 mg in adults), move to a second-line drug like levetiracetam, fosphenytoin, or valproate.

Levetiracetam (Keppra) · Antiepileptic (seizure-preventing drug)

Levetiracetam calms overactive neurons by binding to a protein called SV2A inside brain cells. SV2A sits on tiny bubbles (synaptic vesicles) that store glutamate (the brain's main excitatory signal). By attaching to SV2A, levetiracetam slows down how much glutamate neurons release. Less glutamate means less excitation and less chance of a seizure storm starting or spreading.

1. Levetiracetam enters the bloodstream (given IV or by mouth).
2. It crosses the blood-brain barrier into the brain.
3. It binds to SV2A proteins in nerve endings.
4. Neurons release less glutamate.
5. Seizure activity is prevented or stopped.

SAFETY WATCH

Generally safe; fewer drug interactions and side effects than older seizure medicines. · Can cause mood changes (irritability, depression, anxiety) in some people. · Reduce the dose in people with kidney disease. · Rarely causes serious skin rash (Stevens-Johnson syndrome) — stop the drug and call a doctor if a rash appears.

Haloperidol (Haldol) · Typical antipsychotic (dopamine receptor blocker)

In severe agitation (like delirium or psychosis), too much dopamine activity in certain brain circuits makes the person see or hear things that are not there, believe false things, or become violent. Haloperidol blocks dopamine D2 receptors and calms the overactive circuits, reducing agitation, hallucinations, and delusions.

1. Haloperidol enters the bloodstream (given IV, IM, or by mouth).
2. It crosses the blood-brain barrier into the brain.
3. It blocks dopamine from attaching to D2 receptors.

4. Overactive dopamine circuits calm down.
5. Agitation, hallucinations, and delusions decrease.

SAFETY WATCH

DO NOT USE if the person has Lewy body dementia or Parkinson disease — can cause severe muscle stiffness and worsening confusion. · Check the heart rhythm (QTc interval on ECG) before giving haloperidol — if QTc is long, haloperidol can trigger a dangerous heart rhythm (torsades de pointes). · Can cause muscle stiffness, tremor, and restlessness (extrapyramidal symptoms). · Can cause neuroleptic malignant syndrome (rare but life-threatening: high fever, muscle rigidity, confusion, unstable vital signs). · Older adults are more sensitive — use the smallest dose possible and only when absolutely necessary. · Black-box warning: Increases death risk in older adults with dementia — use only when non-drug approaches fail and the person is a danger to themselves or others.

Red flags

- One-sided weakness, vision loss, or slurred speech (stroke or brain bleed—get a CT scan of the head and call neurology immediately)
- Fever with stiff neck or severe headache (bacterial meningitis—draw blood cultures, start intravenous ceftriaxone plus vancomycin, then do a lumbar puncture to test spinal fluid)
- Pinpoint pupils and slow breathing (opioid overdose—give naloxone intravenously or intramuscularly now)
- Posturing, unequal pupils, or Glasgow Coma Scale 8 or lower (herniation or severe brain injury—intubate, give intravenous hypertonic saline or mannitol to drop brain pressure, call neurosurgery)
- Blood glucose below 55 mg/dL (hypoglycemia—give intravenous dextrose or inject glucagon immediately)

Nursing priorities

1. Check the fingerstick blood sugar (glucose) immediately at the bedside, before waiting for any laboratory results. — Low blood sugar (hypoglycemia below 70 mg/dL) starves the brain of fuel (glucose) and causes confusion, seizures, coma, and permanent brain damage within minutes. It is the fastest reversible cause—giving intravenous dextrose (sugar) can wake the patient up in seconds.

2. Check vital signs (blood pressure, heart rate, respiratory rate, temperature, and oxygen saturation) and put the patient on continuous pulse oximetry and cardiac telemetry monitoring. — Low oxygen (hypoxia) or low blood pressure (hypotension) means the brain is not getting enough oxygen-rich blood, causing confusion. Fever points to infection like meningitis or sepsis. Slow breathing (respiratory depression) signals opioid overdose or rising pressure inside the skull (increased intracranial pressure). Abnormal heart rhythms (arrhythmias) or very fast or slow heart rate can drop blood flow to the brain.
3. Protect the patient from falling by raising both side rails, placing the bed in the lowest position, and keeping the call light within reach; consider a 1-to-1 sitter if the patient is combative or trying to pull out tubes. — Confused patients do not understand where they are or that they are too weak to walk safely. They try to climb out of bed and fall, causing head bleeds (subdural or a pooled collection of blood (hematoma)), broken hips, or other injuries. Falls are the number-one preventable harm in hospitals for patients with altered mental status (confused thinking).
4. Establish intravenous (IV) access with at least one large-bore catheter (18-gauge or 20-gauge) and draw all stat (emergency) blood work at the same time: complete metabolic panel, liver function tests with ammonia level, arterial blood gas, complete blood count, toxicology screen, and troponin. — The IV line lets you give emergency medicines (dextrose for low sugar, naloxone for overdose, antibiotics for meningitis, benzodiazepines for seizures) within seconds. Drawing blood at IV start avoids extra needle sticks. Blood tests show the cause: low sodium (hyponatremia), high calcium (hypercalcemia), kidney failure (uremia), liver failure (high ammonia), low oxygen or high carbon dioxide (from the blood gas), infection (high white blood cells), or heart attack (high troponin).
5. If blood sugar is below 70 mg/dL, give thiamine 100 milligrams into the vein first, then immediately give 50 milliliters of 50% dextrose (D50W)—that is 25 grams of sugar—by slow IV push over 3 to 5 minutes while watching for IV infiltration (the IV fluid leaking into the tissue instead of the vein). — Thiamine (vitamin B1) must go in before sugar in anyone who might drink alcohol heavily or be malnourished, because giving sugar without thiamine can trigger Wernicke encephalopathy—sudden permanent brain damage causing confusion, abnormal eye movements, and trouble walking. D50W is very thick (hyperosmolar) and burns tissue badly if it leaks outside the vein, so check the IV site while you push it and stop immediately if you see swelling.

6. Perform a focused neurological assessment every 15 to 30 minutes: check if the patient knows their name, where they are, and what day it is (orientation times three); check pupil size and reaction to light; check for face droop, arm drift (ask the patient to hold both arms up with eyes closed—one weak arm drifts down), and leg strength; and use the Glasgow Coma Scale to track if the patient is getting better or worse. — Sudden worsening—like new one-sided weakness, unequal pupils (one bigger than the other, called unequal pupils (anisocoria)), or decreasing Glasgow Coma Scale score—means stroke is growing, brain bleed is expanding, or pressure inside the skull is rising (herniation starting). This requires immediate CT scan and neurosurgery or neurology consultation, because the patient may need surgery or clot-busting medicine.
7. If the patient has pinpoint pupils (very tiny pupils) and slow breathing under 10 breaths per minute, give naloxone 0.4 milligrams IV push immediately and be ready to bag-mask ventilate (use an Ambu bag to breathe for the patient) if breathing stays slow or stops. — Opioid drugs (heroin, fentanyl, oxycodone, morphine) activate receptors in the brainstem that slow breathing and shrink pupils. Naloxone blocks those receptors and reverses the overdose within 1 to 2 minutes, but it wears off in 30 to 90 minutes—so the patient can stop breathing again if the opioid is long-acting. Without naloxone or breathing support, the brain does not get oxygen (hypoxia) and dies in 4 to 6 minutes.
8. Keep the patient NPO (nothing by mouth—no food, liquid, or pills) until the cause of confusion is found and the patient is fully awake enough to protect their own airway. — Confused patients cannot swallow safely—they aspirate (breathe in) food, water, or pills into the lungs, causing aspiration pneumonia or choking. Their gag reflex and cough reflex (protective airway reflexes) are weak. If the confusion gets worse, they may need emergency intubation (breathing tube), and a full stomach increases the risk of vomiting and aspiration during intubation.

Sources

- American College of Emergency Physicians (ACEP) Clinical Policy on Acute Altered Mental Status
- American Academy of Neurology (AAN) Practice Parameter for Coma and Altered Consciousness
- Infectious Diseases Society of America (IDSA) Bacterial Meningitis Guidelines
- American Heart Association/American Stroke Association (AHA/ASA) Acute Ischemic Stroke Guidelines

Last reviewed: 2026-07-07