

CLABSI

Bacteria crawl into the bloodstream through a central line (a large IV tube placed near the heart), causing a blood infection.

THE STORY IN ONE SENTENCE

Bacteria crawl into the bloodstream through a central line (a large IV tube placed near the heart), causing a blood infection.

“CLABSI means germs sneak into your blood by hitchhiking along a central line—the line is guilty until you prove it innocent.”

Normal anatomy — the body's job

The blood inside your veins is sterile (germ-free). A central line is a long, thin tube (catheter) that a doctor slides through your skin into a large vein near your heart (such as the internal jugular vein in your neck, the subclavian vein under your collarbone, or the femoral vein in your groin). This line lets doctors deliver medicine, fluids, or nutrition straight into your bloodstream. Your skin is a barrier that normally keeps bacteria out; the line breaks that barrier.

BODY'S JOB

Your skin acts like a fortress wall, keeping billions of bacteria that live on your skin from getting inside your body. Inside your veins, blood flows sterile, and white blood cells (immune soldiers) patrol constantly, hunting for any invaders. The lining of your veins (endothelium) is smooth and slippery so nothing sticks to it. This setup keeps your bloodstream germ-free under normal conditions.

1. Skin forms a continuous, waterproof barrier—bacteria live on the surface but cannot cross into the bloodstream.
2. When a central line is placed, it punctures the skin and stays in place, creating a permanent opening (a bridge from outside to inside).
3. The catheter tip sits in a large vein (such as the superior vena cava, the main vein draining into the heart), delivering fluids or medicine directly into fast-flowing blood.

4. Bacteria on the skin can slide down the outside of the catheter (along the insertion tract) or enter through the catheter hub (the connector where nurses attach IV tubing) and travel into the bloodstream.
5. Over time, bacteria stick to the catheter surface (inside or outside) and form biofilm—a slimy coating that shields them from antibiotics and the immune system, like a fortress for germs.

What breaks

Bacteria living on the skin or hands sneak into the bloodstream by traveling along the central line or through its connectors. Once inside, they multiply and release toxins that make you sick. The line itself becomes a hiding place (biofilm) that antibiotics cannot reach.

CLINICAL TERM

Bacteria colonize (settle and grow on) the external catheter surface at the skin insertion site (extraluminal route) or contaminate the internal catheter lumen (the hollow inside of the tube) via hub manipulation or contaminated connectors (intraluminal route). Biofilm—a glycocalyx matrix (slimy coating of bacterial proteins and sugars)—forms on the catheter surface within 24–48 hours, protecting bacteria from host immune cells and antibiotics. Bacteria then seed the bloodstream, causing catheter-related bloodstream infection (CR-BSI or CLABSI). The line is the portal of entry and the presumed source until blood cultures prove otherwise.

How the body compensates

Early in infection, your body can keep your vital signs near-normal by raising your heart rate and breathing rate to deliver oxygen to tissues fighting the infection. This masks how sick you really are, hiding the fact that you are on the edge of collapse.

1. {'what': 'Heart rate increases (tachycardia—usually over 90–100 beats per minute)', 'why': 'Your heart pumps faster to deliver more oxygen and immune cells (white blood cells) to tissues fighting the infection. This keeps blood pressure stable and organs fed even as blood vessels start to leak and widen.'}
2. {'what': 'Breathing rate increases (tachypnea—usually over 20 breaths per minute)', 'why': 'Your lungs breathe faster to bring in more oxygen and blow off carbon dioxide. This also helps compensate for lactic acid (a waste product) building up in tissues that are not getting enough oxygen (anaerobic metabolism), keeping blood pH from dropping (metabolic too much acid in the blood (acidosis)).'}

3. {'what': 'Blood vessels in the skin and gut constrict (vasoconstriction)', 'why': 'Your body sacrifices blood flow to the skin (you look pale or mottled) and intestines to shunt blood to the brain and heart—your most vital organs. This keeps your blood pressure stable temporarily, even as infection spreads.'}
4. {'what': 'This compensation can mask early sepsis', 'why': 'A single snapshot of vital signs may look stable (blood pressure 110/70, alert and talking), hiding the fact that the body is working overtime to stay stable. Once compensation fails, the patient crashes suddenly—blood pressure drops, confusion sets in, urine output stops—because there is no reserve left.'}
5. {'what': 'Watch for the tipping point', 'why': 'Persistent low blood pressure despite IV fluids (septic shock, defined as systolic BP under 90 or MAP under 65 despite 30 mL/kg fluid bolus), confusion or altered mental status (brain not getting enough oxygen), low urine output (under 0.5 mL/kg/hour, meaning kidneys are failing), or rising lactate (over 2 mmol/L, a blood test showing cells are starving for oxygen) signal decompensation—the body can no longer compensate, and multi-organ failure is beginning.'}
6. {'what': 'Trend vital signs over time, not just one reading', 'why': 'A heart rate climbing from 80 to 110 over six hours, even if blood pressure stays normal, tells you the body is compensating harder—infection is advancing. Stable now does not mean safe. Serial measurements (every 1–4 hours) reveal the trajectory and catch deterioration early.'}

Domino effect

1. Bacteria enter the bloodstream through the central line — Skin bacteria (commonly *Staphylococcus epidermidis*, a germ that lives harmlessly on everyone's skin, or *Staphylococcus aureus*, a more aggressive cousin) migrate along the catheter tract from the insertion site, or hands contaminate the catheter hub during access. Either way, bacteria break through the skin barrier into the sterile bloodstream—a place they should never be.
2. Bacteremia (bacteria in the blood) triggers a whole-body alarm — Bacterial cell walls contain toxins (endotoxins from gram-negative bacteria like *E. coli*, or peptidoglycan and lipoteichoic acid from gram-positive bacteria like Staph). These toxins activate immune cells (macrophages, neutrophils), which release chemical messengers (cytokines: $\text{TNF-}\alpha$, IL-1, IL-6) that sound the alarm throughout the body, causing fever, chills, and widespread inflammation.
3. Fever and rigors (violent shaking chills) appear — Cytokines travel to the brain and reset the body's thermostat (hypothalamus) to a higher temperature (fever, often 101–104°F). Your muscles shiver violently (rigors) to generate heat and reach that new set-point. This is your body trying to slow bacterial growth—most bacteria grow best at 98.6°F, so raising the temperature gives your immune system an edge.

4. A fast heart rate (Tachycardia) (fast heart rate) and fast breathing (tachypnea) (fast breathing) develop — Your heart races (often 100–140 beats per minute) to pump more blood, delivering oxygen and immune cells to infected tissues. Your lungs breathe faster (often 24–30 breaths per minute) to bring in oxygen and blow off carbon dioxide and lactic acid (a waste product made by oxygen-starved cells doing anaerobic metabolism). This keeps you stable temporarily.
5. Sepsis emerges if bacteria overwhelm your defenses — Uncontrolled bacterial multiplication and toxin release cause widespread vasodilation (blood vessels relax and widen, dropping blood pressure), capillary leak (fluid leaks out of blood vessels into tissues, causing swelling and dropping blood volume), and organ dysfunction from poor blood flow (hypoperfusion—organs starve for oxygen). This is sepsis: life-threatening organ dysfunction caused by the body's dysregulated (out-of-control) response to infection.
6. Septic shock occurs if blood pressure drops despite IV fluids — Massive vasodilation and capillary leak drop blood pressure so low (systolic under 90 or MAP under 65 mmHg) that organs—brain, kidneys, liver—do not get enough oxygen. Cells switch to anaerobic metabolism (burning fuel without oxygen), producing lactic acid (lactate over 2 mmol/L). You become confused (brain poor blood flow (hypoperfusion)), stop making urine (kidney poor blood flow), and your skin turns mottled or cold (peripheral shutdown). Without immediate treatment (fluids, antibiotics, pressors to squeeze blood vessels), multi-organ failure and death follow within hours.
7. Removing the line and starting antibiotics stops the cycle — Removing the catheter eliminates the biofilm reservoir—bacteria hiding in the slimy coating on the line surface that antibiotics cannot reach. Without the biofilm factory, the bacteria source is gone. Broad-spectrum IV antibiotics (such as vancomycin to cover Staph plus a gram-negative antibiotic like cefepime or piperacillin-tazobactam) kill circulating bacteria, stopping toxin release. Once the source is gone and antibiotics clear the bloodstream, your immune system regains control, inflammation calms, blood pressure stabilizes, and organs recover.
8. Blood cultures identify the germ and guide antibiotic choice — Before starting antibiotics, nurses draw blood cultures from the central line and a peripheral vein. If the same bacteria grow from both sites, and the central-line culture turns positive at least two hours earlier than the peripheral culture (differential time to positivity), the line is confirmed guilty. Culture results (usually available in 24–72 hours) identify the exact bacteria and tell doctors which antibiotics will kill it, allowing de-escalation from broad-spectrum to targeted therapy (antibiotic stewardship) and reducing side effects and resistance.

Symptoms · what the patient feels, and why

- **Fever** (temperature $\geq 38.0^{\circ}\text{C}$ or $\geq 100.4^{\circ}\text{F}$): Bacteria in the blood release toxins that tell the hypothalamus (the brain's thermostat) to turn up body heat. The higher temperature slows down bacteria and speeds up immune cells—like turning up the oven to kill germs.
- **Rigors** (severe shaking chills): Your muscles shake hard to make extra heat and match the new fever set-point. Rigors mean a big wave of bacteria just dumped into your bloodstream—your body feels the invasion and reacts fast.
- **A fast heart rate (Tachycardia)** (heart rate >90 beats per minute): The heart pumps faster to rush more blood, oxygen, and immune soldiers to the tissues under attack. It is like a fire engine speeding to a burning building.
- **Low blood pressure (Hypotension)** (systolic blood pressure <90 mmHg or drop ≥ 40 mmHg from baseline): Bacterial toxins force blood vessels to relax and widen, and fluid leaks out of the vessels into the tissues. Pressure inside the pipes (vessels) drops dangerously low. When IV fluids cannot bring it back up, that is septic shock—a life-threatening emergency.
- **Altered mental status** (confusion, drowsiness, agitation): Low blood pressure means not enough blood reaches the brain. Without enough oxygen and sugar (glucose), brain cells cannot work right—thinking gets foggy, and the person may seem confused or sleepy.
- **Redness (Erythema)** (redness), warmth, or pus at catheter insertion site: Bacteria growing on the catheter's outside surface infect the skin and tunnel where the line enters. The body sends extra blood (making it red and warm) and white blood cells (which form pus) to fight the invaders at the door.
- **Low urine output (Oliguria)** (urine output <0.5 mL/kg/hour): Low blood pressure means the kidneys get less blood flow. To protect the body, the kidneys hold back water and make very little urine—like a dam closing when the river is too low. It is a red flag that the kidneys are starting to fail.
- **Fast breathing (Tachypnea)** (respiratory rate >20 breaths per minute): Tissues starved of oxygen start making lactic acid (a waste product). The lungs breathe faster to blow off carbon dioxide and balance the acid in the blood—like opening more windows to clear smoke from a room.
- **Hypothermia** (temperature $<36.0^{\circ}\text{C}$ or $<96.8^{\circ}\text{F}$): In severe sepsis, the hypothalamus (the brain's thermostat) can break down completely and temperature drops instead of rising. This is a very bad sign—it means the immune system is overwhelmed and losing control.
- **Thrombophlebitis** (hard, cord-like vein, tender to touch): Bacteria coat the inside of the catheter and trigger a blood clot (thrombus) mixed with inflammation in the vein. The vein feels like a tight rope under the skin because it is swollen, clotted, and infected.

Focused exam

- Fever that starts during or right after using the central line: Fever tied to line use means bacteria just entered the bloodstream from the line itself. The body's alarm goes off because infection is happening in real time.
- Blood drawn from the line grows bacteria at least 2 hours before blood from the arm does: This time difference proves the line is the source, not some other infection site. It is the smoking gun for CLABSI.
- Redness, warmth, swelling, or pus where the line enters the skin: The line punctures skin, and that entry point is where bacteria sneak in. Redness and warmth mean the body is fighting infection right where the skin was broken.
- Fast heart rate, low blood pressure, or shaking chills: These signs mean bacteria are flooding the bloodstream and the body is struggling to fight back. Sepsis—life-threatening organ trouble caused by infection—is starting, and the line is the cause.
- No other clear source of fever after checking lungs, urine, belly, and surgical sites: When fever has no other identifiable source, the central line becomes the prime suspect for CLABSI.
- White blood cell count too high or very low: The immune system is either sending lots of white blood cells to fight infection or becoming overwhelmed and running out because bacteria from the line are winning.

Diagnostics · every test answers a question

- Paired blood cultures: one set drawn from the central line and one set drawn from a vein in the arm at the same time → Is the central line the source of the bloodstream infection?
- Complete blood count (CBC) with differential → Is the immune system responding well to infection, or is it overwhelmed?
- Lactate level → Are the tissues getting enough oxygen, or is sepsis causing blood flow to be too low?
- Procalcitonin (optional helper test) → Is this infection bacterial, and how severe is it?
- C-reactive protein (CRP) → Is there inflammation in the body, and how strong is it?
- Basic metabolic panel (BMP) or comprehensive metabolic panel (CMP) → Are the kidneys, liver, and electrolytes being hurt by sepsis?
- Chest X-ray → Is there pneumonia or another lung infection causing the fever instead of the line?
- Urinalysis and urine culture → Is a urinary tract infection causing the fever instead of the line?

Treatment logic

- Draw paired blood cultures—one set from the central line and one set from a vein in the arm—before starting any antibiotics → If line blood grows bacteria at least 2 hours faster than arm blood, that proves the line itself is the source. This tells you whether to remove the line or try to save it with antibiotics.
- Start IV antibiotics immediately: vancomycin (covers MRSA and coagulase-negative staph from skin) plus an anti-pseudomonal beta-lactam like cefepime, piperacillin-tazobactam, or meropenem → These antibiotics kill the most common CLABSI bacteria—skin organisms like *Staphylococcus aureus* and coagulase-negative staph, plus gram-negative rods like *Pseudomonas*. They stop bacteria from multiplying in the blood and worsening sepsis.
- Remove the central line promptly if bacteria are *S. aureus*, *Candida* (fungus), gram-negative rods like *Pseudomonas*, or if fever and positive cultures persist 72 hours after starting antibiotics → Taking out the line removes the source where bacteria are hiding in biofilm and reproducing. This gives antibiotics a real chance to clear the bloodstream infection and prevents complications like endocarditis—infection of the heart valves—or septic thrombophlebitis, an infected clot in the vein.
- Repeat blood cultures 2 to 4 days after starting antibiotics to confirm bacteremia—bacteria in the blood—is clearing → Negative repeat cultures show the antibiotics and line removal (if done) actually worked. You know the infection is clearing and not turning into complicated bacteremia, septic shock, or endocarditis.
- Transthoracic or transesophageal echocardiogram—heart ultrasound—if *S. aureus* bacteremia is confirmed → This looks for vegetations—bacterial clumps stuck to heart valves causing endocarditis—which is a serious complication requiring 4 to 6 weeks of IV antibiotics instead of 2 weeks, and sometimes valve surgery.
- Antibiotic lock therapy—high-concentration antibiotic solution left sitting in the catheter between uses—only if bacteria are coagulase-negative staph, the line is essential and cannot be removed, and bloodstream infection is uncomplicated → High-dose antibiotic liquid sitting in the line for hours kills bacteria stuck in biofilm on the catheter inner surface. This allows you to save the line without removal.
- Give IV fluids—crystalloid like normal saline or lactated Ringer's—if blood pressure is low or lactate is elevated → Fluids expand blood volume and improve blood flow to organs, reversing tissue poor blood flow (hypoperfusion)—not enough blood reaching the tissues. This can bring lactate down and stabilize blood pressure.

Medications

Vancomycin plus a broad-spectrum beta-lactam (such as cefepime or piperacillin-tazobactam) · Glycopeptide antibiotic (vancomycin) combined with beta-lactam antibiotic

Vancomycin sticks to the building blocks of bacterial cell walls in Gram-positive bacteria (the bacteria that turn purple under a microscope, like Staph and Strep). It blocks the bacteria from linking these blocks together, so the wall gets weak and the cell pops like a balloon. The beta-lactam drug sticks to the construction enzymes (called penicillin-binding proteins, or PBPs) that Gram-negative bacteria (bacteria that turn pink, like *E. coli* and *Pseudomonas*) and some Gram-positive bacteria use to build their walls. When the enzyme is blocked, the bacteria cannot finish building their wall, and they die. Together, these two drugs kill the most common bacteria that sneak into the bloodstream through a central line—Staph from the skin, Strep, *Enterococcus*, *Pseudomonas*, and gut bacteria like *E. coli* and *Klebsiella*.

1. Bacteria living on the skin (especially Staph epidermidis and Staph aureus) or germs that land on the catheter hub crawl into the inside tube of the catheter (the lumen) and form a sticky layer called biofilm—a slime shield that protects them.
2. Some bacteria break off from the biofilm and float into the bloodstream, where the body's alarm system fires off: fever (temperature above 100.4°F or 38°C), fast heartbeat (over 90 beats per minute), fast breathing (over 20 breaths per minute), and a surge in white blood cells (over 12,000 per microliter) or a big drop (under 4,000 per microliter). This whole response is called systemic inflammatory response.
3. Vancomycin spreads into the bloodstream and tissues. It finds the D-Ala-D-Ala tails on the growing peptidoglycan chains and binds to them, blocking the final cross-link step. Without cross-links, the cell wall is too weak to hold the cell together, and it bursts (lyses).
4. The beta-lactam spreads into the bloodstream and tissues, finds the PBP enzymes, and locks onto them. The bacteria cannot finish making peptidoglycan, so the wall stays incomplete and the cell dies.
5. After 24 to 48 hours, the lab reports which bacteria grew in the blood cultures and which antibiotics kill them (called susceptibility results). The doctor then switches to the narrowest antibiotic that still kills the bacteria (called de-escalation), which lowers the risk of side effects and resistance.
6. The catheter must be removed right away if the blood culture grows Staph aureus, Candida (a fungus), Gram-negative rods like *Pseudomonas*, or if bacteria keep growing in the blood after 72 hours of antibiotics. It also must come out if the patient has severe sepsis (organs start to fail—kidneys, lungs, or liver stop working well) or if a clot in the vein fills with pus (called suppurative thrombophlebitis).

7. If the catheter absolutely must stay in place and the bacteria is coagulase-negative Staph (a milder Staph cousin) or Enterococcus, and there are no complications, the doctor may use antibiotic lock therapy: a super-high dose of antibiotic is squirted into the catheter and left there for 8 to 12 hours while the line is not being used, to kill bacteria hiding in the biofilm.
8. Two to four days after starting antibiotics, the doctor draws blood cultures again to make sure no bacteria are growing anymore (called clearance).
9. If Staph aureus is confirmed in the blood, an ultrasound of the heart (called a transthoracic echocardiogram, or TTE) is done to check for endocarditis—an infection of the heart valves—because Staph aureus loves to stick to heart valves and cause serious damage.

SAFETY WATCH

Vancomycin: The doctor checks a blood level (called a trough, drawn right before the next dose) to keep it between 15 and 20 mg/L for serious infections, following IDSA (Infectious Diseases Society of America) guidelines. If the level gets too high or if vancomycin is combined with other drugs that hurt the kidneys (like aminoglycosides or NSAIDs such as ibuprofen), it can damage the kidneys (called nephrotoxicity) or cause hearing loss (called ototoxicity). · Beta-lactams: Some people are allergic. In rare cases, a severe allergic reaction (called anaphylaxis) can happen—throat swelling, trouble breathing, and dangerously low blood pressure. If the patient has reduced kidney function (creatinine clearance below 50 mL per minute), the dose must be lowered to avoid toxic buildup. · The catheter must be pulled out promptly when guidelines say so. Waiting too long raises the risk that bacteria will travel to the heart valves, bones, or other organs (called metastatic infection), cause septic shock (blood pressure dropping below 90 mmHg systolic even after giving fluids), or lead to death.

Red flags

- Low blood pressure (Hypotension) (systolic BP <90 mmHg) despite giving IV fluids—sign of septic shock
- Altered mental status (confusion, extreme sleepiness, or no response)—brain is not getting enough blood
- Low urine output (Oliguria) or no urine output (anuria) (urine output <0.5 mL/kg/hour or zero urine)—kidneys are failing
- Positive blood cultures still growing bacteria 72 hours or more after removing the line and starting antibiotics—may mean endocarditis (infection stuck to a heart valve) or a hidden abscess
- New heart murmur or embolic events (stroke, dead fingers or toes from clots)—bacteria may have seeded the heart valves (endocarditis)

Nursing priorities

1. Obtain two sets of blood cultures—one set drawn from the central line and one set from a peripheral vein in the arm—before any antibiotics are given. — Paired cultures (central and peripheral) help prove the central line is infected; if the central-line culture grows bacteria at least 2 hours before the arm culture, the line is the source. Drawing before antibiotics keeps bacteria alive so the lab can identify the exact germ and test which antibiotics kill it.
2. Start IV antibiotics within 1 hour: vancomycin to cover MRSA (methicillin-resistant *Staph aureus*) and coagulase-negative staph, plus an anti-pseudomonal beta-lactam like cefepime, piperacillin-tazobactam, or meropenem to cover gram-negative bacteria and *Pseudomonas*. — Every hour delay in antibiotics increases mortality (death rate) in sepsis by about 7%; vancomycin covers skin bacteria that enter through the catheter, and the beta-lactam covers gut and water bacteria that cause hospital infections.
3. Assess blood pressure, heart rate, respiratory rate, temperature, oxygen saturation, and mental status every 15 to 30 minutes until stable, and notify the physician immediately if systolic BP drops below 90 mmHg or patient becomes confused. — Septic shock develops when bacteria release toxins that dilate blood vessels, dropping blood pressure and starving organs of oxygen; early recognition lets the team give IV fluids (30 mL per kilogram bolus) and start vasopressors like norepinephrine before organs fail.
4. Measure urine output every hour using a Foley catheter (bladder tube) and report if output falls below 0.5 mL per kilogram per hour. — The kidneys receive 20% of the heart's blood with each beat; when blood pressure drops in sepsis, kidneys shut down to preserve flow to the brain and heart. Low urine output (Oliguria) is an early warning of acute kidney injury and signals the need for more IV fluids or medicines to raise blood pressure.
5. Prepare for central line removal if blood cultures grow *S. aureus*, *Candida* (yeast), or if the patient has septic shock, tunnel infection, or port-pocket infection; coordinate with the physician and have a peripheral IV ready before the line comes out. — *S. aureus* and *Candida* stick to the catheter in a biofilm (a slime layer antibiotics cannot penetrate), so the line must be removed or bacteria keep seeding into the blood, causing endocarditis or septic traveling clots (emboli) (clots of bacteria traveling to lungs, brain, or joints). Removing the source controls the infection.
6. Give IV fluid boluses as ordered—typically 30 mL per kilogram of crystalloid (normal saline or lactated Ringer's) over the first 3 hours—and reassess blood pressure, heart rate, and lung sounds after each liter. — Sepsis causes capillary leak (fluid leaking out of blood vessels into tissues), so the blood volume drops; rapid IV fluids refill the vessels, improve blood pressure, and restore oxygen delivery to organs. Listening to lungs prevents fluid overload in patients with heart failure.

7. Inspect the central line exit site or port pocket daily for redness (erythema) (redness), warmth, tenderness, swelling, or pus-filled (purulent) drainage, and document findings with measurements and photos if available. — Exit-site or tunnel infection (redness and pus tracking along the catheter under the skin) requires line removal even if blood cultures are negative; early detection prevents the infection from spreading into the bloodstream or forming an abscess.
8. Monitor blood-culture results daily and notify the physician immediately if cultures remain positive after 72 hours of appropriate antibiotics and line removal. — Persistent bacteremia (bacteria still growing in blood after 3 days of treatment and source removal) suggests endocarditis (infection stuck to a heart valve), septic thrombophlebitis (infected clot in the vein), or a deep abscess; the patient needs an echocardiogram and longer antibiotic course, often 4 to 6 weeks.

Sources

- IDSA Clinical Practice Guidelines for the Diagnosis and Management of Intravascular Catheter-Related Infection, 2009 (updated 2021)
- CDC Guidelines for the Prevention of Intravascular Catheter-Related Infections, 2011 (updated 2017)
- Surviving Sepsis Campaign International Guidelines for Management of Sepsis and Septic Shock, 2021
- Cochrane Database of Systematic Reviews

Last reviewed: 2026-07-07