

Ccb Overdose

A calcium channel blocker overdose jams the gates that normally let calcium rush into heart muscle, blood vessel muscle, and insulin-making cells, so the heart beats slowly and weakly, blood pressure drops, and blood sugar climbs.

THE STORY IN ONE SENTENCE

A calcium channel blocker overdose jams the gates that normally let calcium rush into heart muscle, blood vessel muscle, and insulin-making cells, so the heart beats slowly and weakly, blood pressure drops, and blood sugar climbs.

“CCB overdose blocks calcium from entering cells, causing a slow weak heart and low blood pressure; we push calcium and high-dose insulin straight into the vein to restart the pump.”

Normal anatomy — the body's job

L-type calcium channels are tiny gates in the walls of heart muscle cells, blood vessel muscle cells, and pancreas insulin cells. When a signal arrives, the gate swings open and calcium rushes in. In the heart, that calcium flood makes the muscle squeeze and passes the electrical beat from the top chambers to the bottom chambers. In blood vessels, calcium makes the muscle tighten to hold pressure steady. In the pancreas, calcium tells the cell to release insulin so sugar can leave the blood and feed your body.

BODY'S JOB

The heart needs calcium rushing in every single beat to squeeze hard enough to push blood out. Blood vessels need calcium to stay firm so pressure doesn't drop. The pancreas needs calcium to know when to release insulin and keep blood sugar normal.

1. An electrical signal arrives at a heart muscle cell and opens the L-type calcium channel gate.
2. Calcium floods into the cell like water through a floodgate.
3. That calcium wave grabs tiny gears inside the muscle fiber and makes them slide past each other—the cell squeezes.
4. At the same time, calcium spreads the electrical signal from the upper heart chambers (atria) to the lower chambers (ventricles) so the beat stays coordinated.

5. In blood vessel muscle, calcium entry makes the vessel tighten just enough to keep blood pressure steady as blood pushes through.
6. In the pancreas, calcium entering tells the cell to release insulin, which unlocks doors on other cells so sugar can leave the blood and go inside to be burned for energy.

What breaks

A calcium channel blocker overdose jams the L-type calcium gates shut all over the body at once. No calcium can rush in, so heart muscle can't squeeze well, the electrical signal between heart chambers slows or stops, blood vessels go limp, and the pancreas stops releasing insulin.

CLINICAL TERM

Overdose of a calcium channel blocker (e.g., amlodipine, diltiazem, verapamil) causes excessive blockade of L-type calcium channels in cardiac myocytes, vascular smooth muscle, and pancreatic beta cells, leading to negative inotropy (weak squeeze), negative chronotropy and dromotropy (slow rate and slow conduction through the AV node), vasodilation (low systemic vascular resistance), and impaired insulin secretion (hyperglycemia).

How the body compensates

The body senses low blood pressure and poor oxygen delivery, so it releases adrenaline and other stress hormones to try to speed the heart and tighten vessels—but the calcium channels are jammed, so these signals can't work. The brain dials itself down to save energy, causing confusion or sleepiness. Muscles switch to backup fuel that makes lactic acid. These responses buy minutes to hours, but they can't fix the jammed gates—so vital signs may look okay at first, then crash suddenly when compensation maxes out.

1. {'what': 'The body releases adrenaline (epinephrine) and norepinephrine from the adrenal glands when pressure sensors detect low blood pressure.', 'why': 'These stress hormones normally tell the heart to beat faster and vessels to tighten—but because calcium channels are blocked, the signal can't get through, so this compensation fails and the patient stays slow and with low blood pressure (hypotensive).'} }
2. {'what': 'The brain reduces its own activity to save oxygen, causing confusion, sleepiness, or even loss of consciousness (altered mental status).', 'why': 'The brain is protecting itself from damage by slowing down when it senses it's not getting enough oxygen-rich blood.'} }

3. {'what': "Muscles that aren't getting enough oxygen switch from normal fuel-burning (aerobic) to a backup system (anaerobic) that makes lactic acid as waste.", 'why': 'This backup system keeps muscles alive for a short time, but lactic acid builds up in the blood (lactic too much acid in the blood (acidosis)) and makes the blood too acidic, which can stop enzymes and the heart from working.'}
4. {'what': 'The kidneys sense low pressure and release renin to try to raise blood pressure and hold onto water.', 'why': "This is a slower backup system that takes hours, and it can't overcome the jammed calcium channels, so it doesn't help in acute overdose."}
5. {'what': 'The patient may look stable for the first few hours, especially with immediate-release pills, then suddenly crash when compensation maxes out or when extended-release pills reach peak levels 6–12 hours later.', 'why': "The body's backup systems can only hold vital signs steady for so long; once they fail or the drug level climbs higher, the heart and vessels give out all at once—this is the dangerous cliff edge to watch for."}

Domino effect

1. Calcium channel blocker floods the bloodstream and binds to L-type calcium channels in the heart, blood vessels, and pancreas, jamming the gates shut. — The drug locks onto the channel like a key stuck in a lock, so calcium can't rush in even when the cell gets a signal to open the gate.
2. Heart muscle can't squeeze hard because no calcium is entering to trigger contraction—the heart beats weakly and slowly (negative inotropy and chronotropy). — Calcium is the switch that makes the muscle fibers slide and squeeze; without it, the heart pumps like a weak battery, and the rate drops because the natural pacemaker cells also depend on calcium to fire.
3. The electrical signal from the top chambers (atria) to the bottom chambers (ventricles) slows or stops at the AV node, causing a slow heart rate (bradycardia) or complete heart block. — The AV node uses calcium channels to pass the signal; when they're jammed, the signal crawls or stops entirely, so the ventricles may not get the command to beat.
4. Blood vessels go limp because their muscle has no calcium to tighten, so blood pressure drops (vasodilation and distributive shock). — Vascular smooth muscle needs calcium to contract and keep vessels firm; without it, vessels relax, the pipes widen, and pressure falls like air leaking from a tire.
5. The pancreas stops releasing insulin because calcium is the trigger, so blood sugar climbs (hyperglycemia) even though cells are starving for fuel. — Beta cells in the pancreas use calcium entry as the 'go' signal to release insulin; when the gate is jammed, no insulin comes out, so sugar piles up in the blood and can't enter cells to be burned.

6. Organs don't get enough oxygen-rich blood because the heart is weak and pressure is low, so muscles make lactic acid (lactic too much acid in the blood (acidosis)) and the brain shuts down to save energy (altered mental status). — When tissues can't get enough oxygen from blood, they switch to anaerobic metabolism that produces lactic acid as waste; the brain senses danger and reduces its activity, causing confusion or sleepiness.
7. The body's stress response (adrenaline, norepinephrine) tries to rescue blood pressure and heart rate but fails because calcium channels are blocked, so the patient stays bradycardic and with low blood pressure (hypotensive) despite high stress hormone levels. — Adrenaline and norepinephrine work by opening calcium channels and other pathways; when calcium channels are jammed by the overdose, this rescue system can't work, and compensation fails.
8. Without treatment, the heart may stop pumping enough blood to keep the brain and organs alive (cardiogenic shock), the rhythm may degrade to complete standstill (asystole), or the patient may die from multi-organ failure. — The heart and vessels can't function without calcium entry; if the blockade isn't reversed with calcium infusion, high-dose insulin, and supportive care, the cardiovascular system collapses completely.

Symptoms · what the patient feels, and why

- Slow heart rate (bradycardia): Calcium channel blockers stop calcium from entering the heart's pacemaker cells, so the heart's natural timer slows down and cannot fire as fast.
- Low blood pressure (hypotension): The medicine relaxes the smooth muscle in blood vessel walls, so the vessels widen and pressure drops; also, the heart pumps less powerfully because calcium is needed for muscle contraction.
- Dizziness or feeling faint: When blood pressure drops and the brain gets less oxygen-rich blood, you feel lightheaded—like standing up too fast.
- Nausea or vomiting: Poor blood flow to the gut and a general stress response from low blood pressure can upset the stomach.
- Confusion or altered mental state: The brain is very sensitive to low oxygen; when blood pressure is too low, not enough oxygen reaches brain cells and thinking becomes muddled.
- Chest pain or shortness of breath: The heart muscle itself may not get enough blood when blood pressure is very low, causing pain; the lungs may fill with fluid if the heart cannot pump effectively (cardiogenic shock).
- High blood sugar (hyperglycemia): Calcium is needed for insulin to be released from the pancreas; blocking calcium prevents insulin release, so sugar builds up in the blood.
- Delayed symptom onset with extended-release pills: Extended-release formulations dissolve slowly in the gut, so the drug enters the bloodstream gradually over many hours—symptoms may not appear until 12 to 24 hours after swallowing.

Focused exam

- Heart rate slower than 60 beats per minute (bradycardia): Calcium-channel blockers (CCBs) stop calcium from entering the heart's natural pacemaker cells. Without calcium, the electrical signals that tell the heart to beat fire much more slowly. When you see a slow heart rate together with low blood pressure, think CCB overdose first.
- Blood pressure lower than 90 systolic (hypotension): CCBs block calcium channels in the smooth muscle that wraps around blood vessels. Without calcium, that muscle cannot squeeze. The vessels stay relaxed and pressure drops. Slow heart plus low pressure together is the signature of CCB overdose.
- Confusion, sleepiness, or trouble waking up (altered mental status): When blood pressure falls, less oxygen-rich blood reaches the brain. Brain cells slow down to save energy. This tells you the overdose is severe enough to hurt blood flow (perfusion) (blood flow to organs).
- Blood sugar higher than 200 mg/dL without a history of diabetes (hyperglycemia): CCBs block calcium channels in the pancreas cells that release insulin. No calcium in means no insulin out, so sugar piles up in the blood. This triad—slow heart, low pressure, high sugar—points directly to CCB overdose and almost nothing else.
- Wet, crackling lung sounds (rales) or pink frothy coughed-up mucus (sputum): When the heart is too weak to pump blood forward, blood backs up into the lungs. Fluid leaks into the air sacs (pulmonary swelling (edema)). The patient may feel short of breath and cough up foam.
- Delayed symptom onset—patient may look fine for 2 to 6 hours after swallowing a long-acting CCB pill: Extended-release CCB tablets (like amlodipine, diltiazem ER, or verapamil SR) release drug slowly. The patient can seem stable at first, then crash suddenly hours later. Never let the clock fool you; if they took a long-acting pill, watch closely for at least 12 to 24 hours.

Diagnostics · every test answers a question

- 12-lead electrocardiogram (ECG) → Is the heart beating slowly, and are there blocks in the electrical system?
- Fingerstick or serum blood glucose → Is the blood sugar high without a history of diabetes?
- Serum lactate → Are the tissues getting enough oxygen, or is the body in shock?
- Basic metabolic panel (BMP) including potassium → Are the kidneys still working, and is potassium dangerously high or low?
- Arterial or venous blood gas → Is the blood too acidic, and is there enough oxygen?
- Serum troponin I or T → Is the heart muscle being damaged?

- Chest X-ray → Is there fluid backing up into the lungs?
- Urine or serum drug screen (if ingestion history is unclear) → Did the patient take other drugs that might explain the symptoms or make treatment harder?

Treatment logic

- Intravenous (IV) calcium: calcium chloride 10–20 mL of 10% solution (1–2 grams) through a central line, or calcium gluconate 30–60 mL of 10% solution (3–6 grams) through a peripheral IV, given over 5–10 minutes and repeated every 15–20 minutes as needed → Calcium floods the space around heart and blood-vessel muscle cells and pushes through the few channels that are not blocked, helping the heart squeeze and the vessels tighten. This directly replaces what the CCB stole. Blood pressure may rise and heart rate may climb within minutes.
- High-dose insulin with dextrose: regular insulin 1 unit/kg IV bolus, then 0.5–1 unit/kg/hour IV infusion, PLUS dextrose 25–50 grams IV bolus followed by dextrose 10% or 50% infusion titrated to keep blood glucose 100–250 mg/dL, with glucose checks every 15–30 minutes → High-dose insulin forces glucose into heart muscle cells to fuel them, making the heart squeeze harder (positive inotropy). It also helps blood vessels tighten. Insulin works even when calcium channels are blocked because it uses a different signal pathway. This is the most important treatment after calcium in severe CCB overdose.
- Vasopressor infusion: norepinephrine (start 0.05–0.1 mcg/kg/min) or epinephrine (start 0.05–0.2 mcg/kg/min), titrated every few minutes to keep systolic blood pressure above 90 mmHg or mean arterial pressure (MAP) above 65 mmHg → Vasopressors force blood vessels to squeeze by activating a different receptor pathway (alpha-adrenergic receptors) that does not need calcium channels. This raises blood pressure quickly and keeps blood flowing to the brain, heart, and kidneys while calcium and insulin start working.
- Atropine 0.5–1 mg IV push, repeated every 3–5 minutes up to a total dose of 3 mg, and/or transcutaneous pacing (external pads) or transvenous pacing (wire threaded into the heart) if atropine does not work → Atropine blocks the vagus nerve signal that slows the heart, allowing the heart to beat faster. Pacing delivers small electrical shocks to force the heart to beat at a safe rate (usually 60–80 beats per minute) when drugs fail.
- Glucagon 3–5 mg IV bolus over 1 minute, then 1–5 mg/hour continuous infusion if the bolus helps → Glucagon activates a backup signal pathway (cyclic AMP) inside heart cells that makes them squeeze harder and beat faster without needing calcium channels. This can work when calcium alone does not, especially in patients on beta-blockers at the same time.

- Intravenous lipid emulsion (ILE): 20% lipid solution 1.5 mL/kg (roughly 100 mL for a 70-kg adult) IV bolus over 1 minute, then 0.25 mL/kg/min (roughly 250–500 mL/hour) continuous infusion for 30–60 minutes, repeated once if needed → Lipid emulsion creates tiny fat droplets in the blood that soak up (absorb) the CCB molecules like a sponge. This pulls the drug away from the heart and blood vessels and traps it in the bloodstream, freeing up the blocked channels. Blood pressure and heart rate may improve over 10–30 minutes.
- Extracorporeal membrane oxygenation (ECMO) or cardiopulmonary bypass in refractory shock or cardiac arrest → ECMO is a machine that pumps and oxygenates the blood outside the body, completely bypassing the failing heart and lungs. This keeps the patient alive for hours to days while the CCB is metabolized and cleared from the body.

Medications

Calcium chloride or calcium gluconate (IV) · Calcium salt

Calcium chloride blocks the heart's muscle from squeezing well and slows electrical signals. We flood the bloodstream with calcium to compete directly with the drug sitting in the heart's doorways (calcium channels) and push the blocker out of the way. The extra calcium restores the muscle's ability to tighten and the heart's rhythm centers to fire.

1. CCB molecules plug the tiny gates (L-type calcium channels) on heart cells so calcium cannot enter.
2. Without calcium, the heart muscle cannot squeeze and the electrical relay between chambers slows or stops.
3. We give a high dose of calcium IV so the blood concentration rises sharply.
4. The surge of calcium pushes CCB molecules off the gates by pure competition—like turning up the volume so you can hear over noise.
5. Calcium rushes back into cells; the squeeze returns and electrical signals speed up again.

SAFETY WATCH

Calcium chloride burns tissue badly if it leaks outside the vein; use a large, secure IV line.

- Can cause dangerous heart rhythms if pushed too fast; give slowly over 5–10 minutes.
- Do NOT mix calcium with bicarbonate in the same IV line—it will form chalk and clog the line.
- Watch the heart monitor continuously during the infusion.

High-dose insulin with dextrose (IV) · Hormone (pancreatic peptide)

CCB overdose shuts down the pancreas so it stops making insulin; heart muscle then cannot pull in sugar or burn it for energy, so the pump weakens even if calcium is restored. We give huge doses of insulin—far more than a diabetic would ever get—to force the heart muscle to

take in sugar and make ATP (the cell's energy currency). Dextrose (sugar water) is given at the same time to keep blood sugar from dropping dangerously low.

1. CCBs block calcium entry into pancreas cells, stopping insulin release.
2. Heart muscle runs out of fuel (ATP) because it cannot move sugar from the blood into the cell without insulin.
3. We give 1 unit of insulin per kilogram as a bolus, then 0.5–1 unit/kg every hour as a drip.
4. Insulin unlocks the doors on heart cells so glucose floods in.
5. The cell burns glucose to make ATP; the heart regains strength to squeeze.
6. We infuse 10 % dextrose (or higher concentration if needed) to replace the sugar insulin pushes out of the blood and prevent low blood sugar (hypoglycemia).

SAFETY WATCH

Check blood sugar every 20–30 minutes; if it drops below 100–150 mg/dL increase the dextrose drip rate. · Watch potassium closely—insulin drives potassium into cells and can cause dangerous low potassium (hypokalemia) within hours. · Replace potassium IV if the level falls below 3.5 mEq/L. · Requires close ICU monitoring with continuous glucose and cardiac telemetry.

Norepinephrine or epinephrine (IV) · Vasopressor (catecholamine)

CCBs relax blood vessels so they open wide and blood pressure falls. These drugs squeeze the vessels tight again by locking onto receptors (alpha-1) on the muscle in vessel walls. They also speed the heart (beta-1 effect) and help it squeeze harder, acting as a temporary bridge while calcium and insulin do their deeper work.

1. CCBs relax smooth muscle in blood vessels; pressure drops and organs do not get enough flow.
2. Norepinephrine or epinephrine is infused continuously into a large vein.
3. The drug binds to alpha-1 receptors on vessel walls, causing them to tighten.
4. Blood pressure rises and blood flow (perfusion) to the brain, kidneys, and heart improves.
5. Beta-1 stimulation speeds the heart rate and strengthens each beat.

SAFETY WATCH

Must go through a central line or very secure peripheral IV—if it leaks into tissue it kills the skin. · Raises heart workload and oxygen demand; can trigger chest pain or arrhythmias. · Monitor blood pressure continuously with an arterial line if possible. · Wean slowly once the patient stabilizes; sudden stop can cause pressure to crash again.

Glucagon (IV or IM) · Pancreatic hormone

Glucagon tells the heart to squeeze harder through a completely different pathway that does not need calcium channels. It binds to its own receptor on heart cells and activates a backup energy system (cyclic AMP) that forces the muscle to contract. Think of it as a side door when the main entrance is locked.

1. CCBs lock the calcium gates, so the usual squeeze signal is blocked.
2. Glucagon binds to glucagon receptors on the heart cell surface.
3. This turns on an enzyme (adenylyl cyclase) that makes cyclic AMP inside the cell.
4. Cyclic AMP tells stored calcium inside the cell to release, bypassing the blocked channels.
5. The heart muscle contracts more strongly even though the channels are still blocked.

SAFETY WATCH

Can cause severe nausea and vomiting; have suction ready and protect the airway. · May raise blood sugar, which is usually harmless in this setting. · Large doses (5–10 mg bolus, then 2–5 mg/hour infusion) are needed—much more than the 1 mg used for low blood sugar. · Effect may wear off in 10–20 minutes; be ready to re-dose or switch strategies.

Intravenous lipid emulsion (Intralipid 20%) · Fat emulsion (rescue therapy)

CCBs are fat-loving molecules. When we infuse a milky solution of tiny fat droplets, the drug molecules leave the heart tissue and dissolve into the fat floating in the blood—like oil soaking up a grease stain. This pulls the poison away from the heart and traps it harmlessly in the bloodstream until the body can clear it.

1. CCB sits in heart muscle, blocking channels and poisoning the pump.
2. We infuse 20 % lipid emulsion 1.5 mL/kg over 1 minute, then 0.25 mL/kg/min for 30–60 minutes.
3. The fat droplets circulate and CCB molecules jump from tissue into the lipid because they are more soluble there.
4. Drug concentration in the heart falls; toxicity eases.
5. The liver gradually breaks down the fat and the trapped drug inside it.

SAFETY WATCH

Can interfere with lab tests (makes blood samples look milky and unreadable) for hours. · Rarely causes fat overload syndrome (trouble breathing, confusion) if huge volumes are given. · May trigger pancreatitis in susceptible patients. · Use only when standard therapies are failing; it is not a first-line treatment.

Atropine (IV) · Anticholinergic (muscarinic antagonist)

CCBs can slow the heart's electrical pacemaker to dangerous levels. Atropine blocks the brake pedal (vagus nerve signals) that naturally slows the heart, allowing the pacemaker to speed up. It only helps the rhythm problem—it does nothing for the squeeze or blood pressure.

1. The vagus nerve releases acetylcholine, which slows the heart's pacemaker.
2. CCBs make this worse by blocking calcium entry needed for pacemaker cells to fire quickly.
3. Atropine blocks acetylcholine receptors on the pacemaker cells.
4. With the brake released, the heart rate rises.
5. Only effective if the pacemaker itself still has some function; if the CCB has completely shut it down, atropine will do nothing.

SAFETY WATCH

If it does not work after 1–2 doses (0.5–1 mg each), stop and move to pacing—repeated doses waste time. · Can cause confusion, dry mouth, blurred vision, and urinary retention (usually not a concern in acute overdose). · In complete heart block, atropine may speed the upper chambers but not the lower ones, worsening the mismatch.

Temporary cardiac pacing (transcutaneous or transvenous) · Device/procedure, not a drug

If the heart's natural pacemaker is poisoned and will not fire, we use an electrical device to send timed shocks through the chest wall or directly into the heart chamber. Each shock forces the muscle to contract, keeping the patient alive until the drug wears off or other treatments take effect.

1. The heart's own pacemaker is blocked and firing too slowly or not at all.
2. Transcutaneous pads are placed on the chest and back, or a pacing wire is threaded into the right ventricle.
3. The device sends electrical pulses at a set rate (e.g. 60–80 beats per minute).
4. Each pulse depolarizes the heart muscle, forcing a contraction.
5. The patient's circulation is maintained artificially until recovery.

SAFETY WATCH

Transcutaneous pacing is painful in an awake patient; give sedation and pain medicine. · Risk of skin burns at the pad sites with prolonged use. · Transvenous pacing requires a skilled operator and carries risk of heart perforation, infection, and bleeding. · Must have backup defibrillator and resuscitation drugs immediately available.

Extracorporeal membrane oxygenation (ECMO) · Device/procedure, not a drug

If the heart is so poisoned that it cannot pump blood at all despite every medication and pacing, ECMO is a machine that takes over completely. Blood is pulled out of the body, run through an artificial lung to add oxygen and remove carbon dioxide, then pumped back in. It buys time—sometimes days—for the drug to clear and the heart to recover.

1. The heart is failing completely; blood pressure and oxygen are critically low.
2. Large tubes (cannulas) are placed into a vein and an artery.
3. Blood is drained out, pumped through a machine that oxygenates it, and returned to the body.
4. Organs receive oxygen and nutrients even though the native heart is barely working.
5. ECMO runs for hours to days while the CCB is metabolized and excreted.

SAFETY WATCH

Risk of severe bleeding (blood must be thinned to prevent clotting in the circuit). · Infection at cannula sites or in the bloodstream. · Limb starved blood flow (ischemia) (loss of blood flow to the leg where the artery cannula is placed). · Stroke from clots or air bubbles. · Requires 24/7 monitoring by trained perfusionists and ICU staff.

Activated charcoal (oral or via nasogastric tube) · Adsorbent

If the overdose was recent (within 1–2 hours) and the patient is awake enough to protect their airway, activated charcoal is a fine black powder that acts like a sponge. CCB pills that have not yet dissolved in the stomach stick to the charcoal, which then carries them out in the stool before they can be absorbed into the blood.

1. Patient swallows or is given CCB pills.
2. Pills begin to dissolve in stomach acid.
3. Activated charcoal (50 grams for adults, 1 gram/kg for children) is given by mouth or NG tube within 1–2 hours.
4. CCB molecules bind tightly to the charcoal surface.
5. The charcoal and bound drug pass through the gut and are eliminated in stool, preventing absorption.

SAFETY WATCH

Do NOT give if the patient is drowsy, vomiting, or cannot protect their airway—risk of aspiration (charcoal in the lungs) which is life-threatening. · Causes black stool and sometimes constipation. · Contraindicated if the patient has a bowel perforation or obstruction. · For extended-release CCB formulas, repeat doses every 4 hours may be considered under specialist guidance.

Whole bowel irrigation with polyethylene glycol (PEG) solution · Osmotic purge

Extended-release CCB pills can sit in the gut for many hours, slowly leaking poison. We flood the intestines with a large volume of salty liquid (the same prep used before colonoscopy) to flush everything through before the pills fully dissolve. Think of it as a fire hose washing out the plumbing.

1. Sustained-release CCB tablets pass slowly through the gut, releasing drug over 12–24 hours.
2. PEG solution (GoLYTELY, CoLyte) is given at 1–2 liters per hour via NG tube or by mouth.
3. The huge volume of fluid overwhelms the gut's ability to absorb it.
4. Stool becomes clear liquid, carrying unabsorbed pills and dissolved drug out before they can enter the blood.
5. Continue until rectal effluent (what comes out) is clear, usually 4–6 hours.

SAFETY WATCH

Do NOT use if there is bowel obstruction, perforation, or severe a stalled gut (ileus) (paralyzed gut). · Patient must be alert or have a protected airway (intubation) to prevent aspiration. · Causes severe diarrhea, electrolyte shifts (watch sodium and potassium), and skin breakdown around the rectum. · Labor-intensive; requires ICU-level monitoring.

Red flags

- Heart rate below 50 beats per minute or systolic blood pressure below 90 mmHg
- Confusion, decreased level of consciousness, or unresponsiveness
- Seizures or signs of cardiogenic shock (cold, clammy skin; weak pulse; rapid breathing)
- Chest pain, severe shortness of breath, or fluid in the lungs on exam
- Ingestion of an extended-release formulation—symptoms can worsen suddenly hours later

Nursing priorities

1. Establish continuous cardiac monitoring and measure vital signs every 5 to 15 minutes, watching for heart rate below 50 or systolic blood pressure below 90. — Calcium channel blockers block calcium entry into heart cells, which slows electrical conduction through the AV node (the heart's natural pacemaker relay station) and weakens heart muscle squeezing, so heart rate and blood pressure can drop suddenly and require immediate treatment escalation.
2. Secure two large-bore IV lines (18-gauge or larger) and prepare to administer IV calcium chloride or calcium gluconate as ordered. — Calcium is the direct antidote that competes with the calcium channel blocker at receptor sites, restoring calcium entry into heart and blood vessel muscle cells to improve heart contraction strength and raise blood pressure, and large IV lines allow rapid high-volume infusions.
3. Check blood glucose immediately and then every 30 to 60 minutes, and administer dextrose if glucose drops below 70 mg/dL. — Calcium channel blockers block calcium channels in pancreatic beta cells (insulin-releasing cells), which shuts off insulin release and causes high blood sugar (hyperglycemia) (high blood sugar), but the high-dose insulin treatment we give can then cause severe low blood sugar (hypoglycemia) (low blood sugar) that damages the brain if not caught early.
4. Prepare high-dose insulin infusion with a dextrose drip at the bedside, double-checking the insulin dose calculation with another nurse before starting. — Insulin forces glucose and calcium into heart muscle cells independent of the blocked calcium channels, restoring the heart's ability to contract strongly (inotropy), and this is often more effective than calcium alone in severe overdoses, but wrong insulin doses can kill through low blood sugar (hypoglycemia).
5. Monitor for signs of fluid overload—lung crackles, increasing shortness of breath, oxygen saturation dropping—especially during calcium and fluid resuscitation. — The weakened heart cannot pump effectively (reduced contractility), so extra IV fluids we give to raise blood pressure can back up into the lungs causing pulmonary edema (fluid flooding the lungs) (fluid in air sacs), which blocks oxygen exchange and can cause respiratory failure requiring intubation.
6. Keep atropine, transcutaneous pacing pads, and vasopressor infusions (norepinephrine, epinephrine, dopamine) immediately available at the bedside. — Severe a slow heart rate (bradycardia) may not respond to calcium or insulin and require atropine (blocks the vagus nerve to speed heart rate) or emergency electrical pacing, and persistent low blood pressure (hypotension) despite treatment may need vasopressors (drugs that squeeze blood vessels) to prevent organ damage from low blood flow.

7. Assess neurological status every 15 minutes using Glasgow Coma Scale or AVPU (Alert, Verbal, Pain, Unresponsive), and immediately report decreasing consciousness or confusion. — Dropping blood pressure means less blood reaches the brain (cerebral poor blood flow (hypoperfusion)), and altered mental status is an early warning that brain tissue is being injured; catching this early allows treatment intensification before permanent brain damage occurs.
8. Document exact time of ingestion and prepare for possible extended monitoring or ICU transfer, especially if extended-release formulation was involved. — Extended-release tablets continue releasing drug for 12 to 24 hours, so a patient who looks stable at hour two can suddenly crash at hour eight when more drug absorbs; delayed toxicity requires prolonged observation and immediate ICU resources if deterioration occurs.

Sources

- American College of Medical Toxicology (ACMT) position statements
- American Heart Association (AHA) ACLS guidelines
- UpToDate calcium-channel blocker poisoning monograph
- Journal of Medical Toxicology consensus guidelines

Last reviewed: 2026-07-07