

Central DI

The back part of the pituitary gland stops releasing enough ADH (the hormone that tells your kidneys to save water), so the kidneys dump huge amounts of pale, watery urine and the body gets dangerously thirsty.

THE STORY IN ONE SENTENCE

The back part of the pituitary gland stops releasing enough ADH (the hormone that tells your kidneys to save water), so the kidneys dump huge amounts of pale, watery urine and the body gets dangerously thirsty.

“Central DI — the brain stops sending the 'hold water' signal, so the kidneys flush out gallons of dilute urine and the patient must drink constantly or sodium will spike and cause confusion, seizures, or coma.”

Normal anatomy — the body's job

Special nerve cells in the hypothalamus (a grape-sized control center deep in the middle of the brain) make ADH and send it down a narrow stalk into the posterior pituitary (the back lobe of the pituitary gland), which stores the hormone and releases it into the bloodstream when the body needs to save water.

BODY'S JOB

ADH travels through the blood to both kidneys and locks onto receptors called V2 receptors on the walls of the collecting ducts—the final stretch of the kidney's filtering tubes where urine becomes concentrated. That signal tells the duct cells to pull water channels called aquaporin-2 up to the surface, opening tiny doors so water flows back into the blood instead of leaving as urine. This keeps your body's water level steady and your blood sodium in the safe range (135 to 145 milliequivalents per liter), even when you sweat, breathe out moisture, or go hours without drinking.

1. Nerve cells in the hypothalamus manufacture ADH (also called vasopressin) and package it into tiny storage bubbles inside the nerve fiber.
2. The bubbles slide down long nerve fibers—like beads on a string—through the pituitary stalk into the posterior pituitary gland at the base of the brain.

3. When sensors in the hypothalamus detect that your blood is getting too salty (osmolality above 285 to 290 milliosmoles per kilogram) or your blood volume is dropping, the posterior pituitary squirts ADH into nearby blood vessels.
4. ADH rides through the bloodstream to both kidneys in about one minute.
5. In the kidney's collecting ducts (the final tubes before urine enters the bladder), ADH docks onto V2 receptors on the duct wall.
6. The receptors trigger the duct cells to move aquaporin-2 water channels from storage inside the cell up to the surface facing the urine.
7. Water that was about to leave as urine flows back through those channels into the blood, concentrating the urine (making it darker and more yellow) and diluting the blood (lowering sodium back to normal).
8. Blood sodium and osmolality return to the safe range, blood pressure stays steady, and the thirst signal turns off.

What breaks

The nerve cells in the hypothalamus that make ADH, the stalk that carries it down to the pituitary, or the back part of the pituitary itself get damaged or destroyed, so ADH production or release drops sharply or stops completely.

CLINICAL TERM

Central diabetes insipidus results from destruction or dysfunction of the hypothalamic magnocellular neurons or the neurohypophyseal tract, leading to deficient synthesis, transport, or secretion of arginine vasopressin (ADH).

How the body compensates

The body has no backup factory for ADH and no other hormone that can do its job, so the only way to replace lost water is through thirst and drinking. As long as the person can sense thirst (which requires an intact hypothalamus) and drink freely, blood pressure and sodium stay near normal—but this balance is fragile and tips fast if water access is blocked or thirst is impaired.

1. {'step': 'Osmoreceptors (special sensors in the hypothalamus that measure how salty the blood is) detect rising blood sodium and osmolality and trigger an intense, powerful urge to drink.', 'why': 'Drinking replaces the water lost in urine liter for liter, diluting the blood back toward normal and keeping sodium from climbing into the danger zone above 145 milliequivalents per liter.'}

2. {'step': 'Heart rate and blood pressure usually stay normal or near-normal if the person drinks enough to match urine output.', 'why': 'Constant drinking refills blood volume as fast as the kidneys dump it, so the heart does not have to speed up and blood vessels do not collapse—yet. The person is in a precarious steady state.'}
3. {'step': 'Urine stays pale, clear, and dilute (specific gravity less than 1.005, osmolality less than 200 milliosmoles per kilogram) no matter how high blood sodium climbs, because the kidneys cannot concentrate it without ADH.', 'why': 'The kidney's collecting ducts never get the signal to open water channels, so they act as if the body has plenty of water even when it does not. The only real compensation is behavioral—drinking more—not a kidney response.'}
4. {'step': 'If the person cannot drink—because they are asleep, sedated, intubated, an infant too young to ask for water, elderly with a blunted thirst drive, or confused from another illness—serum sodium shoots up fast, often climbing several milliequivalents per liter every few hours.', 'why': 'Water loss continues at a rate of several liters per day (sometimes 500 to 1000 milliliters per hour), but water intake stops, so the blood becomes dangerously concentrated and sodium can reach 150, 160, or even 170 milliequivalents per liter within hours to days.'}
5. {'step': 'Severe high sodium (hypernatremia) (sodium above 155 to 160 milliequivalents per liter) pulls water out of brain cells by osmosis, shrinking them and tearing the small bridging veins that connect the brain surface to the skull.', 'why': 'Water always moves from areas of lower salt concentration (inside the brain cells) to higher concentration (the bloodstream). Brain cells try to protect their volume by making extra internal particles (idiogenic osmoles) over one to two days, but if sodium rises faster than that, the cells cannot keep up. The result is brain cell dehydration, confusion, irritability, muscle twitching, seizures, coma, and bleeding inside the skull (subdural or subarachnoid bleeding (hemorrhage)).'}
6. {'step': 'Vital signs (blood pressure, pulse, breathing rate) can look deceptively stable during early high sodium (hypernatremia) because total body water loss happens gradually and thirst-driven intake masks the deficit until intake stops.', 'why': 'The danger is hidden. A patient may look reasonably well while awake and drinking, but once they fall asleep, go to surgery, or become too ill to drink, decompensation is rapid and the safe window to correct sodium (no faster than 10 to 12 milliequivalents per liter per day to avoid cerebral swelling (edema)) becomes very narrow.'}

7. {'step': 'Once the diagnosis is made and desmopressin (DDAVP) is started, a new opposite risk appears: if the DDAVP dose is too high, given too often, or the patient continues drinking out of habit or residual thirst, free water is retained without being peed out, and sodium can drop too low (hyponatremia, below 135 milliequivalents per liter).', 'why': 'DDAVP turns off urine output by restoring the kidney's ability to hold water, but the thirst center in the hypothalamus may take a day or two to reset. Excess water intake without a urinary escape route dilutes the blood, causes brain cells to swell (cerebral swelling (edema)), and can trigger headache, nausea, confusion, seizures, or even death if sodium falls below 120 milliequivalents per liter.'}

Domino effect

1. ADH production or release stops because the magnocellular neurons in the hypothalamus or the pituitary stalk are damaged or destroyed. — Without ADH traveling through the blood to the kidneys, the collecting ducts never get the signal to insert aquaporin-2 water channels into their walls, so water that should return to the blood is dumped into the bladder and lost as urine.
2. Urine output jumps suddenly and dramatically to three to twenty liters per day (polyuria), and the urine looks clear, pale, and watery with a specific gravity under 1.005 and osmolality under 200 milliosmoles per kilogram (normal concentrated urine is 600 to 1200). — The kidney filters about 180 liters of blood every day and normally pulls almost all the water back, leaving one to two liters as urine. Without ADH, the kidney cannot pull water back in the collecting ducts, so nearly all the filtered water leaves the body, flooding the bladder every hour or two.
3. Massive water loss shrinks the blood volume and makes the blood more concentrated, raising plasma sodium above 145 milliequivalents per liter and plasma osmolality above 295 milliosmoles per kilogram. — Water leaves the bloodstream much faster than sodium does (the kidneys still hold onto sodium through other hormones like aldosterone), so the ratio of sodium to water climbs. Blood becomes saltier and thicker.
4. Rising blood osmolality triggers osmoreceptors in the hypothalamus, which fire signals that create intense, unrelenting thirst (polydipsia), and the person drinks constantly—often preferring ice-cold water or ice chips. — The hypothalamus detects the high sodium and turns on the thirst drive to pull water back into the body from the outside and dilute the blood back to normal. Thirst in central DI is often described as more intense and specific than ordinary thirst.

5. If the person can drink freely and has a normal thirst mechanism, sodium may stay near normal or only mildly elevated (145 to 150 milliequivalents per liter), but they are trapped in an exhausting cycle of urinating and drinking around the clock, often every one to two hours, day and night. — Water intake roughly matches water output (input equals output), so the body stays in a precarious, fragile balance. The person cannot leave the house for long, cannot sleep through the night, and must always be near a bathroom and a water source.
6. If water intake stops or falls behind output—during sleep, anesthesia, surgery, nausea, vomiting, loss of consciousness, or in infants, young children, and debilitated adults who cannot ask for or reach water—serum sodium climbs rapidly, often spiking from 145 to 160 milliequivalents per liter or higher within hours to one day. — Urine output does not slow down at all (there is still no ADH), so water keeps pouring out at several hundred milliliters per hour while intake drops to zero. Blood sodium rises at a rate of 1 to 2 milliequivalents per liter per hour, much faster than the brain can adapt.
7. Severe high sodium (hypernatremia) (sodium above 155 to 160 milliequivalents per liter) shrinks brain cells by pulling water out of them, stretches and tears bridging veins between the brain and skull, and disrupts electrical signaling in neurons, causing early symptoms of irritability, restlessness, and intense thirst, then confusion, lethargy, muscle twitching, weakness, fever, seizures, and coma. If untreated or corrected too fast, permanent brain injury or death can occur. — Brain cells cannot tolerate sudden dehydration. The mechanical stress of shrinking tears blood vessels (causing subdural a pooled collection of blood (hematoma) or subarachnoid bleeding (hemorrhage)), and the high sodium disrupts the electrical gradients neurons need to fire properly. If sodium is corrected too quickly (faster than 10 to 12 milliequivalents per liter per day), brain cells swell rapidly as water rushes back in, causing cerebral swelling (edema), brainstem herniation, and death.
8. Once desmopressin (DDAVP, a synthetic analog of ADH) is given—usually as a nasal spray, sublingual tablet, or subcutaneous injection—the kidneys immediately start holding water again. Within one to two hours, urine output drops from liters per hour to normal (50 to 100 milliliters per hour), urine becomes concentrated and yellow, and serum sodium drifts back toward normal over 24 to 48 hours. Close monitoring with frequent sodium checks (every 4 to 6 hours initially) is essential to avoid overcorrection and caused by medical care (iatrogenic) low sodium (hyponatremia). — DDAVP replaces the missing ADH signal by binding to the same V2 receptors in the kidney that natural ADH uses, restoring the kidney's ability to insert aquaporin-2 channels and concentrate urine. Careful dosing (starting low, often 0.05 to 0.1 milligrams DDAVP by mouth or 1 to 2 micrograms subcutaneously) and fluid balance (matching intake to output, not forcing fluids) prevent the sodium from swinging from too high to dangerously low. Patients often need one to two doses per day for life, adjusted based on urine output, thirst, and sodium levels.

Symptoms · what the patient feels, and why

- Making huge amounts of urine every day (often 3–20 liters, which is like 3–20 big soda bottles): The brain normally makes a hormone called ADH (anti-diuretic hormone, or the 'save water' signal) in a tiny region called the hypothalamus and stores it in the pituitary gland (a pea-sized gland at the base of the brain). ADH travels through the blood to the kidneys and tells them to hold on to water. In central DI, the hypothalamus or pituitary is damaged by surgery, a tumor, head injury, infection, or sometimes for no clear reason, so the brain cannot make or release enough ADH. Without this signal, the kidneys never get the command to stop water from pouring out, and the body loses water like a leaky bucket.
- Constant, strong thirst and drinking water all day and night: When you lose so much water in the urine, your blood becomes more concentrated (like juice concentrate before you add water). Sensors in the hypothalamus (the brain's thirst center) notice this change and trigger a powerful thirst signal so you will drink and replace the lost water. People who are awake and can drink usually do this automatically and stay safe, but babies, very sick people, or anyone who cannot drink enough can become dangerously dehydrated.
- Getting up many times at night to pee and drink (nocturia): Because the kidneys keep making huge amounts of urine around the clock—even at night—the bladder fills up over and over, waking the person to empty it and then to drink more water to keep up with the loss. This cycle can happen four, five, or more times every night.
- Urine that looks almost clear or very pale, like water: Normal urine is yellow because the kidneys concentrate waste into a smaller amount of water. In central DI, the kidneys are flushing out huge volumes of water that are barely concentrated at all, so the urine is very dilute and pale—its specific gravity (a measure of how concentrated urine is) is usually very low, under 1.005.
- Feeling tired, weak, or foggy: Constantly losing water and having to drink and pee day and night interrupts sleep and wears the body down. Also, if the thirst response cannot keep up—or if someone cannot drink freely—the blood can become too concentrated, which slows down how cells work and makes the person feel sluggish and confused.
- Dry mouth and dry skin: When the body is losing more water than it is taking in, the skin and mucous membranes (the moist linings of the mouth and lips) lose moisture, like a plant that isn't watered enough starts to wilt. This is an early sign that the body is getting behind on replacing fluids.
- Fast heart rate and low blood pressure (especially when standing up): If water loss gets ahead of drinking, the amount of blood in the vessels drops (called hypovolemia, meaning low blood volume—like a garden hose with less water running through it). The heart has to beat faster to keep enough blood moving to the brain and organs. Blood pressure can fall, especially when standing, because there is not enough fluid volume to fill the system. This drop when standing is called orthostatic low blood pressure (hypotension).

- Confusion, extreme drowsiness, or seizures (if the blood sodium gets dangerously high): When someone with central DI cannot drink enough to match the huge urine losses—for example, after surgery when they are not fully awake, or in a baby who cannot ask for water—the blood becomes too concentrated and the sodium level shoots up (hypernatremia, meaning too much sodium in the blood, usually above 145–150 mEq/L or higher). High sodium pulls water out of brain cells by osmosis (water moves toward the saltier side), shrinking the cells and causing the brain to misfire. This can lead to confusion, sleepiness, muscle twitching, seizures, or even coma if not fixed quickly.
- Weight loss (especially in babies and young children): Babies and small children with central DI lose water faster than they can signal thirst or drink on their own. The constant fluid loss keeps them from growing normally and can cause weight to drop. In infants, this often shows up as failure to thrive (not gaining weight as expected).
- Irritability and fussiness (especially in infants who cannot say they are thirsty): Babies with central DI feel the same powerful thirst and discomfort as adults, but they cannot ask for water or explain what is wrong. They cry, refuse to eat solid food (because they want fluids), and become cranky because their bodies are constantly dry inside.

Focused exam

- Measure urine output over 24 hours — is it more than 3 liters?: Central DI happens when the hypothalamus (the deep brain area that makes hormones) or the back pituitary (where ADH is stored and released) stops putting out ADH, the hormone that tells the kidneys to hold onto water. Without it, the kidneys dump water like an open faucet. Peeing more than 3 liters per day (or more than 40–50 mL per kilogram of body weight in adults) is the main sign. If output is less, you're looking at something else, not Central DI.
- Ask about thirst — does the patient drink huge amounts of water right after peeing?: When water pours out, the blood gets saltier, and the brain's thirst center (in the hypothalamus) fires off a powerful thirst signal. An awake patient drinks gallons to balance it out. If they can't drink (because they're sedated, confused, or in a coma), the salt climbs fast and can cause seizures or brain swelling. Thirst tells you whether the patient can protect themselves.
- Check blood sodium level — is it high, normal, or low?: High sodium (above 145 mEq/L) means the patient is losing water faster than they can replace it — they're dehydrated and at risk for brain damage. Normal or low sodium (135–145 or below 135 mEq/L) means they drank enough or already got DDAVP (the ADH replacement medicine). Sodium is your safety number — watch it closely because it tells you whether the patient is winning or losing the water race.

- Check urine specific gravity or osmolality — is the pee dilute (watery)? In Central DI, pee is very dilute (specific gravity under 1.005 or osmolality under 200 mOsm/kg) because the kidneys can't concentrate it without ADH — it's like trying to make lemonade without adding the powder, just pouring out water. Dilute pee plus high urine output means ADH is missing or broken.
- Ask about recent brain surgery, head trauma, or brain imaging findings: Central DI usually starts after pituitary surgery, a severe head injury, or when imaging finds a tumor pressing on the hypothalamus or pituitary gland. The timing is important — symptoms often appear within days to weeks after the event because it takes time for swelling or damage to knock out the nerve cells that make ADH.
- Check how alert the patient is — awake and talking, or drowsy and confused?: Awake patients can drink when they're thirsty and keep their sodium safe. Patients who are sedated, in a coma, or confused can't drink, so their sodium climbs and they can have seizures or brain swelling. Mental status changes the danger level fast — a confused patient with Central DI is a medical emergency.

Diagnostics · every test answers a question

- Serum sodium (blood sodium) → Is the blood getting too salty because water is leaving the body?
- 24-hour urine output measurement → Is the patient peeing too much?
- Urine osmolality and specific gravity → Is the pee dilute (watery) or concentrated?
- Water deprivation test (supervised in the hospital) → Can the body concentrate urine when it's dehydrated, or is ADH missing?
- Serum osmolality → Is the blood too concentrated (too many particles, not enough water)?
- Brain MRI with focus on the pituitary gland and hypothalamus → Is there damage, a tumor, or missing tissue in the pituitary gland or hypothalamus?
- Morning cortisol and thyroid-stimulating hormone (TSH) with free T4 → Is the rest of the pituitary gland working, or is it damaged too?
- Copeptin level (where available) or direct ADH level (rarely done) → Is ADH actually low or absent in the blood?

Treatment logic

- DDAVP (desmopressin) — nasal spray, tablet under the tongue (sublingual), oral tablet, or IV/subcutaneous injection → Replaces the missing ADH signal, so the kidneys start holding onto water again instead of dumping it. Urine output drops from over 3 liters back down to normal (1–2 liters per day), and extreme thirst goes away. Blood sodium stabilizes in a safe range. DDAVP is a synthetic version of ADH that lasts longer and is easier to dose — typical starting doses are 5–10 mcg nasal spray, 0.1–0.2 mg oral, or 1–2 mcg IV/subcutaneous.
- Let the patient drink freely when thirsty — do not force fluids or restrict them → An awake patient's thirst tells them exactly how much water their body is losing, so they drink just enough to stay balanced. The brain's thirst center (in the hypothalamus) is very precise — it tracks blood salt level and tells the patient to drink when sodium climbs. This prevents both dehydration (from not drinking enough) and water overload (from drinking too much).
- Check blood sodium every 6–12 hours in the first few days, then every 1–2 weeks once stable → Catches high sodium (from dehydration or not enough DDAVP) or low sodium (from too much DDAVP or giving too much free water through an IV) before the patient has seizures or brain swelling. Sodium swings can happen fast in the first days after surgery or injury, so frequent checks keep you ahead of trouble.
- Find and treat the underlying cause — MRI to look for tumors, infection, or inflammation; follow up after pituitary surgery; treat autoimmune hypophysitis with steroids if present → If a tumor is pressing on the hypothalamus or pituitary, removing it can relieve the pressure and let the nerve cells heal. If inflammation (hypophysitis) is damaging the gland, steroids (like prednisone) can stop the attack. In some cases, ADH production can restart on its own over weeks to months after the cause is treated.
- Avoid giving hypotonic (low-salt) IV fluids like half-normal saline (0.45% NaCl) or D5W (5% dextrose in water) in patients taking DDAVP — use normal saline (0.9% NaCl) or let them drink free water instead → Prevents the kidneys from diluting the blood dangerously while DDAVP is forcing them to hold onto water. Hypotonic fluids flood the body with free water when the kidneys can't get rid of it, causing low sodium (hyponatremia below 135 mEq/L), which can lead to seizures or brain swelling. Normal saline matches the blood's salt, so it doesn't dilute anything.
- If the patient can't drink (sedated, on a ventilator, confused), give IV fluids that match urine output exactly — measure pee every 1–2 hours and replace it milliliter for milliliter with normal saline or a similar isotonic fluid → Keeps the patient's water balance neutral when they can't drink for themselves. This prevents dehydration and keeps sodium in a safe range (135–145 mEq/L). It's like filling a bucket as fast as it's leaking — you stay even.

- Replace other missing pituitary hormones if present (cortisol, thyroid, sex hormones, growth hormone) → Keeps the patient alive and healthy when the whole pituitary is damaged (panhypopituitarism). Low cortisol (adrenal insufficiency) is especially dangerous during illness or stress because the body can't fight infection or maintain blood pressure without it. Thyroid hormone keeps metabolism and energy normal.

Medications

Desmopressin (DDAVP) · Synthetic vasopressin analog (lab-made copy of ADH, the water-saving hormone)

Desmopressin is a lab-made copy of ADH (anti-diuretic hormone), the signal your brain normally makes to tell your kidneys to save water. In central DI, your hypothalamus or pituitary gland cannot make enough ADH, so your kidneys dump huge amounts of water into your urine—sometimes 10 to 20 liters a day—leaving you desperately thirsty and peeing constantly. Desmopressin replaces that missing signal. It travels to your kidneys and tells the collecting ducts (the final tubes where urine gets concentrated) to pull water back from the urine into your blood, the same way real ADH would. Instead of gallons of pale, watery pee, you make normal amounts of darker, concentrated urine. You can hold onto the water you drink, so you stop feeling so thirsty and your blood sodium stays in the healthy range.

1. You take desmopressin—either as a tablet by mouth, a nasal spray that you puff into your nose, or an IV shot that goes straight into your vein if you are in the hospital.
2. The drug travels through your bloodstream to your kidneys.
3. Desmopressin locks onto V2 receptors sitting on the cells that line the collecting ducts (the last tube section before urine becomes final).
4. This lock-and-key fit flips a switch inside the cell, telling it to move tiny water channels called aquaporins up to the tube wall (like sliding open little doors in a fence).
5. Water from the forming urine flows back through those aquaporin channels into your blood, instead of leaving your body as pee.
6. Your urine becomes more concentrated (darker yellow, smaller volume—maybe 1 to 2 liters a day instead of 10), and your blood stays properly diluted so your sodium level holds steady in the normal range (135 to 145 mEq/L).
7. Your brain's thirst alarm quiets down because your blood is no longer too salty, so you stop feeling parched all the time.

SAFETY WATCH

Low sodium (Hyponatremia) (dangerously low blood sodium): This is the biggest risk. If you drink too much water while taking desmopressin, your kidneys hold onto nearly all of it because the drug is doing its job—saving water. That floods your blood and dilutes your sodium below 135 mEq/L, sometimes far below. Early warning signs are headache, nausea, confusion, feeling foggy, and muscle cramps. Severe low sodium can cause seizures, coma, brain swelling, and even death. To prevent this, drink only when you are truly thirsty—do NOT chug water on a schedule or try to drink a certain number of glasses a day. Let your body tell you when you need water. Skip one dose of desmopressin each week (or as your doctor advises) to let your kidneys clear out any extra water. Your doctor will check your sodium level regularly, especially when you first start the medicine or change the dose. · Never give IV hypotonic fluids (weak salt solutions like half-normal saline, quarter-normal saline, or D5W—plain sugar water) to someone taking desmopressin, especially if they are unconscious, sedated, or post-surgery and cannot feel or tell you they are full. These fluids pour water into the body faster than the kidneys can get rid of it (because desmopressin blocks water excretion), and sodium can plummet within hours, causing seizures and brain damage. · Children and older adults are at higher risk for water overload and low sodium. Watch them closely. Babies and toddlers cannot tell you they feel sick, so monitor for fussiness, vomiting, weight gain (from retained water), or lethargy. · If you develop a bad headache, confusion, nausea, vomiting, or just feel wrong while on desmopressin, STOP the drug immediately, do not drink any more water, and call your doctor or go to the emergency room. You may need a blood test to check your sodium, and if it is dangerously low, you may need careful IV saline to bring it back up slowly (too fast can also hurt your brain). · Nasal spray may not absorb well if you have a stuffy nose from a cold, allergies, nasal surgery, or scarring inside your nose from chronic inflammation or cocaine use. If the spray stops working, you may need to switch to the tablet or injection form. · Rarely, allergic reactions can happen—rash, itching, swelling of the face or throat, trouble breathing. If this occurs, stop the drug and get emergency help. · Do NOT use desmopressin if you have psychogenic excessive thirst (polydipsia) (compulsive water drinking driven by a mental health condition), because these patients will keep drinking huge amounts of water on top of the drug, causing severe, life-threatening low sodium (hyponatremia).

Red flags

- Making huge amounts of urine right after brain surgery, a head injury, or while very sick—this can mean the hypothalamus or pituitary gland is damaged and the brain's water-control system needs fast treatment.

- Confusion, extreme sleepiness, seizures, or muscle twitching along with massive urination—this means the blood sodium may be climbing dangerously high (hypernatremia) because the person cannot drink enough to keep up.
- A baby or young child who is always thirsty, soaking many diapers with clear urine, irritable, and not growing well—central DI in infants is easy to miss and can cause serious dehydration and brain damage very quickly.
- Sudden onset of uncontrollable thirst and urination in someone with a known pituitary tumor, a history of brain infection (such as meningitis or encephalitis), or autoimmune inflammation of the pituitary—this may signal new or worsening pituitary damage.
- Signs of severe dehydration: sunken eyes, dry lips and tongue, skin that stays pinched when you pull it up (tenting), very little or no tears when crying, or feeling dizzy and faint—this means the body is critically low on water and needs emergency fluids.

Nursing priorities

1. Check blood sodium level (and serum osmolality if available) immediately and then every 6–12 hours until stable, watching for values above 145 mEq/L. — Without ADH, the kidneys dump free water and blood sodium climbs fast (hypernatremia). Sodium above 150 mEq/L shrinks brain cells, causing confusion, seizures, coma, and bleeding inside the brain. Early detection lets us replace water and give DDAVP before dangerous levels occur.
2. Measure and record every drop of urine output and all fluid intake every hour, using a urometer or measuring container at the bedside. — Central DI can cause 500–1000 mL of urine per hour. Tracking input and output shows whether DDAVP is working (urine volume should drop) and whether the patient is drinking enough to match losses. Large negative fluid balance signals worsening dehydration.
3. Give DDAVP (desmopressin) exactly as ordered—nasal spray, sublingual tablet, or IV injection—and watch urine output drop within 1–2 hours. — DDAVP replaces the missing ADH, telling the kidneys to hold onto water. Urine volume should fall and urine concentration (specific gravity or osmolality) should rise. If output stays high, the dose may be too low or the diagnosis may be nephrogenic DI (kidney does not respond to ADH).
4. Let the patient drink water freely whenever thirsty—keep a pitcher of ice water at the bedside and never restrict fluids. — The thirst mechanism is the body's main defense against high sodium in central DI. Forcing the patient to limit drinking causes dangerous high sodium (hypernatremia). Only drinking when thirsty prevents over-replacement and low sodium (hyponatremia), especially after DDAVP starts working.

5. Check urine specific gravity (normal 1.005–1.030) or urine osmolality with every void or every 4 hours to see if the kidneys are concentrating urine. — In untreated central DI, urine specific gravity stays very low (under 1.005) because the kidneys cannot concentrate. After DDAVP, specific gravity should climb above 1.010, proving the kidneys are responding and saving water. Persistently low values suggest wrong diagnosis or nephrogenic DI.
6. Watch for signs of water overload—headache, nausea, confusion, weight gain—especially in the first 24 hours after starting DDAVP, and hold the next dose if sodium drops below 135 mEq/L. — DDAVP can work too well, causing the kidneys to hold too much water and dilute the blood sodium (hyponatremia). Low sodium swells brain cells and causes seizures. Holding DDAVP and restricting fluids for a few hours lets excess water leave through normal kidney function.
7. Assess mental status, muscle strength, and deep-tendon reflexes every 4 hours, and report new confusion, twitching, or seizures immediately. — Both high sodium (from untreated DI) and low sodium (from DDAVP overdose) change brain-cell size and cause neurologic symptoms. Quick changes in sodium—up or down—are more dangerous than stable abnormal levels. Rapid detection prevents permanent brain damage (osmotic demyelination or cerebral swelling (edema)).
8. Coordinate brain MRI or CT scan as ordered to look for pituitary tumors, hypothalamic lesions, or signs of surgery or trauma. — Imaging finds the cause of central DI—tumor, cyst, inflammation, bleeding, or absent posterior pituitary bright spot. Knowing the cause guides whether the patient needs surgery, long-term DDAVP, or treatment for infection or autoimmune disease.

Sources

- Endocrine Society Clinical Practice Guideline on Diagnosis and Treatment of Diabetes Insipidus (2021)
- UpToDate (peer-reviewed clinical reference)
- Pituitary Society consensus statements
- AACE (American Association of Clinical Endocrinologists)

Last reviewed: 2026-07-07