

Child Abuse and Neglect Recognition

Think abuse when the injury story does not match the injury itself or what a child that age can do.

THE STORY IN ONE SENTENCE

Think abuse when the injury story does not match the injury itself or what a child that age can do.

“Watch for harm and neglect in kids. Spot the patterns to save lives. Every state says you must report it—it is the law.”

Normal anatomy — the body's job

Healthy children are cared for by adults who keep them safe, fed, clean, and protected from harm. Injuries in normal childhood are accidental, match the developmental stage (what the child can do at that age), and have a clear, consistent story.

BODY'S JOB

Caregivers watch over kids, provide food and shelter, keep dangerous objects away, supervise play, and get medical help quickly when a child is hurt or sick. The child's bones, skin, and brain develop and heal normally when there is no repeated trauma.

1. A baby cannot roll, crawl, or grab, so injuries in a non-mobile infant are red flags.
2. A toddler learning to walk falls forward onto hands, knees, and forehead—bruises there are normal.
3. Accidents happen suddenly, are witnessed or immediately discovered, and the story stays the same every time you ask.
4. Caregivers seek help right away when a child is hurt, and they cooperate with the medical team without blaming the child.

What breaks

Abuse is when an adult or older child intentionally harms a child—hitting, shaking, burning, sexual contact, cruel words, or failing to give basic needs like food or medical care. The upstream break is the loss of protection and safety.

CLINICAL TERM

Child maltreatment includes physical abuse (non-accidental trauma), sexual abuse, emotional abuse, and neglect (failure to provide adequate food, supervision, medical care, or a safe environment). CDC and AAP define these as acts or failures to act that cause actual or potential harm.

How the body compensates

The child's body tries to survive each injury by clotting blood, healing fractures, and maintaining vital signs, but it cannot protect itself from the next episode of harm. The danger is that vital signs may look stable between injuries while the real threat—ongoing abuse—continues.

1. After a hit or shake, the brain swells and bleeds, but blood pressure and heart rate may stay normal at first because the young body compensates well.
2. Fractures begin to heal and form new bone (callus) within 7–10 days, so seeing fractures at different stages of healing proves repeated injury over time, not one accident.
3. Bruises change color as the body breaks down blood—red or purple when fresh, green or yellow after several days—so multiple bruises in different colors mean multiple episodes of hitting.
4. The caregiver may bring the child in only when the injury is so severe that compensation fails—seizure, unconsciousness, or visible deformity—missing the window to intervene earlier.
5. Siblings at home are at high risk because the same abuser continues the same behavior; protecting the patient means also protecting brothers and sisters.
6. The child cannot escape or change the environment, so the cycle repeats unless the medical team, social work, and child protective services break it.

Domino effect

1. Caregiver loses control, uses force, or neglects basic care. — Abuse starts when the adult fails to protect—either actively harming (shaking a crying baby, hitting with a belt, burning as punishment) or passively neglecting (not feeding, ignoring medical needs). This is the upstream break.

2. Child suffers specific injuries that match the mechanism of harm. — The type of force leaves a signature: shaking causes subdural a pooled collection of blood (hematoma) (bleeding between brain and skull) and retinal hemorrhages (bleeding in the back of the eyes); grabbing and twisting causes metaphyseal fractures (chip fractures at the ends of long bones); dunking in hot water causes symmetric burns with sharp lines; looped cords leave parallel bruises or marks.
3. Body tries to heal—bones form callus, bruises change color, brain swells. — The child's body responds to each injury by clotting, laying down new bone, and mounting an inflammatory response, but these healing signs actually become evidence: different-aged fractures prove repeated trauma, and patterned bruises prove inflicted force, not accidents.
4. Caregiver delays bringing the child in, changes the story, or blames the child. — The abuser knows the injury looks bad, so they wait, hoping it will get better on its own, or they make up a story that does not match (claiming a 2-month-old rolled off the bed when the baby cannot roll yet). This mismatch is a red flag.
5. Medical team sees the pattern: injury plus inconsistent history plus delay. — No single finding proves abuse, but the combination—bruises in protected areas (ears, neck, genitals), fractures in non-mobile infants, burns with clear borders, plus a story that does not fit—points to intentional harm, not accident.
6. Team reports to child protective services and law enforcement (mandatory). — Every healthcare provider is a mandated reporter by law in all 50 states. Reporting is not about proving abuse in the ED—it is about triggering a full investigation to protect this child and any siblings. Failure to report leaves the child at risk of further injury or death.
7. Without intervention, abuse escalates—more severe injuries, permanent disability, or death. — Abusive head trauma has a 25–30% mortality rate and high rates of long-term brain damage in survivors. Repeated fractures can lead to growth problems and chronic pain. The cycle continues until someone outside the home breaks it.
8. Early recognition and reporting stop the cycle and save lives. — When the medical team identifies the pattern, involves social work, and reports to authorities, the child can be placed in a safe environment, the abuser can be held accountable, and siblings can be evaluated and protected. This is the off-ramp from the domino chain.

Symptoms · what the patient feels, and why

- Injury does not match the story or age: A baby who cannot yet roll over should not have bruises from a fall. When the injury is more serious than the explanation, or impossible for that child's age, it raises the question of whether an adult hurt the child instead.

- Many injuries in different healing stages: Bruises change color as they heal—fresh ones are red or purple, older ones turn green, yellow, or brown. Finding bruises of many colors, or fractures (broken bones) that healed at different times, means injuries happened again and again, not just once by accident.
- Waiting too long to come in: A caregiver who caused the injury may delay bringing the child to the doctor out of fear or guilt. The longer the wait for a serious injury like a broken bone or a burn, the more worried the team becomes that something is being hidden.
- Story keeps changing or is different each time: When adults tell different versions of how an injury happened—or change the story between visits—it suggests the true cause is being hidden. Honest accidents usually have clear, steady explanations.
- Bruises in unusual spots (ears, neck, belly, back, buttocks, genitals): Children who play and fall naturally bruise their shins, knees, elbows, and forehead—the bony parts that stick out. Bruises on soft, protected areas like the belly, ears, or inner thighs are much less likely from play and more likely from being hit or grabbed.
- Patterned marks (belt, hand, cord, bite): When a bruise or burn shows the exact shape of an object—like finger marks, a loop from a cord, or teeth marks—it means that object was pressed hard into the skin on purpose, not by accident.
- Child is fearful, withdrawn, or overly watchful of adults: A child who has been hurt at home may become very quiet, avoid eye contact, or flinch when an adult moves quickly. This constant fear and watchfulness happen because the child has learned that adults can be dangerous.
- Poor hygiene, malnutrition, or untreated medical needs: Neglect means caregivers are not meeting the child's basic needs. The child may be very thin (not getting enough food), wearing dirty or torn clothes, have rotten teeth, or have ongoing illnesses that were never treated—all because no adult is taking proper care of them.
- Immersion burns with clear lines or sparing of flexed areas: When a child is forced into hot water, the burn has a sharp, even edge—like a waterline—and may spare the buttocks or feet if the child pulled them up to protect themselves. Accidental splash burns have irregular, scattered edges instead.
- Retinal hemorrhages (bleeding inside the eyes) in an infant: When a baby is shaken violently, the brain moves back and forth inside the skull, and tiny blood vessels in the back of the eyes tear and bleed. This specific pattern of many small bleeds is a strong sign of abusive head trauma (shaken baby syndrome).

Focused exam

- Injury pattern does not match the story—for example, a spiral fracture (twisting break) but the caregiver says the child fell straight down: When the injury and the story do not fit together, you have found a red flag for abuse. Accidents leave predictable injuries. Spiral fractures need twisting force; falling straight does not twist bones.
- Bruises, burns, or fractures in different stages of healing—some fresh, some old, some weeks apart: Multiple injuries at different ages tell you harm happened more than once. Accidents happen once. Repeated injuries across time mean repeated harm.
- High-specificity abuse injuries: looped cord marks, glove-and-stockings immersion burns with sharp waterlines, posterior rib fractures, metaphyseal corner fractures, subdural a pooled collection of blood (hematoma) with retinal hemorrhages: These injuries almost never happen by accident. They are the body's way of showing what was done. Posterior rib fractures mean squeezing. Metaphyseal fractures mean yanking or shaking. Subdural bleeding plus retinal hemorrhages in an infant means abusive head trauma.
- Caregiver story changes each time you ask, or the child is blamed for the injury, or there is unexplained delay in seeking care despite serious injury: The adult's behavior matters as much as the child's body. When stories shift or adults delay getting help, ask why. Abusers often try to cover harm or hope injuries will heal on their own.
- Injuries to soft tissues that do not match developmental stage—bruising on ears, genitals, buttocks, inner thighs, or face in a non-mobile infant: Babies who cannot crawl or walk do not bruise themselves on soft tissues. Accidental bruises happen over bony spots like shins and foreheads once kids start moving. Private areas bruised in an infant is not normal.
- Signs of neglect: severe malnutrition, untreated infections or wounds, failure to thrive without medical explanation, or developmental delay with no follow-up care: Neglect is a form of abuse—when caregivers do not give the child what they need to survive and grow. Neglect can kill just like physical harm can.
- Behavioral red flags: fearfulness around caregivers, age-inappropriate sexual knowledge, extreme withdrawal, overly compliant behavior, or regressive behaviors like bedwetting after being trained: The child's behavior and emotions tell the story when words cannot. These signs mean the child does not feel safe at home.

Diagnostics · every test answers a question

- Skeletal survey (full-body x-rays per AAP protocol: 2 views each limb, skull, spine, ribs, pelvis) → Are there hidden fractures that I cannot see or feel—especially metaphyseal corner fractures, posterior rib fractures, or old healing fractures?

- Non-contrast CT of the head (if any concern for head trauma, altered mental status, vomiting, seizure, or bruising on head/face) → Is there bleeding inside the skull—subdural a pooled collection of blood (hematoma), subarachnoid bleeding (hemorrhage), or brain swelling—that could kill this child or cause permanent brain damage?
- Dilated retinal exam by ophthalmology (mandatory if any suspicion of abusive head trauma) → Are there retinal hemorrhages (bleeding in the back of the eye) that suggest the child was violently shaken?
- Coagulation panel: PT, aPTT, platelet count, and sometimes von Willebrand factor or factor VIII/IX if unexplained bruising → Could a bleeding disorder explain the bruises, or is this pattern truly from harm?
- Liver and pancreatic enzymes: AST, ALT, amylase, lipase (if abdominal trauma suspected—bruising, vomiting, abdominal tenderness) → Is there hidden injury to the liver, spleen, pancreas, or bowel from blunt force to the belly?
- Urinalysis and urine culture (if neglect suspected or unexplained fever, poor hygiene, untreated illness) → Does this child have an untreated infection because no one took care of them?
- Forensic sexual assault examination by trained SANE nurse or child abuse pediatrician (if sexual abuse suspected—genital injury, bleeding, discharge, behavioral signs, or disclosure) → Is there physical evidence of sexual contact or trauma to the genitals, anus, or mouth?
- Screening for sexually transmitted infections: gonorrhea, chlamydia (NAAT), HIV, syphilis, hepatitis B/C if sexual abuse suspected → Did the abuser transmit an infection to the child?

Treatment logic

- Immediate safety assessment and separation from suspected abuser if the child is in danger → Stops ongoing harm and gives the child a safe space where healing can begin. Prevents the next injury or death.
- Detailed documentation with body maps, measurements, and high-resolution photographs of every injury under standard lighting → Creates legal evidence that can be used in court to protect the child. Captures injury patterns that prove abuse—patterns the untrained eye might miss.
- Mandatory report to Child Protective Services and law enforcement immediately (legally required in all 50 U.S. states) → Activates the teams trained to investigate abuse, remove the child from danger if needed, and hold abusers accountable. Connects the child to long-term safety planning and services.

- Hospitalization for medical stabilization, further work-up, and multidisciplinary team coordination (pediatrician, social work, child psychiatry, forensic nurse, child abuse specialist) → Keeps the child medically stable while the team completes the full assessment. Each discipline brings one piece of the picture—medical, legal, emotional, safety.
- Trauma-informed therapy and mental health support for the child (play therapy for young children, cognitive-behavioral therapy for older children) → Helps the child process trauma, rebuild trust, and develop healthy coping skills. Reduces long-term psychological harm and risk of revictimization.
- Medical treatment of injuries: fracture stabilization, wound care, treatment of infections, nutrition support if malnourished → Fixes the immediate physical harm and prevents complications like infection, malunion of fractures, or failure to thrive.
- Education and support for non-offending caregivers (if one parent did not know or was also a victim) → Empowers safe caregivers to protect the child going forward and to understand trauma responses. Strengthens the child's support system.

Medications

No specific medications treat child abuse itself · Child abuse is a social and legal emergency, not a disease treated with drugs

Child abuse means an adult hurts a child on purpose, touches them in a wrong sexual way, says cruel things over and over, or does not give them food, shelter, medical care, or safety. The most important steps are NOT medicines — they are keeping the child safe right now, reporting to the authorities, documenting every injury carefully, and getting experts involved. Medical treatment focuses on injuries the child already has (broken bones, cuts, infections, emotional trauma), not on a pill that stops abuse. The job of the medical team is to recognize the abuse, protect the child, and heal the harm that was done.

1. Step 1: The doctor or nurse notices something wrong — bruises that do not match the story, a pattern of old and new injuries, a child who is very afraid or says someone hurt them, burns in a shape, bite marks, genital injury, a baby with bleeding in the brain and broken ribs when the parent says nothing happened, or a child who is extremely underweight and dirty.
2. Step 2: The medical team takes photos of every injury, writes down exact measurements and colors, draws body maps, and describes what they see in clear words. This record will be evidence.
3. Step 3: The doctor is required by law to report suspected abuse immediately to Child Protective Services and often to the police. Doctors do NOT need to be 100% certain — a reasonable suspicion is enough. Every U.S. state and most countries make this report mandatory.

4. Step 4: If the child is in danger going home, the doctor admits the child to the hospital or CPS places the child with a safe relative or foster family right away.
5. Step 5: Social workers and child-abuse specialists come to the hospital or clinic to help plan safety and support.
6. Step 6: If sexual abuse is suspected, a forensic medical exam is done by a specially trained nurse or doctor (often called a SANE — Sexual Assault Nurse Examiner). This exam collects evidence gently and carefully, treats any sexually transmitted infections, and offers emergency contraception if needed.
7. Step 7: The child may need treatment for specific injuries — casting a broken bone, stitching a cut, antibiotics for an infection, or surgery if internal organs are hurt.
8. Step 8: Mental health support begins as soon as possible. Many abused children develop PTSD (their brain stays on high alert and they have nightmares and flashbacks), depression, or anxiety. Trauma-focused therapy helps them heal emotionally.

SAFETY WATCH

Never send a child back to an unsafe home just because their physical injuries have healed — emotional abuse and neglect are just as dangerous. · Do not interview the child over and over; repeated questioning can be traumatizing and can contaminate their memory. Let trained forensic interviewers do one careful interview on video. · Do not assume a story is true just because a caregiver seems nice or because the family goes to church or has money. Abuse happens in every kind of family. · Do not assume a story is false because the child seems calm, does not cry, or takes time to disclose what happened. Many abused children are too scared or ashamed to tell right away, or have been threatened into silence. · Watch for signs the child is being coached or pressured to change their story. Sometimes non-abusive adults are tricked by an abuser's lies. · If you are a mandated reporter (teacher, doctor, nurse, counselor, coach) and you do NOT report suspected abuse, you can be charged with a crime. Your job is to report — investigators will figure out what really happened.

Red flags

- Injury that cannot happen by accident at that age (e.g. broken bones in a baby who is not yet walking)
- Caregiver story that keeps changing or does not make sense
- Bruises, burns, or fractures in different stages of healing
- Patterned injuries (belt marks, hand prints, bite marks, cord loops)
- Delay in seeking care for a serious injury without a clear reason

Nursing priorities

1. Ensure immediate safety by separating the child from the suspected abuser and placing them in a secure hospital room with supervised visitation only — The first priority is stopping ongoing harm; an abused child returned to an unsafe environment faces high risk of repeated and potentially fatal injury
2. Perform a complete head-to-toe skin examination with the child fully undressed, documenting every bruise, burn, mark, and injury on a body map diagram — Injuries hidden under clothing or in non-accidental patterns (inner thighs, ears, neck, genitals, back) are often missed; thorough documentation creates a permanent medical-legal record
3. Take high-resolution color photographs of all injuries with a ruler for scale, under consistent lighting, from multiple angles before any treatment that might alter appearance — Photographs provide objective evidence for Child Protective Services and law enforcement; bruises and burns change color and fade quickly, so immediate documentation preserves critical forensic evidence
4. File a mandatory report to Child Protective Services and local law enforcement immediately, even if you only suspect abuse — All 50 states legally require healthcare workers to report suspected abuse; failure to report is a crime and leaves the child at risk; reporters have legal immunity when acting in good faith
5. Obtain a skeletal survey (full-body X-rays) for all children under 2 years old with suspected physical abuse, looking for old and new fractures in different healing stages — Babies and toddlers cannot report past abuse; X-rays reveal hidden fractures (ribs, long bones, skull) in various stages of healing that prove repeated trauma over time
6. Assess for life-threatening internal injuries by monitoring vital signs, neurological status, and abdominal signs; obtain CT (computed tomography scan) of the head if altered mental status, vomiting, seizures, or head bruising present — Abusive head trauma (shaking or impact) causes brain bleeding and swelling that can kill within hours; abdominal trauma can rupture organs causing internal bleeding without external bruising
7. Screen for sexual abuse in any child with physical abuse by inspecting the genital and anal areas for bruising, tears, discharge, or other trauma — Physical abuse and sexual abuse often occur together in the same household; genital injuries in young children are almost never accidental and require immediate forensic examination
8. Provide emotional support and age-appropriate explanations to the child without asking leading questions about who hurt them or coaching their responses — Traumatized children need reassurance they are safe and the abuse was not their fault; premature or improper questioning can contaminate forensic interviews and compromise legal cases against perpetrators

Sources

- AAP Clinical Report on Child Abuse Recognition 2023
- CDC Child Abuse and Neglect Prevention
- JAMA Pediatrics guidelines on abusive head trauma
- ACR Appropriateness Criteria for Suspected Physical Abuse—Child 2022

Last reviewed: 2026-07-07