Folate Deficiency Anemia

Without enough folate, the bone marrow makes too few red blood cells that are too big and too fragile to carry oxygen properly.

THE STORY IN ONE SENTENCE

Without enough folate, the bone marrow makes too few red blood cells that are too big and too fragile to carry oxygen properly.

“Folate deficiency anemia happens when your body does not have enough folate (a B vitamin also called B9) to build healthy red blood cells. The bone marrow makes fewer cells that are abnormally large and break apart easily, so oxygen delivery to tissues drops.”

Normal anatomy — the body's job

Folate (vitamin B9) enters the body through food and gets absorbed in the small intestine, then travels in the blood to every cell. Inside the bone marrow, folate works with vitamin B12 and enzymes to build purines and pyrimidines (DNA building blocks) so new red blood cells can divide and mature normally. Healthy red blood cells live about 120 days, carrying oxygen from the lungs to every tissue.

BODY'S JOB

Think of folate as the blueprint ink in a copy shop: every time the bone marrow prints a new red blood cell, it needs fresh ink to copy the DNA instructions clearly so the cell comes out the right size and shape.

1. You eat folate in leafy greens, beans, fortified bread, and orange juice.
2. The upper small intestine (jejunum) absorbs folate and releases it into the bloodstream.
3. Blood carries folate to the bone marrow inside your hip bones, ribs, and skull.
4. Inside marrow stem cells, folate helps enzymes build DNA so the cell can divide into two new cells.
5. Each new red blood cell divides three or four times, shrinking and losing its nucleus to become a mature disc.
6. Mature red blood cells leave the marrow, circulate for 120 days, and carry oxygen to every organ.

What breaks

When folate stores drop too low, the bone marrow cannot make enough DNA for dividing cells. Red blood cell precursors grow larger instead of dividing normally, and many die inside the marrow before finishing. The few that reach the bloodstream are fragile macrocytes (oversized cells) that break apart early, so the blood carries less oxygen.

> **CLINICAL TERM**
> Folate deficiency impairs thymidylate synthesis (the chemical step that builds thymine for DNA), causing megaloblastic erythropoiesis (abnormal red blood cell maturation with giant precursors). Ineffective erythropoiesis (death of precursors in the marrow) and red blood cell destruction (hemolysis) (early red cell destruction) result in macrocytic anemia (low hemoglobin with MCV above 100 fL).

How the body compensates

The body tries to compensate by releasing erythropoietin (EPO, a kidney hormone) to push the marrow harder, but without folate the marrow cannot respond normally. The heart pumps faster and breathing deepens to deliver the limited oxygen available, yet tissue low oxygen (hypoxia) worsens until folate is replaced.

1. {'step': 'Kidneys sense low oxygen in blood and release more EPO', 'detail': 'EPO travels to bone marrow to signal stem cells to make red blood cells faster'}
2. {'step': 'Bone marrow expands and tries to increase red cell production', 'detail': 'Without folate, precursors grow abnormally large and many die before maturing, so output stays low'}
3. {'step': 'Heart rate increases (tachycardia) to circulate existing red cells faster', 'detail': 'Faster pumping tries to deliver oxygen to starving tissues despite fewer carriers'}
4. {'step': 'Breathing rate rises to pull more oxygen into the lungs', 'detail': 'Deeper breaths raise blood oxygen saturation, but fewer red cells mean less oxygen reaches tissues'}
5. {'step': 'Blood flow shifts away from skin and gut toward brain and heart', 'detail': 'Vasoconstriction in less-critical areas preserves oxygen for vital organs'}
6. {'step': 'Over time, compensation fails and chronic low oxygen (hypoxia) damages tissues', 'detail': 'Nerves, heart muscle, and rapidly dividing gut cells suffer most when oxygen and folate remain low'}

Domino effect

1. Folate stores in the liver drop below 5–10 mg (normal is 10–20 mg) after weeks of poor intake or malabsorption — Body stores last only 3–4 months; daily needs are 400 µg but intake or absorption is insufficient
2. Serum folate falls below 3 ng/mL, then red cell folate drops below 140 ng/mL — Blood levels fall first, then the folate inside existing red cells declines as stores empty
3. Bone marrow erythroblasts pause during DNA synthesis and grow abnormally large (megaloblasts) — Thymidylate synthase cannot make thymine without folate cofactors, so cells arrest in S phase and cytoplasm keeps expanding
4. Ineffective erythropoiesis causes many precursors to die in the marrow (intramedullary red blood cell destruction (hemolysis)) — Abnormal nuclear maturation triggers apoptosis; only a fraction of megaloblasts mature and enter blood
5. Hemoglobin and red cell count fall; MCV rises above 100 fL as macrocytes enter circulation — Fewer, larger, fragile cells cannot carry enough oxygen, and the spleen destroys many early
6. Tissue low oxygen (hypoxia) causes fatigue, pale skin (pallor), a fast heart rate (tachycardia), and shortness of breath — Muscles, brain, and heart receive insufficient oxygen; lactic acid builds up during normal activity
7. Rapidly dividing cells suffer next: tongue becomes sore and red (glossitis), diarrhea begins, and neural tube defects occur in fetuses — Gut mucosa and developing nerves need folate for DNA synthesis; deficiency halts cell division in these tissues
8. Severe untreated anemia leads to high-output heart failure, neurologic damage (peripheral neuropathy, rarely subacute combined degeneration if B12 also low), and death — Chronic low oxygen (hypoxia) exhausts the heart; folate deficiency impairs myelin repair, especially when combined B12 deficiency is missed

Symptoms · what the patient feels, and why

- Extreme tiredness that does not improve with rest: Low folate stops cells from making DNA, so the body cannot produce enough red blood cells to carry oxygen to muscles and organs
- Pale or yellowish skin and eyes: Without enough folate, the bone marrow makes large fragile red blood cells (macrocytes) that break apart easily, releasing bilirubin (a yellow pigment)
- Shortness of breath during usual activities: Fewer red blood cells mean less oxygen delivered to the lungs and heart, forcing the body to breathe faster to compensate

- Sore red tongue (glossitis): Folate is essential for rapidly dividing cells in the mouth and tongue; deficiency causes these cells to die off, leaving the tongue smooth and inflamed
- Rapid heartbeat or chest pain: The heart pumps faster to circulate the limited oxygen-carrying red blood cells throughout the body, straining the heart muscle
- Tingling or numbness in hands and feet: Severe folate deficiency can occasionally affect nerve cells, though this is far more common with vitamin B12 deficiency
- Diarrhea and poor appetite: The intestinal lining sheds cells rapidly and depends on folate for renewal; deficiency slows this process and damages the gut lining
- Confusion or difficulty concentrating: The brain requires steady oxygen supply; anemia reduces oxygen delivery to brain cells, slowing mental processing

Focused exam

- Pale conjunctivae (inside lower eyelids) and pale nail beds: These areas are normally pink because of blood vessels near the surface; pale skin (pallor) signals low hemoglobin and confirms anemia
- Smooth red tongue without the normal bumpy surface (atrophic glossitis): Loss of the small projections (papillae) on the tongue is a specific sign of folate or vitamin B12 deficiency affecting rapidly dividing mucosal cells
- Yellow tint to the skin and whites of the eyes (jaundice): Fragile macrocytes break apart in the blood vessels (hemolysis), releasing bilirubin and suggesting more severe or longstanding folate deficiency
- Rapid resting heart rate above 100 beats per minute (tachycardia): The heart compensates for low oxygen-carrying capacity by beating faster; persistent a fast heart rate (tachycardia) indicates the anemia is causing cardiovascular stress
- Decreased vibratory sense or light touch in the feet: Neurologic changes raise concern for concurrent vitamin B12 deficiency (which causes nerve damage); folate alone rarely causes peripheral neuropathy
- An enlarged spleen (Splenomegaly) (enlarged spleen felt below the left rib cage): The spleen may enlarge as it removes defective macrocytes from circulation; an enlarged spleen (splenomegaly) can also point to other causes of red blood cell destruction (hemolysis) or bone marrow disorders

Diagnostics · every test answers a question

- Complete blood count (CBC) with red blood cell indices → Does the patient have anemia, and are the red blood cells larger than normal (macrocytic)?
- Peripheral blood smear → What do the red blood cells look like under the microscope?

- Serum folate level → Is the blood folate concentration low?
- Red blood cell (RBC) folate level → What are the long-term folate stores inside red blood cells?
- Serum vitamin B12 level → Could vitamin B12 deficiency be causing the macrocytic anemia instead of or in addition to folate deficiency?
- Serum methylmalonic acid (MMA) and homocysteine → If B12 or folate levels are borderline, are the body's metabolic pathways actually impaired?
- Reticulocyte count → Is the bone marrow trying to make new red blood cells?
- Lactate dehydrogenase (LDH) and indirect bilirubin → Are red blood cells breaking apart inside the blood vessels (hemolysis)?

Treatment logic

- Oral folic acid supplementation → Restores folate levels within days, allowing the bone marrow to resume normal DNA synthesis and produce healthy red blood cells; hemoglobin typically rises within two to four weeks
- Identify and treat the underlying cause (stop interfering medications, treat celiac disease or alcoholism, adjust dialysis) → Prevents recurrence by removing the reason folate became depleted; for example, stopping methotrexate or treating celiac disease allows normal folate absorption
- Vitamin B12 supplementation if B12 deficiency is also present → Protects nerves and prevents subacute combined degeneration (spinal cord damage); both deficiencies must be treated together because folate alone can worfully mask B12 deficiency
- Dietary counseling on folate-rich foods (spinach, broccoli, lentils, fortified cereals and breads) → Builds sustainable folate intake to maintain normal stores after initial supplementation; fortified grains provide synthetic folic acid that is absorbed more reliably than natural folate
- Preconception and prenatal folic acid (400–800 mcg daily for all women of childbearing age) → Prevents neural tube defects in the developing fetus during the first month of pregnancy, before most women know they are pregnant; reduces the risk of spina bifida and anencephaly by up to seventy percent
- Red blood cell transfusion if hemoglobin is critically low (below 7 g/dL) or patient has severe heart symptoms → Immediately raises hemoglobin and oxygen-carrying capacity, stabilizing patients with dangerous anemia or signs of heart failure (chest pain, severe shortness of breath)

Medications

folic acid · Water-soluble B vitamin (vitamin B9)

Folic acid is a synthetic form of folate that the body converts into tetrahydrofolate (the active form). Tetrahydrofolate carries one-carbon units needed to build purines and thymidine, which are the building blocks of DNA. Without adequate folate, cells cannot divide properly, especially rapidly dividing cells like red blood cell precursors in the bone marrow.

1. Oral folic acid is absorbed in the proximal small intestine (jejunum) by specific folate transporters
2. The intestinal cells and liver convert folic acid to dihydrofolate and then to tetrahydrofolate using the enzyme dihydrofolate reductase
3. Tetrahydrofolate accepts one-carbon groups to become methyltetrahydrofolate and other cofactor forms
4. These cofactors donate one-carbon units for synthesis of thymidine (DNA base) and purines (adenine and guanine)
5. Bone marrow cells use these DNA building blocks to divide and mature into normal-sized red blood cells instead of large fragile macrocytes

> **SAFETY WATCH**
>
> Can mask vitamin B12 deficiency by improving anemia while allowing nerve damage to progress; always check B12 before starting folate · High doses (above 1 mg daily) may hide pernicious anemia on blood tests · Generally very safe; nausea or abdominal discomfort is rare · Interferes with some seizure medications; phenytoin and phenobarbital levels may drop

cyanocobalamin (vitamin B12) · Water-soluble B vitamin (cobalamin)

Cyanocobalamin is a form of vitamin B12 that the body converts into methylcobalamin and adenosylcobalamin, the active forms. These cofactors work with folate in DNA synthesis and are also essential for myelin (nerve coating) formation. B12 must be given along with folate if both deficiencies exist, because treating folate alone will not protect the nerves.

1. Cyanocobalamin is given by mouth or injection and absorbed in the ileum (lower small intestine) with the help of intrinsic factor from the stomach
2. B12 travels in the blood attached to carrier proteins called transcobalamins
3. Inside cells, B12 is converted to methylcobalamin and adenosylcobalamin
4. Methylcobalamin works with folate to convert homocysteine to methionine, regenerating tetrahydrofolate for DNA synthesis

5. Adenosylcobalamin helps convert methylmalonyl-CoA to succinyl-CoA, supporting myelin production and nerve function

SAFETY WATCH

Oral B12 may not work if intrinsic factor is missing (pernicious anemia) or the ileum is damaged; intramuscular injections bypass this problem · Low potassium (Hypokalemia) (low potassium) can occur in the first week as new blood cells are rapidly produced; monitor potassium levels · Very safe with few side effects; injection-site soreness is common with intramuscular shots · Rare allergic reactions to injectable forms

leucovorin (folinic acid) · Reduced folate (active form of folic acid)

Leucovorin is already in the tetrahydrofolate form, so it does not require conversion by dihydrofolate reductase. It is used to rescue normal cells during high-dose methotrexate chemotherapy, because methotrexate blocks dihydrofolate reductase and stops all folate activity. Leucovorin bypasses the blocked enzyme and allows healthy cells to make DNA while the cancer cells remain suppressed.

1. Leucovorin is given intravenously or orally and enters cells directly as folinic acid
2. It is converted into active tetrahydrofolate cofactors without needing dihydrofolate reductase
3. These cofactors supply one-carbon units for DNA synthesis in normal cells
4. Cancer cells affected by methotrexate cannot use leucovorin as effectively because methotrexate also blocks other folate pathways inside cancer cells
5. Normal bone marrow and gut lining recover, preventing dangerous toxicity from methotrexate

SAFETY WATCH

Must be given within 24 to 42 hours after high-dose methotrexate to prevent severe toxicity · Timing and dose are critical; too much leucovorin can reduce methotrexate effectiveness against cancer · Can cause allergic reactions including rash and wheezing · Not interchangeable with folic acid for routine supplementation

Dietary modification (folate-rich foods) · Nutritional intervention

Folate occurs naturally in foods as polyglutamate folate, which is converted to monoglutamate folate in the intestine before absorption. Green leafy vegetables (spinach, kale), legumes (lentils, black beans), and fortified grains provide high folate content. The United States

mandates folic acid fortification of enriched grain products, which has significantly reduced folate deficiency and neural tube defects.

1. Folate polyglutamates in food are broken down to monoglutamates by intestinal enzymes called conjugases
2. Monoglutamate folate is absorbed across the jejunal wall by folate transporters
3. Absorbed folate travels to the liver where it is stored or converted to active forms
4. Tissues throughout the body take up folate from the blood and add glutamate chains back on for storage
5. Stored folate is released as needed to support DNA synthesis in dividing cells

SAFETY WATCH

Natural folate is less bioavailable than synthetic folic acid (about fifty percent absorption vs. eighty-five percent) · Alcohol blocks folate absorption even with adequate dietary intake · Overcooking vegetables destroys folate; steaming is better than boiling · Fortified foods provide the most reliable dietary source in the United States

Packed red blood cell transfusion · Blood product replacement therapy

Transfusion delivers donor red blood cells that immediately carry oxygen to tissues. It is reserved for patients with dangerously low hemoglobin (usually below 7 g/dL) or those with symptoms of heart failure or severe low oxygen (hypoxia) (low oxygen). Transfusion is a bridge therapy while folate supplementation takes time to stimulate the bone marrow to produce new cells.

1. Donor blood is cross-matched to ensure ABO and Rh compatibility with the patient
2. Packed red blood cells are infused intravenously over one to four hours
3. Donor cells circulate immediately and begin delivering oxygen to oxygen-starved (hypoxic) tissues
4. Each unit of packed red blood cells raises hemoglobin by about 1 g/dL in an average adult
5. Temporary improvement in oxygen delivery stabilizes the patient while folate therapy restores bone marrow function over weeks

SAFETY WATCH

Risk of transfusion reactions including fever, allergic reactions, and rare destroying red blood cells (hemolytic) reactions if blood is mismatched · Volume overload and heart failure can occur if transfused too rapidly, especially in elderly patients · Infection transmission risk is very low with modern screening but not zero · Iron overload can develop with repeated transfusions over time

Red flags

- New confusion, difficulty walking, or loss of balance (may indicate B12 deficiency with spinal cord damage, not just folate deficiency)
- Chest pain or shortness of breath at rest (signals severe anemia causing heart strain or heart failure)
- Pregnancy with folate deficiency (high risk of neural tube defects in the developing fetus)
- Hemoglobin below 7 g/dL or signs of shock (pale, sweaty, rapid pulse, low blood pressure)

Nursing priorities

1. Assess neurological status for signs of folate-related nerve damage — Folate deficiency can cause irritability, forgetfulness, and cognitive changes but does not cause the permanent nerve damage (demyelination) seen with B12 deficiency, so early detection of changes helps guide treatment urgency
2. Monitor for signs of tissue low oxygen (hypoxia) from severe anemia — Folate deficiency causes macrocytic anemia (large red blood cells that do not carry oxygen well), leading to chest pain, shortness of breath, dizziness, and fatigue that can progress to heart failure if hemoglobin drops very low
3. Identify and address dietary causes of folate deficiency — Poor intake of leafy greens, fruits, and fortified grains is the most common cause in developed countries, and alcohol use disorder blocks folate absorption and storage, so nutrition education and social support are essential
4. Screen pregnant or reproductive-age women immediately — Folate deficiency in the first trimester causes neural tube defects (serious brain and spine malformations) in the developing fetus, so preconception and early pregnancy supplementation with 400-800 mcg daily is critical
5. Review medications that interfere with folate metabolism — Methotrexate (chemotherapy and autoimmune drug), trimethoprim (antibiotic), phenytoin (seizure medication), and sulfasalazine (inflammatory bowel disease drug) block folate pathways, so patients on these need higher supplementation doses
6. Differentiate folate deficiency from vitamin B12 deficiency before treatment — Both cause identical large red blood cells on blood smear, but giving folate to a B12-deficient patient can mask the anemia while allowing permanent nerve damage to progress, so serum B12 and folate levels must be checked together
7. Educate on proper folic acid supplementation and expected response time — Oral folic acid 1-5 mg daily corrects deficiency within 2-4 weeks in most patients, reticulocyte count (young red blood cells) rises in 3-5 days showing bone marrow recovery, and hemoglobin normalizes in 1-2 months

8. Assess for underlying malabsorption disorders if dietary intake is adequate — Celiac disease (gluten-triggered intestinal damage), inflammatory bowel disease, and tropical sprue (infection causing malabsorption) prevent folate absorption in the jejunum (upper small intestine), requiring treatment of the root cause plus supplementation

Sources

- American Society of Hematology (ASH) guidelines on the evaluation and management of macrocytic anemia and megaloblastic disorders (2021)
- Centers for Disease Control and Prevention (CDC) recommendations for folic acid supplementation for prevention of neural tube defects (2020)
- National Institutes of Health (NIH) Office of Dietary Supplements fact sheet on folate for health professionals (2022)
- American College of Obstetricians and Gynecologists (ACOG) Practice Bulletin on neural tube defects and periconceptional folic acid supplementation (2017, reaffirmed 2021)

Last reviewed: 2026-08-29