

Marfan Syndrome

Marfan syndrome is a connective tissue disorder caused by defective fibrillin-1 protein, leading to fragile elastic fibers in the aorta, eyes, and skeleton that stretch and tear under normal body stress.

THE STORY IN ONE SENTENCE

Marfan syndrome is a connective tissue disorder caused by defective fibrillin-1 protein, leading to fragile elastic fibers in the aorta, eyes, and skeleton that stretch and tear under normal body stress.

“Marfan syndrome is a genetic condition where the body makes faulty connective tissue. The main protein that gives strength and elasticity to blood vessels, eyes, and bones does not work properly. This causes dangerous stretching of the aorta, lens problems in the eyes, and tall stature with long limbs.”

Normal anatomy — the body's job

Fibrillin-1 is a structural protein that forms microfibrils (tiny fibers) in the extracellular matrix (the scaffolding between cells). These microfibrils give strength and elasticity to connective tissues throughout the body, especially in blood vessel walls, the lens suspensory ligaments in the eye, and the periosteum (bone covering) and ligaments. Fibrillin-1 also stores and regulates transforming growth factor beta (TGF- β), a signaling molecule that controls cell growth and tissue repair.

BODY'S JOB

Fibrillin-1 works like the elastic waistband in sweatpants—it stretches when you need it to but snaps back to shape and holds everything in place securely.

1. The FBN1 gene on chromosome 15 provides instructions to make fibrillin-1 protein
2. Cells called fibroblasts make and secrete fibrillin-1 into the extracellular matrix
3. Fibrillin-1 molecules link together to form long microfibrils that create a mesh network
4. This microfibril network combines with elastin protein to form elastic fibers in vessel walls, ligaments, and other tissues

5. The elastic fibers stretch during movement or blood pressure pulses, then recoil back to original shape
6. Fibrillin-1 microfibrils also bind and store TGF- β , keeping its activity controlled and balanced

What breaks

In Marfan syndrome, a mutation (spelling mistake) in the FBN1 gene causes the body to make abnormal fibrillin-1 protein or not enough of it. Without properly working fibrillin-1, the microfibrils are weak and break easily. Tissues that need to stretch and recoil—like the aorta wall, the eye lens ligaments, and bone ligaments—become fragile and overstretch under normal daily stress.

CLINICAL TERM

Most Marfan cases result from missense or nonsense mutations in FBN1 that produce structurally defective fibrillin-1 or cause haploinsufficiency (only one working gene copy making half the normal protein amount). Defective microfibrils cannot properly sequester TGF- β , leading to excessive TGF- β signaling that weakens the extracellular matrix and promotes proteolytic degradation (enzyme breakdown) of elastic fibers.

How the body compensates

The body has limited ability to compensate for defective fibrillin-1. Fibroblasts may increase production of other extracellular matrix proteins like fibronectin to try to reinforce weak tissues, but these cannot fully replace fibrillin-1 function. The left ventricle may hypertrophy (thicken) to maintain blood pressure when the aorta dilates, but this adds more stress to the already weak aortic wall.

1. {'step': 'Increased extracellular matrix protein synthesis', 'detail': 'Fibroblasts sense tissue weakness and make more fibronectin, collagen, and other structural proteins to try to strengthen connective tissue scaffolding'}
2. {'step': 'Left ventricular hypertrophy in response to aortic dilation', 'detail': 'As the aorta stretches and loses elasticity, the left ventricle must generate higher pressure to eject blood, causing heart muscle to thicken over time'}
3. {'step': 'Upregulation of matrix metalloproteinase inhibitors', 'detail': 'Some tissues release tissue inhibitors of metalloproteinases (TIMPs) to slow down enzyme breakdown of remaining elastic fibers'}
4. {'step': 'Increased collagen deposition in vessel walls', 'detail': 'Smooth muscle cells deposit more rigid collagen in the aortic wall to prevent rupture, but this stiffens the artery and raises blood pressure'}

5. {'step': 'Reflexive vasoconstriction to maintain blood flow (perfusion) pressure', 'detail': 'When the dilated aorta reduces effective forward flow, baroreceptors (pressure sensors) trigger sympathetic nervous system activation to constrict peripheral vessels'}
6. {'step': 'Myopia development from elongated eye globe', 'detail': 'The eye attempts to focus light on the retina despite abnormal lens position by elongating the eyeball, which causes nearsightedness'}

Domino effect

1. Defective fibrillin-1 causes progressive aortic root dilation — Each heartbeat stretches the weak aortic wall slightly more, and the vessel loses the ability to recoil, so diameter increases gradually over years
2. Aortic regurgitation develops as the root dilates — When the aortic root widens, the three valve leaflets cannot meet properly in the center, allowing blood to leak backward into the left ventricle during diastole (relaxation phase)
3. Left ventricular volume overload and eccentric hypertrophy — The ventricle must pump both the regurgitant volume and the normal stroke volume, so it dilates and thickens to handle the extra load
4. Increased wall stress accelerates aortic dilation — According to Laplace's law, wall tension equals pressure times radius, so as the aorta gets wider, wall stress increases exponentially, creating a vicious cycle of worsening dilation
5. Aortic dissection occurs when intimal tear allows blood into the vessel wall — Extreme wall stress causes the inner aortic layer (intima) to tear, and high-pressure blood cleaves through the weakened media layer, separating wall layers
6. Blood flow to branch vessels is compromised by the dissection flap — The false lumen (dissection space) can compress the true lumen or occlude branch vessel origins, cutting off blood to the brain, kidneys, intestines, or limbs
7. Cardiac tamponade (the heart squeezed by fluid around it) develops if dissection ruptures into the pericardium — Blood fills the pericardial sac around the heart, preventing ventricular filling and causing cardiogenic shock
8. Death from aortic rupture, tamponade, or end-organ malperfusion — Sudden massive bleeding (hemorrhage) into the chest or abdomen causes exsanguination (bleeding to death), or critical organs fail from lack of oxygen delivery

Symptoms · what the patient feels, and why

- Tall stature with long arms, legs, and fingers: Defective fibrillin-1 protein (a connective tissue building block) allows bones to grow longer than usual because growth plates stay open longer and cartilage is less restrained

- Chest pain or pressure: Weakened aortic wall stretches and tears, or mitral valve leaflets prolapse backward into the left atrium causing starved blood flow (ischemia) or dissection
- Vision problems or lens dislocation: Zonules (tiny fibers made of fibrillin that hold the lens in place) stretch or break, letting the lens move out of position
- Shortness of breath with exertion: Mitral or aortic valve regurgitation (backward leak) forces the heart to pump the same blood twice, or spontaneous pneumothorax (lung collapse) reduces oxygen exchange
- Pectus deformity (sunken or protruding breastbone): Weak costal cartilage (rib connective tissue) and rib overgrowth allow the sternum to cave inward or push outward
- Joint hypermobility or frequent dislocations: Ligaments (bands that hold bones together) have abnormal fibrillin so they overstretch and fail to stabilize joints
- Sudden severe back or abdominal pain: Aortic dissection (tear in the aortic inner layer) lets blood split the wall layers apart, or aortic aneurysm (balloon-like bulge) leaks or ruptures
- Skin stretch marks not related to weight change: Dermis (deep skin layer) has defective fibrillin so collagen bundles separate under normal tension, leaving visible red or white lines

Focused exam

- Arm span-to-height ratio greater than 1.05 and upper-to-lower body segment ratio less than 0.86: Disproportionate long limbs meet skeletal criteria in the revised Ghent nosology (2010 diagnostic system) and suggest growth dysregulation from fibrillin deficiency
- Positive wrist sign (thumb overlaps fifth finger around opposite wrist) or thumb sign (thumb tip extends beyond ulnar border of palm): These signs indicate arachnodactyly (spider-like fingers) and score as skeletal features; they reflect long metacarpals and ligament laxity
- Pectus carinatum or pectus excavatum requiring surgery: Severe chest deformity meets skeletal system criteria and may compress the heart or lungs, reducing cardiac output or lung volumes
- Mid-to-late systolic click and late systolic murmur at apex: Indicates mitral valve prolapse (leaflet bulges into left atrium) with regurgitation; severe regurgitation causes left atrial enlargement and atrial fibrillation (an irregular, quivering heartbeat) risk
- Early diastolic decrescendo murmur at left sternal border, bounding pulses, wide pulse pressure: Signs of aortic regurgitation (valve does not close fully); chronic severe regurgitation enlarges the left ventricle and leads to heart failure
- Blood pressure difference greater than 20 mmHg between arms, or new diastolic murmur with severe chest pain: Suggests acute aortic dissection (inner aortic layer tears); type A dissection (involves ascending aorta) requires emergency surgery within hours

Diagnostics · every test answers a question

- Transthoracic echocardiography with aortic root Z-score → Is the aortic root diameter enlarged for the patient's age and body surface area, and is there valve regurgitation?
- CT angiography or MR angiography of the entire aorta → Are there aneurysms (abnormal dilations) or dissection flaps in the thoracic or abdominal aorta?
- Slit-lamp ophthalmologic examination → Is the lens displaced (ectopia lentis), and are there other fibrillin-related eye findings?
- Genetic testing for FBN1 mutation → Is there a pathogenic variant in the FBN1 gene that encodes fibrillin-1 protein?
- Skeletal radiographs (spine, chest) → What is the severity of scoliosis, kyphosis, or pectus deformity, and is surgical correction needed?
- Electrocardiogram and 24-hour Holter monitor → Are there arrhythmias such as atrial fibrillation (an irregular, quivering heartbeat) or ventricular ectopy that increase stroke or sudden death risk?
- Dural ectasia imaging (lumbar MRI) → Is the dural sac (membrane surrounding spinal cord) ballooned outward, causing back pain or nerve symptoms?
- Pulmonary function tests → Is there restrictive lung disease from chest-wall deformity or a history of pneumothorax reducing lung capacity?

Treatment logic

- Beta-blocker therapy (atenolol or metoprolol) → Slows heart rate and reduces aortic wall stress (force per unit area) by lowering blood pressure and the speed of ventricular contraction, slowing aortic root growth by approximately 0.1 cm/year
- Preventive (Prophylactic) aortic root replacement surgery → Replaces the dilated ascending aorta and aortic valve with a graft (tube) and mechanical or bioprosthetic valve, preventing rupture or dissection; 30-year survival exceeds 80% post-surgery
- Angiotensin II receptor blocker (ARB) therapy (losartan) → Blocks TGF- β (transforming growth factor beta) signaling that drives aortic wall remodeling and smooth muscle loss, reducing aortic root growth rate in some studies by 0.1–0.2 cm/year
- Competitive and contact sports restriction → Avoids sudden blood pressure spikes and chest trauma that can trigger dissection or worsen valve regurgitation; dynamic low-intensity exercise (walking, swimming) is safe and improves fitness
- Mitral or aortic valve repair or replacement → Restores competent valve closure, reduces left ventricular volume overload, and prevents progression to heart failure or pulmonary high blood pressure (hypertension); mitral repair has lower long-term complications than replacement

- Annual cardiovascular imaging surveillance → Detects progressive aortic dilation before rupture or dissection; ascending aorta diameter and growth rate determine surgical timing per ACC/AHA guidelines (surgery at ≥ 5.0 cm or ≥ 4.5 cm with risk factors)

Medications

Atenolol · Cardioselective beta-1 adrenergic receptor blocker

Atenolol blocks beta-1 receptors on heart cells so adrenaline (epinephrine) cannot bind and speed up the heart. Slower heart rate and weaker contractions lower blood pressure and reduce the force hitting the aortic wall with each heartbeat. Less mechanical stress on the aorta slows aneurysm growth.

1. Atenolol binds to beta-1 receptors on the heart cell surface
2. Blocks cyclic AMP production inside the cell
3. Reduces calcium entry into myocytes, weakening contraction strength
4. Slows electrical firing in the sinoatrial node, lowering heart rate
5. Decreases cardiac output and systemic blood pressure, reducing aortic wall tension

SAFETY WATCH

A slow heart rate (Bradycardia) (heart rate < 50 bpm) and low blood pressure (hypotension) · Fatigue and exercise intolerance from reduced cardiac output · Bronchospasm in patients with asthma or COPD (chronic obstructive pulmonary disease) · Avoid abrupt discontinuation (can cause rebound a fast heart rate (tachycardia) and high blood pressure (hypertension))

Losartan · Angiotensin II receptor blocker (ARB)

Losartan blocks angiotensin II from binding to AT1 receptors on blood vessels and the heart. This lowers blood pressure and also reduces TGF- β (a signaling protein that damages aortic tissue in Marfan syndrome). Less TGF- β means less breakdown of elastin (stretchy fibers) and collagen in the aortic wall.

1. Losartan binds to AT1 receptors, blocking angiotensin II attachment
2. Prevents vasoconstriction (blood vessel narrowing) and aldosterone release
3. Reduces TGF- β activation in the aortic wall
4. Decreases matrix metalloproteinase (enzymes that break down elastin and collagen) activity
5. Slows aortic dilation and improves wall structural integrity

SAFETY WATCH

High potassium (Hyperkalemia) (high potassium) especially with renal insufficiency or potassium supplements · Low blood pressure (Hypotension) and dizziness, particularly in volume-depleted patients · Contraindicated in pregnancy (causes fetal renal failure and death) · Monitor creatinine; may worsen renal function in bilateral renal artery narrowing (stenosis)

Warfarin · Vitamin K antagonist anticoagulant

Warfarin blocks vitamin K epoxide reductase, the enzyme that recycles vitamin K in the liver. Without active vitamin K, the liver cannot make clotting factors II, VII, IX, and X. Thinner blood prevents clots from forming on mechanical heart valves (which all Marfan patients with valve replacements may have).

1. Warfarin inhibits vitamin K epoxide reductase in hepatocytes
2. Depletes active vitamin K cofactor needed for gamma-carboxylation
3. Clotting factors II, VII, IX, X are synthesized but remain inactive (non-carboxylated)
4. Coagulation cascade cannot form fibrin clots efficiently
5. INR (international normalized ratio) rises to therapeutic range 2.0–3.0 for valve patients

SAFETY WATCH

Narrow therapeutic window; requires frequent INR monitoring · Major bleeding risk (intracranial bleeding (hemorrhage), gastrointestinal bleed) if INR >4 · Many drug and food interactions (vitamin K in green leafy vegetables reduces effect) · Contraindicated in pregnancy (teratogenic, causes fetal bleeding)

Celiprolol · Beta-1 selective blocker with beta-2 agonist activity

Celiprolol blocks beta-1 receptors to slow the heart and lower blood pressure, like atenolol. It also mildly stimulates beta-2 receptors in blood vessels, causing vasodilation (widening). This combination may reduce aortic stiffness more than pure beta-blockers. Some European studies show benefit in vascular Ehlers-Danlos, and it is sometimes used off-label in Marfan.

1. Celiprolol blocks beta-1 receptors, reducing heart rate and contractility
2. Simultaneously activates beta-2 receptors on peripheral vessels
3. Causes mild vasodilation, lowering systemic vascular resistance
4. Reduces pulse wave velocity (stiffness measure) in the aorta
5. Lowers aortic wall stress through combined heart-rate and blood-pressure effects

SAFETY WATCH

Similar a slow heart rate (bradycardia) and low blood pressure (hypotension) risks as other beta-blockers · Beta-2 agonism may worsen asthma less than non-selective agents, but caution still needed · Drug availability limited outside Europe · Monitor for fatigue and dizziness

Pneumococcal and influenza vaccines · Preventive (Prophylactic) immunization (non-pharmacologic prevention)

These vaccines train the immune system to recognize *Streptococcus pneumoniae* and influenza virus. After vaccination, B cells produce antibodies that bind to bacterial capsule or viral surface proteins. If the patient later encounters the real pathogen, preformed antibodies quickly neutralize it, preventing pneumonia. Marfan patients with chest-wall deformities or prior pneumothorax have reduced lung reserve, so preventing respiratory infections is critical.

1. Vaccine antigens (protein or polysaccharide fragments) enter the bloodstream
2. Antigen-presenting cells carry fragments to lymph nodes
3. B lymphocytes recognize antigens and mature into plasma cells
4. Plasma cells secrete IgG antibodies specific to pneumococcus or influenza
5. Memory B cells remain in circulation for rapid response on re-exposure

SAFETY WATCH

Injection-site pain and low-grade fever for 1–2 days · Rarely, allergic reaction to vaccine components (egg protein in some flu vaccines) · Pneumococcal conjugate (PCV13) and polysaccharide (PPSV23) vaccines recommended; timing per CDC schedule · Annual influenza vaccine needed due to viral strain changes

Genetic counseling and prenatal diagnosis · Non-pharmacologic reproductive intervention

Genetic counselors explain the 50% autosomal dominant inheritance risk, review family pedigree, and offer preimplantation genetic testing (PGT) or prenatal diagnosis. PGT involves in-vitro fertilization, embryo biopsy at 5 days, and FBN1 gene sequencing before uterine transfer. Alternatively, chorionic villus sampling (CVS) at 10–13 weeks or amniocentesis at 15–20 weeks can test fetal DNA. These methods allow parents to make informed reproductive choices.

1. Genetic counselor obtains detailed family and medical history
2. FBN1 sequencing identifies pathogenic variant in affected parent
3. For PGT: embryos biopsied and tested; only unaffected embryos transferred

4. For CVS or amniocentesis: fetal cells analyzed for FBN1 mutation
5. Parents receive risk assessment and decision-support counseling

SAFETY WATCH

CVS carries ~0.3% miscarriage risk; amniocentesis ~0.1% · PGT requires hormonal stimulation and invasive egg retrieval · Ethical, financial, and emotional considerations require multidisciplinary support · Negative prenatal test does not rule out de novo mutations in later offspring

Red flags

- Sudden severe chest or back pain (possible aortic dissection—activate code team and order STAT CT angiography)
- New heart murmur with fainting (syncope) or heart failure symptoms (acute valve rupture or severe regurgitation—urgent cardiology consult)
- Blood pressure difference >20 mmHg between arms (dissection extending into brachiocephalic vessels—immediate imaging)
- Shortness of breath with unilateral absent breath sounds (spontaneous pneumothorax—needs chest X-ray and possible chest tube)

Nursing priorities

1. Monitor for aortic dissection (tear in the large blood vessel leaving the heart) and rupture — Marfan syndrome weakens the aortic wall, making it prone to life-threatening tears. Early detection through regular blood pressure checks and assessment of chest or back pain can save lives.
2. Assess cardiovascular status including heart sounds and murmurs — Mitral valve prolapse (backward bulging of a heart valve) and aortic regurgitation (leaky aortic valve) are common in Marfan syndrome and can lead to heart failure if not monitored and treated.
3. Monitor for spontaneous pneumothorax (collapsed lung) — Patients with Marfan syndrome have weak connective tissue in the lungs, increasing risk for sudden lung collapse. Watch for sudden chest pain, shortness of breath, or decreased oxygen levels.
4. Screen for vision changes and lens dislocation — Ectopia lentis (displaced lens in the eye) occurs in more than half of Marfan patients and can cause severe vision problems. Early referral to ophthalmology preserves vision.
5. Educate about activity restrictions and avoidance of contact sports — High-intensity exercise and contact sports increase aortic wall stress and risk of dissection. Patients need clear guidance on safe physical activities to prevent sudden death.

6. Administer beta-blockers as prescribed and monitor heart rate and blood pressure — Beta-blockers slow heart rate and reduce force of heart contractions, decreasing stress on the aorta. This slows aortic root dilation and reduces dissection risk.
7. Provide genetic counseling education and family screening referrals — Marfan syndrome is autosomal dominant (passed from parent to child with 50% chance). Family members need screening because early diagnosis and treatment improve outcomes.
8. Assess for scoliosis (curved spine) and joint hypermobility complications — Skeletal abnormalities cause chronic pain and functional limitations. Early orthopedic intervention and physical therapy can prevent progression and improve quality of life.

Sources

- American Heart Association (AHA) 2022 Guidelines for Management of Patients with Thoracic Aortic Disease
- American College of Medical Genetics and Genomics (ACMG) 2021 Practice Guidelines for Genetic Evaluation of Marfan Syndrome
- National Marfan Foundation 2023 Clinical Guidelines for Diagnosis and Management
- European Society of Cardiology (ESC) 2020 Guidelines on the Diagnosis and Treatment of Aortic Diseases

Last reviewed: 2026-08-29