

Personality Disorder and Crisis Stabilization

Her moods, sense of self, and relationships swing wildly from moment to moment.

THE STORY IN ONE SENTENCE

Her moods, sense of self, and relationships swing wildly from moment to moment.

“Borderline personality disorder in crisis stems from a vulnerable brain emotion-regulation circuit that was hurt by early rejection, invalidation, or trauma.”

Normal anatomy — the body's job

A healthy brain keeps your emotions in a manageable range, your sense of self steady, and your view of other people balanced—even when stress or disappointment happens.

BODY'S JOB

The prefrontal cortex (the front part of the brain that plans and pumps the brakes on feelings) talks to the amygdala (the alarm center deep in the brain that flags danger). The prefrontal cortex calms the amygdala down when a threat turns out to be small. Serotonin (a brain messenger that steadies mood) and other neurotransmitters keep this conversation smooth. The result: you feel upset for a bit, then settle. You know who you are. You trust that one argument won't destroy a relationship.

1. Stress or disappointment arrives (someone says no, you make a mistake, a friend cancels plans).
2. Your amygdala fires an alarm: 'This is bad!'
3. Your prefrontal cortex checks the facts: 'Is this really dangerous, or just uncomfortable?'
4. If it's just uncomfortable, the prefrontal cortex tells the amygdala to quiet down.
5. Serotonin and GABA (the brain's main calming signal) carry the 'settle down' message.
6. Your heart rate drops. Your breathing slows. The emotion fades to a manageable level.
7. You keep a steady picture of yourself and the other person: 'I'm still me. They're still mostly good. This moment doesn't define everything.'

What breaks

The prefrontal cortex—the very part that's supposed to calm big feelings—never develops properly when a child grows up in an invalidating or traumatic environment. Early and repeated stress physically shrinks this brake region and leaves the amygdala on constant high alert. The brain learns that emotions are overwhelming and people are unpredictable, so it never builds the wiring for steady self-image or trust.

CLINICAL TERM

Prefrontal-limbic dysregulation: reduced prefrontal cortical volume and function (especially dorsolateral and ventromedial regions) plus amygdala hyperreactivity. Neurotransmitter abnormalities include low serotonin tone and altered endogenous opioid response. Gene-environment interaction: heritable emotion-regulation vulnerability amplified by childhood invalidation, neglect, or abuse. The resulting phenotype is affective instability, identity disturbance, and interpersonal hypersensitivity.

How the body compensates

The body tries to manage unbearable emotions through short-term behavioral crutches—self-harm for endorphin relief, splitting to simplify overwhelming ambiguity, and frantic efforts to avoid abandonment. These strategies work just long enough to prevent total breakdown but deepen the cycle and raise suicide risk.

1. {'what': 'Self-injury', 'why': "Cutting, burning, or hitting releases endorphins (the body's natural painkillers) and briefly overrides emotional pain with physical sensation. It gives a sense of control when feelings are chaotic. But it becomes a learned reflex and carries infection risk, scarring, and accidental deeper injury."}
2. {'what': 'Splitting (all-good / all-bad thinking)', 'why': "When the prefrontal cortex can't hold a nuanced view of people, the brain defaults to extremes: 'You're perfect' or 'You're terrible.' This simplifies overwhelming ambiguity but destroys relationships when the person inevitably disappoints and is suddenly 'all bad.'"}}
3. {'what': 'Frantic efforts to avoid abandonment', 'why': "The amygdala reads even neutral cues (a delayed text, a tired look) as proof of abandonment. She calls repeatedly, threatens, or clings. The effort is exhausting and often pushes people away—confirming her worst fear."}
4. {'what': 'Dissociation under peak stress', 'why': "When emotional pain exceeds the brain's capacity, the circuit that anchors 'this is me in my body right now' shuts off. She feels unreal, detached, or outside herself. It's a last-ditch circuit breaker to prevent total overload, but it is terrifying and disorienting."}

5. {'what': 'Suicidal gestures and attempts', 'why': "Some are true attempts driven by unbearable pain and hopelessness. Some are desperate signals: 'Please see how much I'm suffering.' Both are dangerous—five to ten percent will die by completed suicide over their lifetime. Every gesture must be taken seriously."}
6. {'what': 'Polypharmacy', 'why': 'Patients and providers reach for multiple medications (antidepressants, mood stabilizers, antipsychotics, benzodiazepines) trying to calm symptoms. No drug fixes the core emotion-regulation deficit. Overmedication brings side effects, drug interactions, and false hope that a pill will solve what only skill-building therapy can address.'}

Domino effect

1. A child grows up in an environment that invalidates, ignores, or punishes her emotions—or worse, exposes her to abuse, neglect, or chaos. — The prefrontal cortex is still wiring up during childhood. It needs a steady, validating environment to learn how to regulate feelings. When that environment is absent or hostile, the brake circuits never develop properly. The amygdala, getting no calming signal from above, stays on high alert. The brain learns: 'Emotions are overwhelming. People are unpredictable. I can't trust anyone to stay.'
2. Her prefrontal cortex remains weak, and her amygdala stays hyperreactive into adulthood. — Chronic stress physically shrinks prefrontal gray matter and enlarges the amygdala. Serotonin tone stays low. The result: she has no dimmer switch on feelings. Small stresses trigger intense anger, fear, or emptiness that flood her completely.
3. Her sense of self becomes unstable—she doesn't know who she is or what she wants. — The prefrontal cortex normally holds a steady narrative: 'This is me. These are my values. These are my goals.' Without a working prefrontal brake, her self-image shifts with each emotion. When she's angry, she's bad. When she's loved, she's good. There's no stable core to anchor to.
4. She lives in terror that people will abandon her and scans every interaction for proof. — Her hyperactive amygdala reads neutral cues—a delayed text, a distracted look—as threats. The broken prefrontal cortex can't reality-check those alarms, so every small thing feels like abandonment. She clings, demands reassurance, or lashes out pre-emptively.
5. She splits people into all good or all bad, flipping between the two in moments. — A healthy brain holds contradictions: 'This person is mostly kind but sometimes frustrating.' Her damaged prefrontal cortex can't do that nuanced math. Instead, she defaults to extremes. When you're close, you're perfect. When you say no or disappoint her, you're suddenly all bad. The flip is instant and total.

6. She hurts herself—cutting, burning, hitting—to manage unbearable emotional pain. — Self-injury releases endorphins, the body's natural painkillers, and replaces chaotic emotional pain with controllable physical pain. It works in the moment, so the brain learns it as a go-to tool. But it's dangerous (infection, scarring, accidental deep injury) and deepens shame and isolation.
7. She makes suicidal gestures or attempts when the pain peaks. — Some gestures are cries for help: 'Please see how much I'm suffering.' Some are true attempts to end unbearable pain. The broken prefrontal cortex can't generate hope or plan a future, and low serotonin removes the brake on impulse. Both types are life-threatening—five to ten out of every hundred will die by completed suicide.
8. Under extreme stress, she feels unreal, detached from her body, or like she's watching herself from outside. — This is dissociation: the brain's last-ditch circuit breaker. When emotional pain exceeds capacity, the circuit that says 'this is me, right here, right now' shuts off to prevent total system overload. It feels like being a ghost or watching a movie of your own life. It's terrifying but protective—a way to survive unbearable moments by disconnecting.

Symptoms · what the patient feels, and why

- Frantic efforts to avoid real or imagined abandonment: Early attachment wounds—times when caregivers were unpredictable or left emotionally—wire the amygdala (the brain's alarm center that flags danger) to see separation as a life-or-death threat. The person will beg, threaten, or cling to keep others close.
- Unstable, intense relationships that swing between 'you're perfect' and 'you're the worst': Splitting happens when the prefrontal cortex (the front of the brain that weighs pros and cons) is overwhelmed by strong feelings from the amygdala. The person cannot hold both good and bad in mind at once, so they flip all-or-nothing—loving someone one moment, hating them the next.
- Unstable sense of self—who I am, what I want, and what I believe change often: The sense of self is built and held steady by connections between the prefrontal cortex and deeper emotion centers like the anterior cingulate cortex (the brain's self-awareness hub). When those circuits are weak or stormy, identity feels like shifting sand—goals, values, and even career plans can change suddenly based on mood or who is nearby.
- Impulsive behavior in at least two areas that can cause harm (spending, sex, substance use, reckless driving, binge eating): Low serotonin (a brain messenger that steadies impulses and mood) and an overactive reward circuit in the ventral striatum (driven by dopamine surges) make it hard to pause before acting. The prefrontal cortex's brakes are weak, so urgent feelings turn straight into action without weighing the cost.

- Recurrent suicidal behavior, gestures, threats, or self-harm (cutting, burning): Self-harm releases endorphins (the body's natural pain-killers) that briefly numb unbearable emotional pain. Suicidal thoughts spike when the person feels trapped, invisible, or like a burden. Both are attempts to manage feelings the brain cannot soothe on its own.
- Mood swings—irritability, anxiety, or sadness lasting hours to a few days: The limbic system (the brain's emotion hub, especially the amygdala) fires intensely and the prefrontal cortex cannot dial it down fast enough. External triggers—rejection, criticism, even a look—can flip mood in minutes. Low GABA (the brain's main calming signal) makes the swings harder to stop.
- Chronic feelings of emptiness: When the brain's reward and meaning circuits in the ventral tegmental area are underactive (low dopamine and serotonin), life feels flat and hollow, like a room with no furniture. The person often tries to fill the void with risky behavior, relationships, or substances.
- Intense, hard-to-control anger or physical fights: The amygdala overreacts to perceived slights, flooding the body with adrenaline and cortisol (the body's main stress hormone). The prefrontal cortex, which normally pumps the brakes on rage, is overwhelmed, so anger explodes before the person can think it through.
- Stress-related paranoia or feeling detached from your body or reality (dissociation): Under extreme stress, cortisol floods the brain and the hippocampus (the brain's memory and context center) goes offline. The brain may shut off awareness—like flipping a circuit breaker—to protect itself, leaving the person feeling numb, unreal, or watching themselves from outside.

Focused exam

- Mood swings every few minutes or hours—angry, then tearful, then panicky—all in one conversation: In Borderline Personality Disorder, the prefrontal cortex (the front part of the brain that pumps the brakes on big feelings) is smaller and fires more weakly than it should. She cannot smooth out her emotions, so they spike and crash like a roller coaster.
- Fresh cuts, burns, or scars on forearms, thighs, or wrists: Self-harm in Borderline Personality Disorder is not a suicide attempt—it is an emergency brake. When feelings flood too hard, hurting herself releases tension and feels like relief. This tells you her emotion-control circuit is overwhelmed.
- All-or-nothing talk about the same person: 'She is perfect' on Monday, 'She is evil' on Wednesday (splitting): Borderline Personality Disorder damages the part of the brain that blends good and bad into a realistic picture. She sees people as all good or all bad, never in between. This drives relationship chaos and fear of being left.

- Suicide threat or self-harm right after a perceived rejection—even a missed text or a canceled plan: Fear of abandonment is the core wound in Borderline Personality Disorder. When she feels left behind, her brain's survival alarm (the amygdala, which flags danger) fires like she is in mortal danger. A neutral event feels like the end of the world.
- Feeling unreal, numb, or watching herself from outside her body during stress (dissociation): When the stress-calming circuit overloads, the brain disconnects from the body and emotions as a last-ditch protection. This tells you her prefrontal cortex has shut down under the pressure.
- Deep, empty loneliness; frantic efforts to avoid being alone, even staying in harmful relationships: Borderline Personality Disorder wires the brain to feel abandoned too easily and to panic without someone nearby. This drives the cycle of crises, clinging, and explosive relationship patterns.

Diagnostics · every test answers a question

- Urine drug screen (UDS) → Is substance use driving impulsivity, mood swings, or the crisis?
- Comprehensive metabolic panel (CMP) including glucose, calcium, sodium, potassium, creatinine, liver enzymes → Is a medical cause (low glucose, high calcium, liver failure, kidney failure) mimicking mood swings or confusion?
- Thyroid-stimulating hormone (TSH) → Is thyroid disease causing mood instability or impulsivity?
- Blood alcohol level → Is alcohol intoxication or withdrawal causing the crisis or clouding her thinking?
- Structured clinical interview using DSM-5-TR criteria for Borderline Personality Disorder (5 or more of 9 criteria present for longer than 1 year across many situations) → Does this pattern of abandonment fear, unstable identity, impulsivity, mood swings, emptiness, anger outbursts, self-harm, and dissociation fit Borderline Personality Disorder?
- Columbia-Suicide Severity Rating Scale (C-SSRS) → Is there a current plan and intent to die, or is this an impulsive gesture without lethal planning?

Treatment logic

- Dialectical Behavior Therapy (DBT) for 12 or more months: weekly individual therapy, weekly skills group (mindfulness, distress tolerance, emotion regulation, interpersonal effectiveness), phone coaching for crises, and therapist team consultation → Teaches the brain to tolerate distress without splitting into all-good or all-bad thinking. Strengthens the prefrontal cortex's ability to pause and regulate when emotions flood. Reduces self-harm, suicidal gestures, and relationship chaos by 50 to 70 percent in trials.

- SSRI (sertraline 50 to 200 mg daily or fluoxetine 20 to 60 mg daily) for 6 to 8 weeks, then reassess mood swings and impulsivity → Raises serotonin (a brain messenger that steadies mood and slows racing thoughts) to stabilize the brake pedal on emotions. Reduces the frequency and intensity of mood swings. May reduce impulsive aggression.
- Mood stabilizer (valproate 500 to 1500 mg daily or lamotrigine 100 to 200 mg daily) if impulsivity and self-harm persist after SSRI trial → Slows the electrical noise in the brain circuit that makes her act without thinking. Reduces self-injury urges and anger outbursts. Lamotrigine also lifts depressive symptoms without triggering mania.
- Low-dose atypical antipsychotic (aripiprazole 2 to 10 mg daily or olanzapine 2.5 to 10 mg daily) only if brief psychotic symptoms or severe dissociation occur during crises → Dampens dopamine (a brain messenger that drives intense thoughts and feelings) to reduce the feeling of unreality and disconnection during acute stress. Helps her think clearly enough to use coping skills.
- Brief crisis hospitalization (3 to 7 days) only if active suicide plan with means and high intent; discharge to intensive outpatient DBT immediately → Removes immediate lethal danger and stabilizes her so she can return to therapy and community supports. Short stays prevent learned helplessness and hospital dependence.
- Coordinate social supports: stable housing, sleep hygiene plan (consistent bed and wake time, no screens 1 hour before bed), substance-use treatment if needed, and validate her emotional pain while setting clear, consistent boundaries on harmful behavior → Removes the daily stress that floods her stress-brake circuit. Consistent routines and reliable adults show her that people can be trustworthy—the opposite of her fear of abandonment. Sleep deprivation worsens impulsivity and mood swings.

Medications

Dialectical Behavior Therapy (DBT) · First-line psychotherapy (not a medication)

DBT is a type of talk therapy built for people who feel emotions very intensely and have trouble riding them out without doing something they regret. It teaches four main skills. First, mindfulness: paying attention to right now without judging yourself. Second, distress tolerance: getting through a crisis without making it worse—like holding ice instead of cutting, or using paced breathing when you want to scream. Third, emotion regulation: learning to name what you feel, understand why it showed up, and cool it down before it takes over. Fourth, interpersonal effectiveness: asking for what you need and saying no without burning bridges or exploding. The therapist listens and says your pain is real (that is validation), but also coaches you to try new ways to cope. Standard DBT runs once a week for at least a year, with a skills group, plus phone coaching between sessions when a crisis hits. Over time, this

repeated practice teaches your brain to pause between feeling and acting, so emotions feel less like tsunamis and more like waves you can ride.

1. Every week in one-on-one sessions, you and the therapist walk through a recent crisis and spot the thought or feeling that sparked the urge to act—maybe to hurt yourself, lash out, or quit a relationship.
2. Skills training teaches you concrete steps to tolerate distress: hold ice in your hand, splash cold water on your face, breathe slowly in a pattern, distract yourself with a puzzle or loud music. The urge can pass without harm.
3. Phone coaching between sessions gives you real-time support when a crisis hits at home, so you do not fall back into old patterns when no one is watching.
4. Over months and months, this practice strengthens the prefrontal cortex's control over the amygdala, so the alarm does not hijack you as often. Emotions still come, but you can let them pass instead of drowning in them.

SAFETY WATCH

At the start, you agree to a safety contract: you promise to call the therapist or a crisis line before acting on any urge to hurt yourself or end your life. · If you miss four sessions in a row or keep refusing to try the skills, standard DBT may pause until you re-commit. The therapy only works if you show up and do the work. · Watch for burnout in the therapist, family, or friends who are supporting you. This is hard work for everyone, and your support team needs breaks and their own care too. · If you are actively hearing voices that are not there (psychotic), so revved up you cannot sleep for days (acutely manic), or so drunk or high you cannot focus, those crises need to be stabilized first before DBT can restart.

Mentalization-Based Therapy (MBT) · Second-line psychotherapy (not a medication)

MBT helps you step back and think about what is going on in your own mind and in other people's minds. When feelings run high, people with personality disorders often lose this ability—you might assume everyone else is attacking you, or you cannot figure out why you feel so desperate. MBT, done weekly for 12 to 18 months, teaches you to pause and wonder: What am I feeling right now? Why might I be feeling this? What might the other person be thinking or feeling? Then you test those guesses before reacting. This slows the jump from emotion to action and makes relationships less explosive. You learn to hold two ideas at once—like 'I feel hurt' and 'Maybe they did not mean to hurt me'—without spiraling into rage or despair.

1. The therapist asks you to describe a recent fight or crisis. Then they gently ask: What do you think was going through your mind? What might the other person have been thinking? What clues did you see?

2. By practicing these questions over and over, you rebuild the habit of pausing before assuming the worst. You learn that your first guess about someone's motive might be wrong.
3. Over time, the prefrontal cortex and the brain's social-thinking circuits reconnect. You get better at mentalizing—seeing yourself and others as thinking, feeling beings whose actions have reasons.

SAFETY WATCH

If thinking about your feelings becomes overwhelming, the therapist must dial back and help you calm down and ground before continuing. · MBT assumes you can talk and reflect. If you are actively dissociating (feeling completely disconnected from your body) or psychotic, stabilize that first. · Track suicide risk at every session. Reflecting on pain can sometimes make it feel sharper in the short term before it gets better.

Selective Serotonin Reuptake Inhibitor (SSRI) — e.g. fluoxetine (Prozac), sertraline (Zoloft) · Antidepressant

SSRIs are used when a personality disorder comes with major depression or severe anxiety that therapy alone has not lifted. They block the pump that vacuums serotonin (a brain messenger that steadies mood, sleep, and impulse control) back into the nerve cell after it delivers its message. When you block that pump, more serotonin stays in the gap between cells and keeps the calming, mood-lifting signal going. This helps you feel less hopeless, less irritable, and a bit more able to use the therapy skills you are learning. SSRIs do NOT treat the core personality pattern—they only cushion the depression or anxiety layered on top, so therapy can work better.

1. You swallow the pill. It travels through your stomach into your blood and crosses the barrier into your brain.
2. The drug parks on the serotonin reuptake pump on the sending nerve cell and blocks it, like putting a cork in a drain.
3. Serotonin that would have been sucked back in stays in the synapse longer, so the calming, steadying signal keeps going.
4. Over 4 to 6 weeks, the extra serotonin nudges your brain to grow new connections between cells and dials down the amygdala's alarm volume. You start to feel steadier and less crushed by hopelessness.

SAFETY WATCH

In the first few weeks, some people—especially those under age 25—feel MORE restless, agitated, or have stronger thoughts of suicide. Check in closely with your doctor during this window. If the thoughts get urgent, call your doctor or a crisis line right away. · Common side effects: nausea (especially in the first week), headache, trouble reaching orgasm, or feeling emotionally flat (like you cannot cry or get excited). · Do not stop the medicine suddenly. Your brain has adjusted to the extra serotonin, and stopping cold can cause dizziness, electric-shock feelings (called brain zaps), or rebound anxiety. Taper slowly with your doctor's help. · Watch for serotonin syndrome if you take other drugs that boost serotonin, like tramadol (a pain pill), certain migraine medicines (triptans), or the herbal supplement St. John's wort. Symptoms: confusion, fever, fast heart rate, sweating, muscle jerks. This is rare but dangerous. Go to the ER if it happens. · SSRIs can flip some people into mania if they have hidden bipolar disorder. If you suddenly have racing thoughts, need almost no sleep, feel invincible, or start wild spending or risky sex, stop the SSRI and get a full mood evaluation right away.

Mood stabilizer — e.g. lamotrigine (Lamictal), valproate (Depakote) · Anticonvulsant used off-label for emotion regulation

Mood stabilizers are prescribed when you swing between rage and despair several times a day, or when impulsive behavior—like spending sprees, dangerous driving, binge drinking, or drug use—keeps wrecking your therapy progress. These drugs were designed to prevent seizures (the electrical storms in the brain that cause convulsions), but they also calm overactive electrical firing in the limbic system (your brain's emotion center). Lamotrigine works by slowing the release of glutamate (the brain's main 'go' signal), which dampens emotional surges. Valproate boosts GABA (the brain's main 'stop' signal) and also blocks the sodium channels that let nerve cells fire too fast. Either way, your brain's emotional thermostat steadies, so feelings do not spike as wildly or as often.

1. You take the pill every day. It enters your blood and crosses into your brain.
2. Lamotrigine parks on sodium channels and slows how fast they open, so nerve cells cannot fire as frantically. It also dials down the release of glutamate, the 'go' signal.
3. Valproate increases GABA (the calming messenger) and blocks sodium channels, together muffling the electrical storms that trigger sudden rage or despair.
4. Over weeks, the emotional highs and lows flatten out. You still feel things, but the swings are not as violent. Impulsive urges lose some of their urgency, so you have a moment to think before you act.

SAFETY WATCH

Lamotrigine: Start at a very low dose and go up slowly. A rash can appear in the first 8 weeks. Most rashes are harmless and go away, but a severe rash called Stevens-Johnson syndrome can be life-threatening. If the rash spreads fast, blisters, peels, or comes with fever, mouth sores, or eye redness, stop the drug immediately and go to the ER. ·

Valproate: Can cause weight gain (sometimes a lot), hair thinning, hand tremor, and—rarely—liver damage or pancreatitis (inflammation of the organ that makes digestive enzymes). Your doctor will check liver enzymes and blood counts at the start and then periodically. Valproate is extremely dangerous to a developing fetus—it causes neural tube defects (the brain and spine do not form right). If you can become pregnant, you must use reliable birth control and talk through the risks before starting. · Both drugs can make you drowsy at first. Do not drive or use heavy machinery until you know how you react. · Do not stop suddenly. Taper slowly to avoid rebound mood swings or, with valproate, a risk of seizure even if you never had one before.

Low-dose atypical antipsychotic — e.g. olanzapine (Zyprexa), quetiapine (Seroquel), aripiprazole (Abilify) · Antipsychotic used off-label for brief psychotic or dissociative episodes

In a personality crisis, some people briefly lose touch with reality. You might see or hear things that are not there, believe everyone is plotting to hurt you, or feel completely disconnected from your own body (like you are watching yourself from outside). A low dose of an atypical antipsychotic can quiet these episodes. These drugs mainly block dopamine D2 receptors. Dopamine is a brain messenger tied to reward, motivation, and—when it floods certain brain circuits—delusions (false beliefs) and hallucinations (seeing or hearing things that are not real). Blocking D2 receptors dials down the false signals. Some of these drugs also block serotonin 5-HT_{2A} receptors, which helps with mood and impulsivity. The goal is **SHORT-TERM** use—days to a few weeks—just long enough to restore your grip on reality, then taper off and rely on therapy. These are not meant to be taken every day for months unless you have a separate psychotic disorder like schizophrenia.

1. You swallow the pill. It travels into your blood and crosses into your brain.
2. The drug parks on dopamine D2 receptors and blocks them, so the flood of dopamine cannot trigger false beliefs or voices.
3. Some also block serotonin 5-HT_{2A} receptors, which calms agitation and smooths out wild mood swings.
4. Within hours to days, your thinking clears. Paranoia fades. You can tell what is real again and re-engage with therapy.

SAFETY WATCH

Metabolic effects: Weight gain, rising blood sugar, and rising cholesterol are common, especially with olanzapine and quetiapine. Your doctor will check fasting glucose and lipids at the start, at 3 months, and then every year. If you already have diabetes or prediabetes, these drugs need close monitoring. · Sedation: These drugs can make you very sleepy, especially at first. Do not drive, use machinery, or make big decisions until you know how you react. · Movement side effects: Even the newer 'atypical' antipsychotics can cause stiffness, inner restlessness (akathisia—it feels like you have to pace or bounce your legs), or—rarely—tardive dyskinesia (involuntary chewing, tongue movements, or facial grimaces that may not go away). Using the lowest dose for the shortest time lowers this risk. · Olanzapine and quetiapine carry higher weight-gain and diabetes risk than aripiprazole. Your doctor may choose aripiprazole if you are already overweight or have diabetes. · Do not use antipsychotics just to chemically restrain someone because they are upset, angry, or demanding. Use them only for true psychosis (losing touch with reality) or severe dissociation.

Avoid polypharmacy · Prescribing principle (not a drug)

People in personality crises often end up on four, five, or more psychiatric medications at once because every new symptom gets a new pill added. This is called polypharmacy, and it rarely helps. The pills interact with each other, side effects pile up (tiredness, weight gain, confusion), and it becomes impossible to tell which drug is doing what. Current guidelines say: pick ONE target symptom—like depression, impulsivity, or a brief psychotic break—and use ONE medication for it. Give it a fair trial (usually 6 to 8 weeks) at the right dose. If it does not work, stop it. Add a second drug only if a completely separate symptom needs treatment (for example, an SSRI for crushing depression PLUS a short-term antipsychotic for a psychotic episode that is layered on top). Never add a drug just to 'try something' or because you feel pressured. Keep the medication list short and the reason for each pill crystal clear.

SAFETY WATCH

At every visit, review the medication list with your doctor and ask: Is this still needed? What is it doing? Can we taper one off? · Watch for drug-drug interactions, especially if you see multiple doctors who do not talk to each other. Bring a complete list of every pill, supplement, and over-the-counter medicine you take to every appointment. · Remember: more pills do not equal better care. Therapy plus one or two carefully chosen medications almost always beats a long, confusing medication list.

Brief psychiatric hospitalization · Crisis intervention (not a medication)

If you are acutely suicidal (planning to end your life right now), have just made a serious suicide attempt, or are so psychotic or dissociated that you cannot keep yourself safe, a short hospital stay—typically 3 to 7 days—can break the immediate danger. The hospital removes access to means of harm (pills, sharps, belts, cords), provides 24-hour monitoring so you are never alone, rapidly adjusts medications if needed, and connects you to outpatient therapy and social supports before you leave. The goal is stabilization, NOT long-term residence. Longer stays often make personality crises worse—being locked up can deepen feelings of helplessness, trigger more self-harm to get attention or relief, and delay you learning to cope in the real world with real supports.

1. You are admitted to a locked psychiatric unit where anything that could be used for self-harm—razors, shoelaces, cords, glass—is secured or removed.
2. A psychiatrist evaluates you for other conditions layered on top of the personality pattern—like major depression, bipolar disorder, or psychosis—and adjusts medication if needed.
3. A social worker or case manager helps line up what you will need after discharge: outpatient therapy appointments, housing, food, transportation, insurance, medication refills.
4. Once you can promise to keep yourself safe (contract for safety) and have a solid plan in place, you are discharged—usually within a week.

SAFETY WATCH

Do not admit someone to the hospital just because they are very upset or threatening self-harm as a way to communicate distress. Admit only when the risk of death or serious injury is truly imminent right now and outpatient safety planning has failed. Otherwise you can accidentally teach that threatening harm is the way to get care. · Discharge planning should start on day one, so you do not sit in the hospital for weeks with nothing happening. · Make sure your outpatient therapist knows you are being discharged and can see you within a few days. Gaps in care—when you leave the hospital and have no one to call—are when crises reignite. · Watch for trauma triggers in the hospital. Being restrained, forced to take medication, strip-searched, or denied privacy can retraumatize you and make the crisis worse, not better.

Validate feelings, not behaviors · Therapeutic approach (not a medication)

People with personality disorders often feel that no one understands how much pain they are in. When a doctor, nurse, or therapist dismisses your feelings—saying 'You are overreacting,' 'Calm down,' or 'It is not that bad'—you either shut down and stop talking, or you escalate (yell, threaten, self-harm) to prove how bad it really is. Validation means the clinician says out loud that your feeling makes sense given what you have been through, even if the behavior

that followed is not okay. For example: 'It makes total sense that you felt abandoned and crushed when your friend canceled plans at the last minute. That would hurt anyone who has been left before. And also, cutting yourself to cope is not safe, and we need to find another way together.' This calms the emotional storm just enough that you can hear feedback about the behavior. It is a core piece of DBT and should guide every single clinical conversation.

1. You describe a crisis or an intense feeling to the clinician.
2. The clinician names the feeling and acknowledges that it is real and understandable: 'That sounds incredibly painful. I can see why you felt that way.'
3. Then the clinician separates the feeling from the behavior: 'Your pain is real and makes sense. The way you tried to cope—cutting, yelling, storming out—put you or someone else in danger, so let us figure out a safer way to get through it next time.'
4. This two-step (validate the feeling, then redirect the behavior) keeps you engaged and listening instead of defensive and shut down, so you can actually absorb what comes next.

SAFETY WATCH

Do not validate behavior that harms you or others. Only validate the underlying feeling that drove the behavior. · Be consistent. If a clinician validates you in one visit and dismisses you in the next, you lose trust and stop opening up. · If you interpret validation as permission to keep doing the harmful behavior, the clinician must clarify immediately: 'I hear your pain and it is real. I am not saying the cutting is okay. We have to find a different way.'

Address social determinants of health (housing, substance treatment, therapy access, food, transportation) · Integrated care principle (not a medication)

Personality crises almost never happen in a vacuum. You might be homeless, not know where your next meal is coming from, actively using alcohol or drugs to numb the pain, stuck in an abusive relationship, or sleeping three hours a night because your apartment is loud, cold, and unsafe. No amount of medication or therapy will stabilize you if you do not have a safe place to sleep, regular meals, and access to the supports you need to survive day to day. Every treatment plan must include a checklist: Are you housed, or are you sleeping on a friend's couch or in a shelter? Do you have Medicaid or insurance to pay for therapy and medications? If you are using opioids, alcohol, or methamphetamine, are you offered medication for addiction treatment (like buprenorphine for opioids or naltrexone for alcohol) at the same visit, not told to 'get clean first'? Is your sleep environment safe and quiet? Do you have food? Can you get to appointments, or do you need bus tokens or rides? Addressing these social

determinants is not optional or extra—it is the foundation on which all psychiatric treatment rests. Without it, nothing else works.

1. At the first visit, the doctor or social worker screens for housing, food security, substance use, insurance, and access to care. They ask directly: Where are you sleeping? Do you have enough food? Are you using drugs or alcohol? Can you get to appointments?
2. They connect you to a social worker or case manager who can help you apply for benefits (like Medicaid, SNAP food stamps, disability), find transitional housing or a shelter bed, or arrange medical transportation.
3. If you have opioid use disorder or alcohol use disorder, they offer medication for addiction treatment on the spot—buprenorphine, methadone, or naltrexone—because waiting for you to 'get clean' almost never works and people die waiting.
4. They make sure you have a phone (even a free government phone) and a stable address where appointment reminders, prescription refills, and benefit letters can reach you.
5. They revisit these supports at every follow-up visit, because social stability can crumble fast—you can lose housing, run out of food, or relapse in a week.

SAFETY WATCH

Do not prescribe medications you cannot afford or cannot pick up. Check the cost and whether your pharmacy is open and reachable before writing the prescription. · Do not discharge someone from the hospital to the street or to an unsafe home. That guarantees they will be back in the ER within days. · If you are in active danger from a partner, parent, or other family member, the team must connect you to domestic violence services, a safety plan, and—if needed—emergency shelter right away. · Be honest about wait times. If therapy has a three-month wait list, say so up front and offer a support group, peer support line, or crisis hotline in the meantime so you are not alone.

Red flags

- Active suicidal plan or recent attempt—always means immediate safety evaluation and often psychiatric admission
- Severe dissociation or psychotic symptoms (hearing voices, paranoia that others are plotting against you)—can mean the person is in crisis and at high risk for self-harm
- Escalating self-harm (deeper cuts, more dangerous methods like ligatures or overdoses)—signals worsening distress and rising suicide risk
- Substance intoxication during a crisis—sharply raises impulsivity and the chance of lethal action
- Stating they feel like a burden or have no reason to live—strong predictors of suicide attempt in the near future

Nursing priorities

1. Assess suicide risk every shift using a structured tool (ask about plan, means, intent, and recent self-harm), and remove dangerous items (belts, cords, sharp objects, pills) from the room if risk is present. — The brain's emotion-regulation circuits (prefrontal cortex and amygdala connections) are disrupted in personality disorders, so distress can escalate to life-threatening action within minutes; environmental safety reduces impulsive self-harm.
2. Set clear, consistent limits on behavior (visit times, phone calls, staff check-ins) and communicate the same plan across all shifts so every nurse responds the same way. — Inconsistent responses (one nurse allowing extra time, another saying no) trigger abandonment fears and anger, which activate the amygdala (emotion center) and worsen emotional storms; predictability calms the nervous system.
3. Teach and practice one grounding skill each shift (hold ice, name five things you see, deep breathing for six counts in and six out) and coach the patient to use it when distress starts, before it peaks. — The prefrontal cortex (thinking brain) goes offline during emotional crises, so practicing skills when calm builds automatic pathways; early use of skills can stop the amygdala hijack that leads to impulsive self-harm.
4. Respond calmly and matter-of-factly to emotional outbursts or threats, without extra attention or anger, and return to normal interaction as soon as the patient is calm. — The brain's reward system (dopamine pathways) learns quickly; if crises bring more staff time or dramatic responses, the behavior increases; neutral responses teach that calm communication gets needs met.
5. Validate feelings (say 'I hear that you're in a lot of pain right now') while redirecting harmful behavior (add 'and I can't let you hurt yourself; let's try the ice instead'). — Validation activates the brain's social-safety circuits (ventral vagal system), which calms the fight-or-flight response; pairing it with redirection teaches that feelings are okay but dangerous actions are not.
6. Administer prescribed medications (SSRI like sertraline, mood stabilizer like valproate, or short-term benzodiazepine like lorazepam 0.5 to 1 mg as needed) on time, and watch for sedation or worsening suicidal thoughts in the first two weeks of an SSRI. — SSRIs increase serotonin (a calming brain chemical) but can temporarily increase agitation and suicide risk in young adults before mood improves; mood stabilizers dampen overactive emotional circuits in the limbic system.
7. Collaborate with the treatment team to create a written safety plan that lists the patient's warning signs, coping skills, people to call, and reasons to stay safe, and review it with the patient daily. — The hippocampus (memory center) stores concrete plans better than vague advice; having a written, rehearsed plan gives the prefrontal cortex a roadmap to follow when the amygdala hijacks decision-making during a crisis.

8. Document objective observations (exact words, actions, times) rather than judgments ('patient is manipulative'), and share patterns with the psychiatrist to guide Dialectical Behavior Therapy referral or medication changes. — Personality pathology involves lifelong patterns, not single events; tracking triggers, behaviors, and responses over days reveals the patterns that DBT targets and shows whether medications are reducing emotional intensity or impulsivity.

Sources

- American Psychiatric Association DSM-5-TR (2022)
- NICE Borderline Personality Disorder Guideline (2023)
- Cochrane Review: Psychological Therapies for BPD (2020)
- Journal of Clinical Psychiatry: Neurobiological Basis of BPD (2023)

Last reviewed: 2026-07-07