

Postpartum Depression

Postpartum depression can start any time during pregnancy or in the first year after giving birth.

THE STORY IN ONE SENTENCE

Postpartum depression can start any time during pregnancy or in the first year after giving birth.

“After birth, a sudden drop in hormones plus sleep loss and huge life stress can push the brain's mood circuits into a stuck-down state that won't lift on its own.”

Normal anatomy — the body's job

A healthy brain keeps mood steady through a balance of chemical messengers—serotonin (steadies mood and sleep), dopamine (sparks joy and motivation), and norepinephrine (powers energy and focus)—working across the limbic system (the deep emotion centers) and the prefrontal cortex (the front part that steadies feelings and keeps perspective).

BODY'S JOB

The limbic system reads your emotional world and sets your baseline feeling tone. The prefrontal cortex acts like the brakes, calming big feelings and helping you see the bigger picture. The hypothalamus (a small control center deep in the brain) talks to the pituitary gland (a pea-sized gland just below the brain) and the adrenal glands (two small glands sitting on top of your kidneys) to manage your stress response—this is called the HPA axis. When everything works, these parts keep your mood in a healthy range, let you feel joy, give you energy to care for your baby, and help you sleep when you get the chance.

1. Pregnancy floods the body with estrogen and progesterone (the main pregnancy hormones), which keep serotonin, dopamine, and norepinephrine levels high.
2. After delivery, estrogen and progesterone drop sharply within 24 to 48 hours—sometimes by 90 percent or more.
3. In most women, the limbic system and prefrontal cortex adapt to the hormone drop and brain-chemical levels rebalance within a few weeks.
4. Sleep, social support, and low stress give the brain time and space to adjust.

5. Mood stays steady, joy returns, energy comes back, and the new mother can bond with her baby.

What breaks

The same hormone crash that ends pregnancy can push brain-chemical levels so low that the mood circuits get stuck in a down state. The prefrontal cortex (the part that steadies feelings) and the limbic system (the deep emotion centers) cannot pull mood back up on their own. Lost sleep, isolation, past depression, childhood trauma, and big life stress make the stuck state much worse and much harder to escape.

CLINICAL TERM

Postpartum depression is a major depressive episode with onset during pregnancy or within the first year postpartum, driven by the rapid withdrawal of estrogen and progesterone after delivery, which destabilizes serotonin, dopamine, and norepinephrine signaling in the limbic system and prefrontal cortex. The HPA axis (hypothalamic-pituitary-adrenal stress circuit) becomes overactive, flooding the body with cortisol (the main stress hormone). Risk is amplified by sleep deprivation, lack of social support, prior mood disorder, childhood adversity, and psychosocial stressors.

How the body compensates

In the early days, the body tries to push through with adrenaline (the fight-or-flight hormone) and sheer willpower. A new mother may look like she is coping—going through the motions of feeding, changing, and soothing her baby—while inside, sadness, guilt, and scary thoughts are growing. This is the dangerous window: she may not ask for help because she feels ashamed, fears judgment, or believes she should be able to handle it alone.

1. {'stage': 'First days to weeks', 'what': 'The body runs on adrenaline and the drive to protect the baby. The mother may look tired but functional.', 'hidden': 'Inside, mood is sinking, joy is gone, and guilt or scary thoughts are starting—but she hides it, fearing she will be seen as a bad mother.'}
2. {'stage': 'Weeks to months', 'what': 'Sleep debt piles up, social isolation grows (especially if she cannot leave the house or feels too ashamed to reach out), and the brain's reward circuits (powered by dopamine) shut down further.', 'hidden': 'She may stop eating well, stop caring for herself, withdraw from her partner or family, and feel numb or detached from her baby—while still going through the motions of care.'}

3. {'stage': 'Critical danger point', 'what': 'Suicidal thoughts appear, or she has intrusive thoughts of harming the baby (these are unwanted, terrifying thoughts—not desires—but they cause overwhelming guilt and fear).', 'hidden': 'She may not tell anyone because the shame is so great. Suicide is a leading cause of maternal death in the first year postpartum. In rare cases (<1 in 1,000), postpartum depression can tip into postpartum psychosis (a break from reality with hallucinations or delusions), which is a psychiatric emergency because the risk of harm to mother or baby is high.'}

Domino effect

1. Within hours after delivery, estrogen and progesterone drop by up to 90 percent. — These hormones kept serotonin, dopamine, and norepinephrine levels high during pregnancy. When they crash, brain-chemical levels fall with them—like a wave pulling back from the shore.
2. Serotonin (the brain messenger that steadies mood and sleep) drops, and the limbic system (the deep emotion centers) cannot stay balanced. — Low serotonin makes the amygdala (the alarm center) more sensitive to threat and the prefrontal cortex (the steady-feeling center) less able to calm it down. Sadness feels heavier, anxiety spikes, and sleep stays broken even when the baby sleeps.
3. Dopamine (the messenger that sparks joy and motivation) falls, and the brain's reward system goes quiet. — The nucleus accumbens (the reward and pleasure center) stops lighting up in response to the baby's smile, a warm meal, or time with loved ones. Nothing feels good. This is anhedonia—the loss of joy—and it is one of the core signs of depression.
4. The HPA axis (the stress circuit from brain to adrenal glands) gets stuck in overdrive, flooding the body with cortisol. — High cortisol keeps the body in a state of alarm, breaks down deep sleep, weakens the prefrontal cortex further, and amplifies guilt and negative thoughts. The brain interprets normal new-parent stress as overwhelming danger.
5. Sleep deprivation from nighttime baby care prevents the brain from clearing waste, rebalancing chemicals, or repairing mood circuits. — Deep sleep is when the brain does its housekeeping—flushing out toxins, restoring serotonin and dopamine, and strengthening the prefrontal cortex. Without it, the stuck-down state gets worse every day.
6. Guilt, shame, and scary intrusive thoughts flood in, and the mother withdraws or hides her feelings. — The weakened prefrontal cortex cannot stop the amygdala from firing alarm signals or block unwanted, frightening thoughts (like images of harm coming to the baby). She feels like a bad mother, fears judgment, and often does not ask for help.

7. In severe cases, suicidal thoughts appear, or—rarely—postpartum psychosis develops (hallucinations, delusions, or a break from reality). — When brain chemistry stays broken, despair can grow into thoughts of self-harm or escape. Postpartum psychosis is a psychiatric emergency and requires immediate hospitalization because the risk of harm to mother or baby is very high.
8. Without treatment—therapy, medication, support, and rest—the depression deepens, mother-infant bonding suffers, and long-term outcomes for both worsen. — Early intervention (screening with tools like the Edinburgh Postnatal Depression Scale, starting therapy such as cognitive-behavioral or interpersonal therapy, considering SSRIs if needed, mobilizing family support, and protecting sleep) can reverse the circuit breakdown and restore mood. Untreated postpartum depression can last months to years and harm the child's emotional and cognitive development.

Symptoms · what the patient feels, and why

- Feeling sad, empty, or hopeless most of the day, nearly every day: After birth, estrogen and progesterone (hormones that were sky-high during pregnancy) crash suddenly. This steep drop affects serotonin and dopamine (brain messengers that regulate mood and pleasure), leaving the emotional centers of the brain—especially the limbic system (the brain's feeling hub)—struggling to stay balanced.
- Loss of interest or pleasure in things that used to feel good: The reward pathways in the brain—powered by dopamine (the 'feel-good' messenger)—lose their signal strength when hormone levels plummet after delivery. The nucleus accumbens (the brain's reward center) stops lighting up the way it should, so even caring for the baby or hobbies can feel flat and empty.
- Extreme tiredness or no energy, even after rest: Sleep deprivation from newborn care disrupts the hypothalamus (the brain's control tower for sleep and energy). At the same time, the body is recovering from birth, inflammation is still settling, and thyroid hormone levels can dip temporarily—all draining the body's fuel reserves.
- Trouble bonding with or feeling connected to the baby: Oxytocin (the 'bonding hormone' released during skin-to-skin contact and breastfeeding) may not rise normally when depression dampens the brain's emotional circuits. The prefrontal cortex (the brain's planning and emotion-regulating center) also struggles to process warm feelings when serotonin is low, making connection feel forced or absent.
- Feeling worthless, guilty, or like a bad mother: The anterior cingulate cortex (a brain region that tracks mistakes and self-judgment) becomes overactive when serotonin drops. This creates a loop of harsh self-criticism and exaggerated guilt, even over small things or situations beyond the mother's control.

- Trouble concentrating, remembering things, or making decisions: The prefrontal cortex and hippocampus (the brain's memory and decision-making centers) need steady dopamine and norepinephrine (brain messengers that sharpen focus and memory) to work well. When these messengers drop after childbirth, thinking gets foggy and choices feel overwhelming.
- Changes in appetite—eating much more or much less than usual: The hypothalamus controls hunger signals. When serotonin and dopamine fall, the brain's ability to sense fullness or hunger gets scrambled, leading some mothers to lose interest in food and others to seek comfort in eating.
- Thoughts of harming yourself or the baby, or thoughts that you or the baby would be better off dead: Severe depletion of serotonin (which helps regulate mood and impulse control) combined with relentless stress and sleep loss can overwhelm the prefrontal cortex's ability to stop dark, intrusive thoughts. The brain's safety brakes weaken, and dangerous ideas that would normally be dismissed can feel compelling or even logical.
- Hearing voices, seeing things that aren't there, or having bizarre beliefs (signs of postpartum psychosis): In rare cases, the sudden hormone crash after birth—combined with inflammation, autoimmune triggers, or genetic vulnerability—can destabilize dopamine pathways so severely that the brain loses its grip on reality. The thalamus (the brain's sensory relay station) and the temporal lobes (which process sounds and perceptions) begin to misfire, creating hallucinations and delusions.

Focused exam

- Edinburgh Postnatal Depression Scale (EPDS) score ≥ 10 : The EPDS is a 10-question form that catches postpartum depression early. A score of 10 or higher means the brain's mood circuit—the parts that keep feelings steady and make you feel okay—is stuck in the 'off' position and needs help now.
- Flat or sad face, slow speech, poor eye contact: When the mood circuit breaks in postpartum depression, the face and voice show it first—like a battery running low, the whole system slows down. This tells you the problem is real and in the brain, not just tiredness from caring for a baby.
- Mom says she has no joy in her baby or anything else (anhedonia): Loss of joy is the core break in postpartum depression—the nucleus accumbens (the brain's pleasure and reward center) goes offline, so nothing feels good anymore. This separates postpartum depression from normal new-mom tiredness, which still allows moments of happiness.
- Direct question: 'Have you thought about harming yourself or your baby?' Answer is yes or unclear: Suicide is the top cause of death in mothers in the year after birth. Postpartum depression makes these thoughts real and scary. You must ask every time and write down the answer—it can save a life.

- Sleep and appetite changed even when baby is asleep or fed (not just from baby care): In postpartum depression, the hypothalamus (the brain's control tower for sleep, hunger, and body rhythms) breaks—mom can't sleep even when she has the chance, and food tastes like nothing. This proves the brain chemistry is wrong, not just the schedule.
- Signs of psychosis: mom sees or hears things not there, or fears she will hurt the baby with no reason to: Postpartum psychosis is a different, more dangerous condition—it means the break from reality is happening, usually from too much dopamine (a brain messenger that when too high causes hallucinations and delusions). This is a separate emergency—hospital now.

Diagnostics · every test answers a question

- Edinburgh Postnatal Depression Scale (EPDS) → Is the mom's mood low enough to be depression, not just normal tiredness?
- Thyroid-stimulating hormone (TSH) and free T4 → Is the thyroid gland (the body's metabolism control center in the neck) working right, or is it broken and causing the low mood?
- Complete blood count (CBC) → Does the mom have anemia (low red blood cells) from blood loss at birth or low iron, which can make tiredness and sadness worse?
- Ferritin (iron storage level) → Are the body's iron stores empty, even if the blood count looks okay?
- 25-hydroxyvitamin D level → Is vitamin D (which the brain needs to make mood messengers) too low?
- Urine drug screen (if history or concern for substance use) → Is the mom using substances (alcohol, opioids, cannabis, stimulants) that could be causing or worsening the mood symptoms?

Treatment logic

- EPDS screening at every postpartum visit (2 weeks, 6 weeks, and any time concern arises) → Catches depression early before it deepens. A score of 10 or higher means the brain's stress-stopping circuit—the prefrontal cortex (the front of the brain that pumps the brakes on big feelings) and the hippocampus (the memory center that helps make sense of stress)—is already struggling. Finding it now stops the spiral before it gets worse.
- Sertraline 50-200 mg daily (SSRI antidepressant) or escitalopram 10-20 mg daily, started right away → These medicines block serotonin (a brain messenger that steadies mood and sleep) from being vacuumed back up too fast, so more stays in the space between brain cells. This lets the mood circuit work again. They take 2-4 weeks to show full effect, but some relief starts earlier.

- Weekly talk therapy: cognitive-behavioral therapy (CBT) or interpersonal therapy (IPT) → CBT rewires the thinking patterns that feed the depression—like breaking the loop of 'I am a bad mother' by testing that thought against real evidence. IPT helps fix relationship problems and role changes (like becoming a mom) that trigger the depression. Both strengthen the prefrontal cortex's ability to regulate feelings.
- Brexanolone IV (60-hour continuous infusion in hospital) or zuranolone oral (2.5 mg daily × 14 days) for moderate-to-severe cases → These target the allopregnanolone pathway—allopregnanolone is a brain steroid (made from progesterone) that calms the amygdala (the brain's alarm center) by boosting GABA (the brain's main calming signal). After birth, allopregnanolone drops sharply, and these medicines restore it. They work in days instead of weeks, which matters when mom is suffering hard or having thoughts of self-harm.
- Sleep support: dedicated night helper (partner, family, doula, or night nurse) so mom gets at least one 5-hour block of unbroken sleep each night → Sleep loss is both cause and symptom—it amplifies the broken stress circuit by keeping cortisol (the body's main stress hormone) high and blocking the hippocampus from clearing out stress memories. Protecting sleep lets the brain reset and makes every other treatment work better.
- Ask directly every visit: 'Do you have thoughts of harming yourself or your baby?' If yes → immediate same-day psychiatric evaluation, safety plan (remove guns, pills, sharp objects), and consider hospitalization → Identifies severe suicidal risk or postpartum psychosis (a medical emergency where the brain loses touch with reality). These need hospital care, often with antipsychotic medicines like olanzapine or haloperidol for psychosis, and close monitoring to prevent tragedy.
- Lactation support + safe medicine choice: sertraline and escitalopram pass into breast milk in tiny amounts (safe); brexanolone and zuranolone are also safe; avoid paroxetine and benzodiazepines → If breastfeeding matters to mom, choosing the right medicine lets her stay connected to her baby and keep the bonding and health benefits of nursing while treating her brain. This reduces guilt and preserves the relationship.

Medications

Sertraline (Zoloft) · SSRI (selective serotonin reuptake inhibitor)

Sertraline blocks the pump that normally sucks serotonin (the brain messenger that steadies mood and sleep) back out of the gap between nerve cells. When serotonin stays in the gap longer, the receiving cell gets a stronger, steadier signal. Over a few weeks, this helps lift the dark cloud of depression and makes it easier to feel joy again. Sertraline passes into breast milk in very tiny amounts — the lowest of all SSRIs — so doctors often pick it first when a mom is breastfeeding.

1. Sertraline enters the bloodstream and crosses into the brain.

2. It lands on the serotonin transporter (the pump on the sending nerve cell that normally pulls serotonin back inside).
3. By blocking that pump, sertraline lets serotonin stay in the synapse gap longer.
4. The receiving nerve cell picks up more serotonin signal, which over time helps rebalance mood circuits.
5. After 2–4 weeks of daily use, the brain adjusts and depression symptoms begin to lift — mood improves, sleep steadies, and interest in life returns.

SAFETY WATCH

Watch for worsening sadness, thoughts of harming yourself or the baby, or new agitation in the first few weeks — call the doctor immediately if these appear. · Common early side effects: upset stomach, loose stools, headache, and trouble falling asleep. These usually fade after the first week or two. · Do NOT stop suddenly — sertraline must be tapered down slowly to avoid withdrawal symptoms like dizziness, electric-shock feelings, and rebound anxiety. · Safe during breastfeeding at standard doses (sertraline passes into milk in the smallest amounts of any SSRI), but discuss with your doctor so the baby can be watched. · Rarely, SSRIs can trigger serotonin syndrome (too much serotonin, causing confusion, fast heart rate, sweating, and muscle jerking) if combined with certain other medicines — tell every provider what you are taking.

Escitalopram (Lexapro) · SSRI (selective serotonin reuptake inhibitor)

Escitalopram works the same way sertraline does — it blocks the pump that pulls serotonin back out of the gap between mood nerve cells, letting the calming, mood-lifting signal stay stronger. It is very clean and selective, meaning it targets serotonin pumps tightly and causes fewer side effects for many people. Escitalopram does pass into breast milk in slightly higher amounts than sertraline, but is still considered compatible with breastfeeding under medical supervision.

1. Escitalopram is absorbed into the bloodstream and crosses into the brain.
2. It binds to the serotonin transporter on the sending nerve cell, blocking the pump.
3. Serotonin stays in the synapse longer, strengthening the signal to the receiving cell.
4. Over 2–4 weeks, the mood circuits rebalance and depressive symptoms ease — sadness lifts, sleep improves, and energy returns.
5. Full benefit usually appears by 6–8 weeks of daily use.

SAFETY WATCH

Watch for new or worse thoughts of harming yourself or the baby, especially in the first few weeks — contact your provider immediately if this happens. · Common early side effects: nausea, headache, dry mouth, and drowsiness or trouble sleeping. Most fade within a week or two. · Do NOT stop suddenly — taper slowly under medical guidance to avoid withdrawal symptoms like dizziness, brain zaps, and irritability. · Safe during breastfeeding for most women, but the baby should be monitored by the pediatrician. · Rarely can cause serotonin syndrome if combined with other serotonin-boosting drugs — always share your full medication list with every provider.

Brexanolone (Zulresso) · Neuroactive steroid (allopregnanolone analog)

Brexanolone is a lab-made copy of allopregnanolone, a natural calming hormone the body makes from progesterone. During pregnancy, allopregnanolone levels soar to help keep mood steady. After delivery, they crash fast — in some women, this sudden drop overwhelms the brain's calming circuits and triggers severe depression. Brexanolone refills that calming signal by turbocharging GABA (the brain's main brake pedal), which quiets overactive fear and sadness circuits. Unlike SSRIs, brexanolone can lift depression in hours to days, not weeks.

1. Brexanolone is given as a continuous IV drip over exactly 60 hours in a certified medical facility.
2. It enters the brain and docks onto GABA-A receptors, making them much more sensitive to GABA (the brain's natural calming messenger).
3. When GABA lands on these supercharged receptors, the calming signal becomes much stronger, quieting the overactive fear and sadness circuits.
4. Depressive symptoms — hopelessness, severe sadness, inability to bond with the baby — often begin to lift within 24–48 hours.
5. The benefit lasts after the infusion ends, though some women need additional antidepressant treatment to maintain improvement.

SAFETY WATCH

Requires a certified treatment center and continuous pulse-oximetry monitoring because brexanolone can cause sudden severe sleepiness and loss of consciousness. · You cannot care for your baby alone during the infusion — a support person must be present the entire time. · Common side effects: extreme drowsiness, dry mouth, flushing, and dizziness. · Very expensive (often over \$30,000 per treatment) and not always covered by insurance — financial counseling and prior authorization are essential. · Safe during breastfeeding, but close monitoring of the baby is recommended during and after the infusion.

Zuranolone (Zurzuvae) · Neuroactive steroid (allopregnanolone analog)

Zuranolone works just like brexanolone — it mimics allopregnanolone, the natural calming hormone that drops sharply after childbirth. It supercharges GABA-A receptors (the brain's main calming switches), letting GABA (the brain's brake signal) work much more powerfully. This quiets the overactive fear and sadness circuits that drive postpartum depression. The major advantage of zuranolone is that it comes as a once-daily pill taken for 14 days, so you can take it at home instead of needing a 60-hour hospital IV infusion.

1. Zuranolone is swallowed once daily in the evening for 14 days.
2. It is absorbed into the bloodstream and crosses into the brain.
3. It docks onto GABA-A receptors and makes them far more responsive to GABA (the brain's natural calming messenger).
4. With stronger GABA signaling, the overactive fear and sadness circuits quiet down.
5. Many women feel better within a few days, and improvement often continues even after the 14-day course ends.

SAFETY WATCH

Can cause severe drowsiness — do NOT drive, operate machinery, or care for your baby alone until you know how it affects you. · Common side effects: sleepiness, dizziness, diarrhea, fatigue, and cold-like symptoms. · Take in the evening with a fatty meal to improve absorption. · Do NOT drink alcohol while taking zuranolone — it increases the risk of dangerous sedation. · Very expensive (often several thousand dollars for the 14-day course) and may not be covered by insurance — prior authorization and financial assistance programs are often needed. · Pregnancy must be avoided while taking zuranolone and for one week after the last dose — use reliable contraception. · Safety during breastfeeding is not fully established — discuss risks and benefits with your doctor.

Red flags

- Any thought of harming yourself or the baby
- Hearing voices, seeing things that are not real, or believing things that make no sense to others (possible postpartum psychosis)
- Unable to care for yourself or the baby, or feeling completely detached from the baby
- Severe anxiety or panic attacks that make it hard to function
- Rapid mood swings, confusion, or bizarre behavior that started suddenly after delivery

Nursing priorities

1. Ask every postpartum patient about thoughts of self-harm or harming the baby at every visit using direct questions — Suicide is the leading cause of death in the postpartum year and infanticide (killing the baby) happens in severe untreated postpartum depression, so direct screening saves lives even though it feels uncomfortable to ask
2. Complete the Edinburgh Postnatal Depression Scale (EPDS) screening tool at the 2-week and 6-week postpartum visits and any time the mother or family expresses concern — The EPDS is a validated 10-question screening test that catches postpartum depression early because many mothers hide their symptoms due to shame or fear of being judged as a bad mother
3. Assess whether the mother can safely care for herself and the baby, including feeding, diapering, and recognizing when the baby needs help — Severe depression causes psychomotor retardation (slowed movement and thinking) and poor concentration that prevent the mother from meeting the baby's basic needs for nutrition and safety
4. Identify and document the mother's support system including who lives at home, who can help with baby care, and whether the family believes postpartum depression is real — Social isolation and lack of practical help increase cortisol (stress hormone) levels that worsen depression, and family members who do not understand mental illness may prevent the mother from getting treatment
5. Teach the mother and family that postpartum depression is a medical illness caused by hormone changes and brain chemistry, not a character weakness or bad mothering — Shame and guilt about having depression delay treatment by an average of 6 months, and understanding the biological cause of depression reduces self-blame and increases treatment acceptance
6. Coordinate same-week mental health referral for therapy and psychiatry evaluation, and confirm the mother has transportation to get to appointments — Cognitive-behavioral therapy (talk therapy that changes negative thought patterns) works as well as medicine for mild to moderate postpartum depression, and combining both treatments gives the best outcomes
7. Review safe medication use during breastfeeding and reassure mothers that sertraline and escitalopram are compatible with breastfeeding — Many mothers refuse antidepressant medicine because they fear harming the baby through breast milk, but untreated maternal depression harms infant brain development more than the tiny amount of medicine in breast milk
8. Schedule follow-up contact within 48 to 72 hours after starting antidepressant medicine to check for worsening depression or new suicidal thoughts — Antidepressants can increase suicide risk in the first 2 weeks of treatment before mood improves, especially in women under age 25, so close monitoring in the early treatment period prevents self-harm

Sources

- ACOG Practice Bulletin No. 212: Screening and Diagnosis of Mental Health Conditions During Pregnancy and Postpartum, 2024
- American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5-TR)
- U.S. Preventive Services Task Force (USPSTF) Recommendation on Perinatal Depression, 2023
- Postpartum Support International Clinical Guidelines

Last reviewed: 2026-07-07