

Septic Arthritis

Bacteria invade the joint space, fill it with pus, and destroy cartilage fast—hours matter.

THE STORY IN ONE SENTENCE

Bacteria invade the joint space, fill it with pus, and destroy cartilage fast—hours matter.

“Septic arthritis: one hot, swollen, red joint the patient refuses to move, plus fever. Tap it now, start IV antibiotics, then wash it out in the OR—delay kills the joint.”

Normal anatomy — the body's job

A healthy joint is a sealed, sterile pocket lined by a thin tissue called the synovium, which makes slippery fluid so the bones glide without grinding. No bacteria live inside.

BODY'S JOB

The synovium acts like a quiet factory: it filters blood, pulls out water and nutrients, and releases just enough slippery fluid (synovial fluid) to keep the cartilage-coated bone ends sliding smoothly—like oil in a hinge. The joint capsule around it is a tough bag that keeps germs out and fluid in.

1. Picture two bones meeting, each capped with shiny white cartilage (like the smooth plastic tip on a chair leg).
2. Around them is a stretchy bag (the joint capsule), and inside that bag is a slick lining (the synovium) that sweats out a few drops of clear, honey-thick fluid.
3. That fluid fills the tiny space between the cartilage caps so they never touch bone-on-bone—every time you bend your knee or elbow, the cartilage glides on that slippery film.
4. Normally, the synovium is thin, pink, and germ-free—it has its own blood supply but keeps bacteria out because the capsule is sealed tight.

What breaks

Bacteria get into the sealed joint space, multiply, and trigger a massive immune attack that fills the joint with pus and destroys the cartilage cushion within hours.

CLINICAL TERM

Bacterial invasion of the synovial space, most often *Staphylococcus aureus* (including MRSA) in all ages, *Streptococcus* (including Group A and B) in neonates and young children, *Kingella kingae* in toddlers, and *Neisseria gonorrhoeae* in sexually active adolescents and adults. Entry routes: hematogenous seeding (bacteremia from skin infection, endocarditis, IV drug use, or central line), direct inoculation (penetrating trauma, arthrocentesis, intra-articular injection), or contiguous spread (adjacent osteomyelitis or soft-tissue abscess).

How the body compensates

The body tries to wall off the infection with swelling, fever, and pain signals, and the child guards the joint by refusing to move it—these are protective reflexes, not the disease itself, but they buy only a few hours before cartilage is destroyed.

1. {'what': 'Fever and systemic inflammatory response', 'how': 'White cells release cytokines (chemical alarms like interleukin-1 and tumor tissue death (necrosis) factor) that reset the brain's thermostat in the hypothalamus, raising body temperature to make the environment less friendly for bacteria.', 'cost': 'Fever helps slow bacterial growth but also makes the child miserable, signals sepsis risk, and does not stop cartilage destruction.'}
2. {'what': 'Joint a fluid collection (effusion) (swelling) and heat', 'how': 'Blood vessels in the synovium dilate and leak fluid and white cells into the joint, trying to dilute bacteria and deliver more immune fighters; increased blood flow makes the skin over the joint hot and red.', 'cost': 'Swelling raises pressure inside the rigid joint capsule, which crushes cartilage and chokes off its nutrient supply—so the very response meant to help also speeds damage.'}
3. {'what': 'Pain and refusal to move (guarding)', 'how': 'Nerve endings in the joint capsule fire when stretched by pus; the brain interprets this as severe pain and triggers muscle spasm around the joint to hold it still and prevent further injury.', 'cost': 'Guarding protects the joint from motion that would hurt, but it also delays diagnosis if the limb looks 'just stiff' rather than infected—and every hour of delay lets more cartilage dissolve.'}
4. {'what': 'Localized redness and warmth', 'how': 'Increased blood flow and inflammatory mediators dilate capillaries in the skin and tissue overlying the infected joint, making it visibly red and warm to touch.', 'cost': 'These signs alert you to infection, but they appear only after bacteria have already been multiplying for hours—by the time the joint looks angry, cartilage damage is underway.'}

Domino effect

1. Bacteria enter the joint space — Most commonly, bacteria (especially *Staph aureus*) travel through the bloodstream from a distant infection—skin abscess, pneumonia, infected IV site—and seed the synovium, which is rich in blood vessels. In toddlers, a tiny cut or throat infection can spill *Kingella* into the blood; in sexually active teens and adults, gonorrhea spreads from the genitals. Less often, a puncture wound (stepping on a nail, a dog bite, a contaminated injection) injects germs directly, or an adjacent bone infection tunnels through.
2. Bacteria multiply on the synovium and in the synovial fluid — The warm, nutrient-rich synovial fluid is an ideal culture medium with no immune cells ready to fight at first. Bacteria double every 20–30 minutes, quickly reaching massive numbers. The synovium, normally a quiet factory, becomes inflamed and leaky.
3. The immune system floods the joint with white blood cells (neutrophils) — Chemical alarms (cytokines, chemokines) summon millions of neutrophils from the blood into the joint. These cells try to engulf and kill bacteria by releasing enzymes (proteases, elastase) and toxic oxygen radicals. The fluid turns thick, cloudy, and yellow-green—classic pus.
4. Pressure inside the joint skyrockets; the joint swells, turns hot and red, and hurts with any movement — The joint capsule is a tough, inelastic bag—pus has nowhere to expand, so pressure climbs. High pressure compresses the tiny blood vessels that feed cartilage, starving it of oxygen and nutrients. Nerve endings in the capsule fire intensely, causing severe pain. Blood flow to the synovium increases (heat, redness). The child holds the limb completely still because even a tiny bend stretches the capsule and hurts.
5. Enzymes from dying white cells digest the cartilage cushion — The same proteases meant to kill bacteria also break down collagen and proteoglycans—the proteins that make cartilage smooth and springy. Bacterial toxins add to the damage. Cartilage has no blood supply and cannot repair itself; once the matrix is chewed up, it is gone forever.
6. Fever spikes as the infection spreads into the bloodstream (bacteremia, sepsis risk) — Cytokines released in the joint spill into the bloodstream, signaling the hypothalamus (the brain's thermostat) to raise body temperature. If bacteria escape the joint and multiply in the blood, sepsis can develop—blood pressure drops, organs fail—especially in infants, elderly, and immunocompromised patients.

7. Without urgent treatment, the cartilage is destroyed, the joint space collapses, and permanent arthritis sets in — If pus sits in the joint for more than 24–48 hours, cartilage is so damaged that even after the infection clears, bone grinds on bone. The joint becomes stiff, painful, and deformed. In children, damage to the growth plate near the joint can stunt bone growth. That is why septic arthritis is treated as a surgical emergency: tap the joint (arthrocentesis) to confirm pus and identify the bug, start IV antibiotics immediately (before culture results), and wash out the joint in the operating room (arthrotomy or arthroscopy) to remove all pus and infected tissue—speed is the only way to save the cartilage.
8. Trace every clue backward to find the upstream source — If you see a hot, swollen knee in a child who will not walk, ask: Why is there pus in the joint? → Bacteria got in. How? → Bloodstream (check for skin infections, recent illness, IV lines), direct poke (ask about trauma, bites, injections), or spread from nearby bone (osteomyelitis). Finding and treating the source (drain an abscess, pull an infected line, treat gonorrhea) prevents recurrence and stops the bacteria from seeding other joints or the heart valves.

Symptoms · what the patient feels, and why

- Hot, red, swollen joint (usually just one): Bacteria invade the joint space and multiply. The immune system floods the area with white blood cells and fluid to fight back, which causes heat, redness, and swelling—like a battle raging inside the joint.
- Severe pain in the joint that gets worse with any movement: The infected fluid stretches the joint capsule (the tight sac around the joint) like an overfilled water balloon, and the immune chemicals irritate the nerve endings, making every tiny move excruciating.
- Not wanting to move or put weight on the joint: Moving the joint or pressing on it hurts so much that the body instinctively holds it still and refuses to use it—a kid might refuse to walk if the knee or hip is infected, and an adult might cradle an infected shoulder.
- Fever and chills: Bacteria release toxins into the bloodstream, and the hypothalamus (the brain's thermostat) cranks up body temperature to help the immune system fight harder—so the person feels hot and shaky.
- The joint looks bigger or puffier than the same joint on the other side: Pus (dead white cells and bacteria) and inflammatory fluid collect inside the infected joint, physically expanding it like a balloon filling with water, so comparing left knee to right knee or left elbow to right elbow shows the difference clearly.
- Warmth when you touch the skin over the joint: Blood vessels dilate (widen) near the infection to rush more immune cells to the fight, and all that extra blood flow heats up the skin—like holding your hand over a heater.

- **Fast heart rate:** The body is under stress from infection, so the sympathetic nervous system (the 'fight-or-flight' system) and circulating cytokines (immune alarm signals) speed up the heart to pump more oxygen and white cells where they're needed.
- **Recent skin infection, IV drug use, or a prosthetic (artificial) joint in the same spot:** Bacteria on the skin can ride the bloodstream into a joint, especially if there's a cut, boil, or needle puncture. Fake joints have no blood supply to patrol for germs, so bacteria stick more easily and can hide on the plastic or metal surface.

Focused exam

- **One joint that is hot, swollen, red, and extremely tender** — usually the knee, hip, shoulder, ankle, or wrist: Septic arthritis picks one joint nearly every time because bacteria land in one spot from the bloodstream. The heat and puffiness mean pus is building inside the joint capsule, not just a bruise or sprain. This is bacteria multiplying in a space that should be completely sterile.
- **The person refuses to move that joint even a tiny bit** — any motion causes severe pain: This refusal to move is the body's protective splinting reflex. Moving the joint squeezes the pus-filled capsule, which sends pain signals screaming through nerve endings. Compare the healthy side: the difference is dramatic and immediate.
- **Fever, usually above 101°F (38.3°C)**, though 30–40% of children and some adults have no fever at all: Fever means the immune system is fighting bacteria that spread through the bloodstream. The absence of fever does NOT rule out septic arthritis — infants, older adults, and people with weak immune systems often have no fever even with a raging joint infection.
- **Recent skin break, surgery, IV line, or history of IV drug use near the affected joint or anywhere on the body:** These are the entry doors bacteria use to get into the bloodstream. Septic arthritis rarely appears out of nowhere — trace back to the break in the skin barrier or the medical device that gave bacteria access to blood.
- **The limb is held in a bent, relaxed position** — hip bent and turned outward, knee slightly bent, elbow partly flexed: This posture makes the joint capsule expand to its largest volume, which lowers the pressure from pus inside and eases pain. The body naturally finds this position without thinking about it.

Diagnostics · every test answers a question

- **Joint aspiration (arthrocentesis)** — synovial fluid cell count, Gram stain, culture, and crystal exam → Is this infection inside the joint space, and if so, what bacteria is causing it?
- **Blood cultures** (two sets from separate sites before starting antibiotics) → Are bacteria circulating in the bloodstream, and what type are they?

- Inflammatory markers — ESR (erythrocyte sedimentation rate) and CRP (C-reactive protein) → Is there active inflammation in the body right now?
- Complete blood count (CBC) with differential → How hard is the immune system working to fight infection?
- Plain X-ray of the affected joint → Is there bone damage, gas in the soft tissues, or a foreign body like a prosthetic joint?
- MRI of the joint (if diagnosis is unclear or prosthetic joint is involved) → Is infection spreading into bone, surrounding muscle, or abscess pockets that need surgical drainage?
- Gonorrhea and Chlamydia nucleic acid amplification test (NAAT) from urine or genital swab in sexually active patients → Is this septic arthritis caused by *Neisseria gonorrhoeae*, which spreads from genital infection?

Treatment logic

- Urgent sterile joint aspiration (arthrocentesis) BEFORE starting any antibiotics → Removes synovial fluid for culture, Gram stain, cell count, and crystal exam so the lab can identify the exact bacteria and test which antibiotics kill it. Aspiration also relieves some pressure and pain by draining pus. Starting antibiotics before aspiration can make cultures falsely negative and leave you guessing the germ.
- Empiric IV antibiotics — vancomycin PLUS either ceftriaxone or cefepime, started immediately after joint aspiration → Kills the most common bacteria before you know the exact type: *Staphylococcus aureus* (including MRSA, covered by vancomycin), *Streptococcus* species, *Neisseria gonorrhoeae*, and gram-negative rods like *E. coli* and *Pseudomonas* (covered by ceftriaxone or cefepime). Once culture and sensitivity results return, narrow to the single best antibiotic for that specific germ.
- Urgent surgical drainage — arthroscopic or open irrigation and debridement of the joint within 24 hours → Physically washes out pus, fibrin clots, dead white blood cells, and bacteria from the joint space with sterile saline. Removes inflammatory debris that blocks antibiotics from reaching bacteria. Stops bacteria from eating away cartilage and bone. Large joints (knee, shoulder, ankle) can often be cleaned arthroscopically; smaller or deep joints (hip, sacroiliac) may need open surgery. Repeat washout may be needed if pus reaccumulates.
- Continue IV antibiotics for 2 weeks, then switch to oral antibiotics to complete 4–6 weeks total for native joints (longer for prosthetic joints, osteomyelitis, or MRSA) → The long antibiotic course ensures bacteria die completely and do not hide in scar tissue, biofilm, or bone. Switching to oral antibiotics after 2 weeks keeps drug levels high in the blood and joint without needing a hospital stay, as long as the person can swallow pills and the bacteria are sensitive to an oral drug.

- Identify and treat the source of bacteria — remove infected IV lines, drain skin abscesses, treat endocarditis, address IV drug use wounds → Stops new bacteria from seeding the bloodstream and landing in the joint again. Fixes the upstream break in the skin or blood vessel barrier, not just the downstream joint infection. Without source control, bacteria keep entering and the infection can come back even after antibiotics stop.
- Remove or revise prosthetic joint implant if infection does not clear with DAIR (debridement, antibiotics, and implant retention) within 2 weeks or if infection is chronic → Bacteria form a slimy biofilm on metal and plastic implants that shields them from antibiotics and white blood cells. Early infection (less than 3 weeks after surgery) may be saved by washout and antibiotics alone. Late or chronic infection almost always requires removing the implant, treating with 6 weeks of IV antibiotics, then placing a new implant in a staged procedure months later.
- Pain control with acetaminophen or NSAIDs (ibuprofen, naproxen) and physical therapy once acute infection is controlled → Reduces pain and inflammation so the person can start moving the joint gently, which prevents permanent stiffness and muscle wasting. Physical therapy stretches scar tissue and rebuilds strength after weeks of holding the joint still.

Medications

Vancomycin · Glycopeptide antibiotic

Vancomycin kills bacteria by blocking them from building their protective cell walls. Think of bacteria like tiny water balloons — they need a tough outer shell to stay intact. Vancomycin stops them from making that shell, so the bacteria burst and die. It works especially well against MRSA (methicillin-resistant *Staph aureus*), a stubborn germ that often causes joint infections and doesn't respond to many other antibiotics.

1. Vancomycin enters the bloodstream through an IV line, travels through the blood, and reaches the infected joint.
2. It crosses into the joint space and synovial fluid, where bacteria are growing and destroying cartilage.
3. Vancomycin binds tightly to the building blocks bacteria use to make their cell walls — specifically a part called D-alanyl-D-alanine.
4. By blocking this step, bacteria can't finish building their walls. Without a complete wall, the bacteria can't hold their shape or protect themselves.
5. The bacteria's insides spill out, the germ dies, and the immune system clears away the debris.
6. Over days to weeks, the infection load drops, inflammation calms, and the joint can start to heal — though damage already done to cartilage may be permanent.

SAFETY WATCH

Kidney damage — vancomycin can harm the kidneys, especially at high doses or if used with other kidney-toxic drugs. Blood tests (creatinine) are checked regularly, and the dose is adjusted if kidney function drops. · Hearing loss — high levels can damage the inner ear, causing ringing in the ears (tinnitus) or permanent hearing loss. Blood levels are monitored to keep the drug in a safe range. · Red man syndrome — if the IV infusion runs too fast, vancomycin can trigger flushing, itching, and a red rash on the face, neck, and chest. Slowing the infusion prevents this. · Low blood counts — rarely, vancomycin can lower white blood cells or platelets. Blood counts are monitored during long courses. · Allergic reactions — true allergy (hives, swelling, trouble breathing) is uncommon but requires stopping the drug immediately.

Ceftriaxone · Third-generation cephalosporin antibiotic

Ceftriaxone kills bacteria by attacking their cell walls from a different angle than vancomycin. It sneaks into the bacteria's wall-building machinery and jams the gears, so the wall can't finish forming. The bacteria lose their structure and burst. Ceftriaxone covers a wide range of germs, including *Neisseria gonorrhoeae* (the gonorrhea germ, a common cause of septic arthritis in sexually active young adults), many gram-negative bacteria (like *E. coli* and *Klebsiella*), and some gram-positive bacteria (like *Streptococcus*). It's often paired with vancomycin at the start because together they cover almost all the bacteria that cause joint infections.

1. Ceftriaxone is given through an IV or a muscle injection (IM), enters the bloodstream, and travels to the infected joint.
2. It crosses into the joint space and reaches the bacteria multiplying there.
3. Ceftriaxone binds to proteins on the bacteria's surface called penicillin-binding proteins (PBPs), which normally help weave the cell wall together.
4. By blocking these proteins, the bacteria can't link the wall's building blocks. The wall becomes weak and full of holes.
5. Water rushes into the bacteria through the holes, the germ swells, bursts, and dies.
6. The immune system mops up the dead bacteria, inflammation drops, and the joint begins to heal — though cartilage damage may already be done.

SAFETY WATCH

Allergic reactions — people allergic to penicillin have a small risk (about 2–5%) of also being allergic to ceftriaxone. Watch for rash, hives, swelling, or trouble breathing. ·

Diarrhea and *C. difficile* colitis — ceftriaxone kills good bacteria in the gut, which can let *C. difficile* (a dangerous gut germ) overgrow and cause severe, watery, bloody diarrhea. If diarrhea starts during or after treatment, tell the doctor right away. · Gallbladder sludge — ceftriaxone can form crystals in the gallbladder, especially in children or people getting high doses for weeks. This usually goes away after stopping the drug. · Low blood counts — rarely, it can drop white cells, red cells, or platelets. Blood counts are monitored. · Do NOT mix with calcium-containing IV fluids in newborns — it can form dangerous crystals.

Oral antibiotics (step-down therapy) · Varies — often fluoroquinolones (e.g. levofloxacin, ciprofloxacin), trimethoprim-sulfamethoxazole, cephalexin, or amoxicillin-clavulanate, chosen based on culture results

After about 2 weeks of IV antibiotics in the hospital, many patients can switch to pills taken by mouth to finish the treatment course at home. The specific pill depends on which bacteria grew in the joint fluid culture and which antibiotics it's sensitive to. Oral antibiotics work the same way as IV antibiotics — they kill bacteria by blocking cell wall building, protein making, or DNA copying — but they're absorbed from the gut into the bloodstream instead of going straight into a vein. Switching to pills is safe once the infection is clearly improving (less pain, less swelling, fever gone, blood markers like CRP dropping) and the bacteria are known to be susceptible. The total antibiotic course for a natural (not prosthetic) joint is usually 4–6 weeks; for a prosthetic (artificial) joint, it may be months or even lifelong.

1. The patient swallows the antibiotic pill, which dissolves in the stomach or small intestine.
2. The drug is absorbed through the gut lining into the bloodstream.
3. It circulates to the infected joint, crosses into the joint space, and reaches the bacteria.
4. The antibiotic blocks a critical step in the bacteria's survival — building walls, making proteins, or copying DNA — so the bacteria die.
5. Over weeks, the last remnants of infection are cleared, inflammation resolves, and the joint stabilizes — though any cartilage or bone damage is permanent.

SAFETY WATCH

Finish the full course — stopping early can cause the infection to come roaring back, sometimes with bacteria that are now resistant to the antibiotic. · Fluoroquinolones (like levofloxacin, ciprofloxacin) — can cause tendon rupture (especially the Achilles tendon in the heel), nerve damage (tingling, numbness), and mood changes. Avoid in children and pregnant women unless no other option exists. · Trimethoprim-sulfamethoxazole — can cause rash (including serious skin reactions like Stevens-Johnson syndrome), low blood counts, high potassium, and kidney problems. Blood tests are monitored. · Cephalexin and amoxicillin-clavulanate — can cause diarrhea, allergic reactions, and (with amoxicillin-clavulanate) liver inflammation. Watch for yellowing of the eyes or skin. · Drug interactions — some oral antibiotics interfere with blood thinners, diabetes drugs, or seizure medicines. Always tell the doctor and pharmacist about every medication you take. · Take with food or on an empty stomach as directed — absorption can vary, and some drugs cause less stomach upset with food.

Red flags

- Fever above 101°F (38.3°C) plus a hot, swollen joint—infection can destroy cartilage in hours
- Refusal to move or bear weight on a joint in a child—especially hip or knee—may be septic arthritis and needs same-day evaluation
- Joint pain plus signs of sepsis: confusion, very low blood pressure, fast breathing, or mottled skin—bacteria may be spreading through the whole body
- Prosthetic (artificial) joint that suddenly becomes painful, warm, or swollen—infection in a fake joint almost always needs surgery to clean out

Nursing priorities

1. Keep the patient NPO (nothing by mouth) until the orthopedic surgeon evaluates them, because urgent surgical washout may be needed within hours. — The infected joint needs drainage of pus and inflamed tissue (irrigation and debridement) under anesthesia to prevent cartilage destruction; delaying surgery while waiting for the stomach to empty risks permanent joint damage.
2. Protect the painful joint by keeping it immobilized in a neutral position with pillows or a splint, and avoid any passive or active range-of-motion exercises. — Movement spreads bacteria and inflammatory enzymes (proteases) throughout the joint space, which destroys cartilage and bone faster; immobilization reduces pain and limits further tissue damage.

3. Draw blood cultures from two different sites before giving any antibiotics, and prepare the patient for immediate joint aspiration by the physician. — Blood cultures grow bacteria in about 50 percent of septic arthritis cases and identify the exact germ; joint fluid analysis (synovial fluid white blood cell count usually over 50,000 cells per microliter with over 90 percent neutrophils) confirms infection before antibiotics alter results.
4. Administer IV antibiotics (vancomycin and ceftriaxone) within one hour after joint fluid is obtained, using a large-bore IV catheter for reliable access. — Empiric antibiotics (broad-spectrum coverage before culture results) must cover *Staphylococcus aureus* including MRSA (methicillin-resistant *Staph aureus*), *Streptococcus* species, and gram-negative bacteria to stop cartilage destruction, which begins within 8 hours of infection.
5. Monitor vital signs every 2 to 4 hours, watching for fever spikes, a fast heart rate (tachycardia) (fast heart rate over 100), low blood pressure (hypotension) (low blood pressure), or worsening pain despite pain medicine. — Septic arthritis can trigger sepsis (body-wide infection response) with shock, especially in elderly or immunocompromised patients; early recognition of systemic inflammatory response syndrome (SIRS) allows rapid fluid resuscitation and ICU transfer if needed.
6. Assess neurovascular status (circulation, sensation, and movement) of the limb distal to the infected joint every 4 hours, comparing with the opposite side. — Joint swelling and pus under pressure can compress blood vessels and nerves, causing compartment syndrome (dangerously high pressure in the muscle compartments) or starved blood flow (ischemia) (tissue oxygen deprivation); loss of pulses or sensation requires emergency surgical decompression.
7. Give scheduled IV acetaminophen or IV ketorolac for pain control rather than waiting for the patient to request pain medicine, avoiding opioids if possible. — Severe pain from joint capsule stretching and inflammation responds better to around-the-clock anti-inflammatory medicines and acetaminophen; opioids cause sedation that masks mental status changes from sepsis and make it harder to assess worsening infection.
8. Examine the skin over the entire body for wounds, rashes, pustules (pus-filled bumps), or IV drug track marks, and document any findings with photos if hospital policy allows. — The source of bacteremia (bacteria in blood) must be identified and treated; skin infections like cellulitis (skin and soft tissue infection) or endocarditis (heart valve infection) with septic traveling clots (emboli) cause pinpoint bleeding spots (petechiae) (small red spots) or purple bleeding blotches (purpura) (purple patches) that guide diagnosis and treatment duration.

Sources

- IDSA Clinical Practice Guideline for the Diagnosis and Treatment of Native Vertebral Osteomyelitis in Adults (2015)

- British Society for Rheumatology guideline for management of the hot swollen joint in adults (2018)
- CDC Septic Arthritis overview
- AMA JAMA Rational Clinical Examination series on the adult with monoarticular arthritis

Last reviewed: 2026-07-07