Neutropenia

A plain-English guide for you and your family

WHAT IS HAPPENING

Neutrophils are white blood cells made in your bone marrow. Their only job is to hunt down and kill bacteria before those germs can cause infection. Normally you have between 1,500 and 8,000 neutrophils in every drop of blood, ready to swarm any cut, scratch, or invader. When something breaks that system—chemotherapy killing off the marrow cells that make neutrophils, medicines like clozapine or sulfamethoxazole stopping production, viruses like HIV or Epstein-Barr infecting the marrow, your own immune system attacking young neutrophils (autoimmune dangerously low infection-fighting cells (neutropenia)), a genetic condition you were born with (like Kostmann syndrome), or a bone-marrow disease like leukemia or aplastic anemia crowding them out—your neutrophil count drops. Once it falls below 1,500 you are neutropenic, and bacteria that your body normally stops cold can now multiply. If the count drops under 500 (severe dangerously low infection-fighting cells), even the germs living harmlessly in your mouth, gut, or on your skin can slip into your blood and cause sepsis, a body-wide infection that can kill in hours. Every sign—fever without an obvious source, mouth sores, a racing heart, dropping blood pressure—points back to this one upstream problem: missing guards. The body tries to cope by speeding up the heart to move what immune cells are left, but it cannot make new neutrophils fast enough. None of this is random. Trace the weird findings backward and you will see they all flow from the same break: too few neutrophils to fight bacteria.

What you might feel

- Frequent infections (like sinus infections, skin boils, or bladder infections that keep coming back)
- Fever without an obvious cause
- Mouth sores or painful gums
- Sore throat or trouble swallowing
- Skin infections or redness around small cuts
- Feeling unusually tired or weak
- No symptoms at all (found on routine blood test)

- Pneumonia or lung infection

Call the doctor right away if...

- Fever above 100.4°F (38°C) — this can signal a serious infection spreading fast when neutrophils are very low
- Shaking chills or feeling very sick all over — may mean bacteria are in the bloodstream (sepsis)
- Trouble breathing or chest pain — suggests lung infection (pneumonia), which can become life-threatening quickly
- Confusion or sudden weakness — signs that an infection is affecting the brain or that sepsis is causing organs to fail
- ANC below 500 cells/µL (severe dangerously low infection-fighting cells (neutropenia)) — the body has almost no neutrophil defense left, and any infection can turn dangerous within hours

What the care team will do

- Check temperature every 4 hours and notify the provider immediately if it reaches 100.4°F (38°C) or higher.
- Start broad-spectrum IV antibiotics within 1 hour if the patient has fever and ANC below 1500.
- Place the patient in a private room with strict handwashing for all visitors and staff if ANC is below 500.
- Avoid rectal temperatures, suppositories, enemas, and invasive procedures unless absolutely necessary.
- Inspect the mouth, skin folds, IV sites, and perirectal area every shift for redness, swelling, or drainage.
- Hold all live vaccines (MMR, varicella, nasal flu vaccine) and avoid contact with anyone who received a live vaccine in the last 2 weeks.
- Teach the patient to avoid fresh flowers, potted plants, raw fruits and vegetables, and undercooked meat if ANC is under 500.
- Administer G-CSF (filgrastim or pegfilgrastim) as ordered, usually starting 24 hours after chemotherapy ends.

Medicines you may be given — and why

Broad-spectrum antibiotics (e.g., piperacillin-tazobactam, cefepime, or meropenem)

When neutrophils (the white blood cells that fight bacteria) drop dangerously low, the body loses its main defense against infections. Bacteria can multiply fast and cause life-threatening infections (a condition called feverish (febrile) dangerously low infection-fighting cells (neutropenia) when fever appears). Broad-spectrum antibiotics act like substitute soldiers: they kill a wide range of bacteria by breaking open the bacteria's protective outer wall or stopping the bacteria from building new proteins they need to survive. This buys time until the body can rebuild its own neutrophil army.

Filgrastim (G-CSF: granulocyte colony-stimulating factor)

Neutrophils are made in the bone marrow (the spongy center of bones). Chemotherapy, some drugs, or bone marrow diseases can shut down that factory. G-CSF is a natural messenger molecule (a protein) that tells the marrow 'make more neutrophils, faster.' Filgrastim is a lab-made copy of that messenger. It binds to receptors (docking stations) on young white blood cells in the marrow and pushes them to mature into neutrophils and release into the bloodstream. It shortens the dangerous window when your infection-fighting cells are too low.

Intravenous immunoglobulin (IVIG)

In autoimmune dangerously low infection-fighting cells (neutropenia), the immune system mistakenly makes antibodies (attack proteins) that tag neutrophils for destruction, like friendly fire. IVIG is a concentrated solution of antibodies pooled from thousands of healthy donors. When you give it by vein, it floods the bloodstream with normal antibodies that calm the immune attack. The exact way it works is still being studied, but IVIG seems to block the bad antibodies from sticking to neutrophils, distract the spleen and other organs from destroying neutrophils, and dampen the overactive immune signals. It gives the neutrophil count a chance to recover.

Where this information comes from

- American Society of Hematology (ASH) 2023 guidelines
- National Comprehensive Cancer Network (NCCN) febrile neutropenia guideline
- UpToDate neutropenia overview
- CDC healthcare-associated infection prevention

Last reviewed: 2026-07-07