

# Surgical Site Infection

Surgical site infection means germs got into the surgical cut and started multiplying—open the wound to let the pus drain out, send a sample to the lab to find out which germ it is, and give the right antibiotic to kill it.

## THE ONE-LINE TRUTH

A surgical site infection is an infection that starts in the cut made during surgery, appearing within 30 days of the operation—or up to 1 year later if an implant (like a mesh, screw, or artificial joint) was placed.

## The chain — how it unfolds

1. Bacteria enter the surgical wound during or right after the operation. — why: The cut breaks the skin's protective wall, and bacteria—most often *Staphylococcus aureus* (a tough germ that lives on everyone's skin and nose, including the drug-resistant kind called MRSA) or gut bacteria like *Escherichia coli* if surgery was on the belly—land on the exposed tissue underneath. Normally, neutrophils (germ-eating white blood cells) swarm in and kill them within hours, but if the patient has diabetes (high blood sugar slows neutrophils and feeds bacteria like fertilizer), smokes (nicotine chokes blood flow so fewer immune cells and less oxygen arrive), is malnourished (not enough protein and vitamins to make new white blood cells and repair tissue), or has an implant (bacteria stick to the foreign surface and hide in a slimy biofilm shield), the germs multiply faster than the immune system can stop them.
2. The body floods the wound with white blood cells and fluid, causing redness, warmth, swelling, and pus. — why: Infected tissue releases chemical alarms—histamine, bradykinin, and cytokines (alarm proteins like interleukin-1)—that make nearby blood vessels widen and leak. Neutrophils squeeze out of the bloodstream and into the wound, where they eat bacteria and dead tissue and then die themselves, forming pus (a thick yellow or green fluid of dead germs, dead neutrophils, and liquefied tissue). This flood of cells and fluid is the body's attempt to wall off the invader—the redness, heat, and swelling are not the disease itself, but the immune response trying to contain it. The wound becomes tender because the swelling stretches nerve endings.

3. If pus is not drained and bacteria keep multiplying, they burrow deeper into fascia and muscle or spill into the bloodstream. — why: Pus trapped under closed skin acts like a warm, moist greenhouse for bacteria—full of food (dead cells and fluid). Pressure builds and bacteria follow the path of least resistance, sliding along fascial planes (the slippery layers that separate muscle groups) or breaking through tiny blood vessels into the bloodstream. Fierce bacteria like group A *Streptococcus* or *Clostridium perfringens* (which makes gas gangrene) release poisons that blow up tissue and blood vessels directly (causing tissue death (necrosis), meaning the tissue dies), creating more dead tissue and cutting off blood supply, which lets infection spread even faster along whole limbs or body regions.
4. The whole body reacts with fever, racing heart, and a surge in white blood cell production. — why: Macrophages (large scavenger immune cells) at the infection site release cytokines—interleukin-1, interleukin-6, and tumor tissue death (necrosis) factor (alarm proteins)—into the bloodstream. These messengers travel to the hypothalamus (the brain's thermostat in the center of the brain) and reset it higher, causing fever (usually 100.4 to 104 degrees Fahrenheit or 38 to 40 degrees Celsius). The bone marrow (spongy tissue inside bones where blood cells are made) is signaled to crank out more neutrophils, spiking the white blood cell count above 12,000 per microliter. The brainstem's cardiovascular center (in the medulla, the lowest part of the brain) speeds the heart to deliver more oxygen and immune cells to tissues. All of this is the body trying to contain the infection—fever slows how fast bacteria can reproduce, and more white cells means more soldiers to fight—but these defenses can only buy time; they cannot cure the infection without antibiotics and draining the source of pus.
5. If bacteria reach the bloodstream in large numbers, sepsis and septic shock develop. — why: Bacteria and their poisons flood the blood and trigger a huge, body-wide immune response. Cytokines make blood vessels everywhere leaky and floppy, so blood pressure drops and fluid leaks out of vessels into tissues (causing whole-body swelling and starving organs of oxygen). The heart races to keep up, but eventually can't maintain pressure. Lactic acid (a waste product from burning fuel without enough oxygen, called anaerobic metabolism) builds up in the blood because tissues are choking. Kidneys, liver, lungs, and brain begin to fail from lack of oxygen and blood flow. This is septic shock—a life-threatening spiral that, without emergency treatment (IV fluids, antibiotics within one hour of recognizing sepsis, and surgery to drain or remove the infected source), can lead to multi-organ failure and death within 24 to 48 hours.

6. The wound edges may split open (dehiscence) as pus and pressure tear apart fragile healing tissue. — why: Infection destroys the new collagen fibers (the body's scaffolding protein) that are knitting the wound edges together, and rising pressure from trapped pus literally rips the cut open. Dehiscence is the body's crude release valve—better to have an open draining wound than a closed pocket of pus (abscess) that spreads deeper. But an open wound heals much more slowly, requires weeks of packing gauze into it and changing dressings daily, and leaves a wider, weaker scar. If an implant or deep organ is now exposed to the outside, the risk of deeper infection climbs sharply.
7. Every finding—redness, warmth, pus, fever, high white count, fast heart rate, low blood pressure—is the body's response to the same root problem: bacteria multiplying in the wound. — why: These are not separate diseases; they are the immune system and cardiovascular system reacting at local, regional, and body-wide levels to try to contain and fight the infection. The real disease is the unchecked bacterial growth. Treatment must trace backward to the source—drain the pus, cut out dead tissue, remove the infected implant if present, and give antibiotics matched to the exact germ found when wound fluid is sent to the lab and grown in culture—because treating only the fever or the high white count without fixing the source will fail. The body's defenses will eventually wear out, and the patient will crash into septic shock and organ failure if the infection is not controlled at its starting point.
8. The window to turn things around closes fast once septic shock begins. — why: Sepsis guidelines (the Surviving Sepsis Campaign, an international expert panel) say the 'golden hour' matters most—blood cultures and broad-spectrum antibiotics (antibiotics that hit many kinds of bacteria) must be given within 60 minutes of spotting sepsis, and source control (surgery to drain pus or cut out dead tissue) should happen within 6 to 12 hours. Every hour of delay raises the risk of death. Once blood pressure cannot be held up even with IV fluids and vasopressor drugs (medicines like norepinephrine that squeeze blood vessels tight), and organs are failing, the death rate climbs above 30 to 40 percent. The key is to spot the downhill trend early—watch for rising heart rate, falling urine output, worsening pain, spreading redness, or confusion—and act before the patient tips into shock, because stopping the slide is far easier than trying to rescue someone already in free fall.

## Spot it — what you see, and why

---

- Redness and warmth around the surgical cut: Germs (bacteria) get into the wound and multiply. The body's defense system sends extra blood to the spot to fight them off. That extra blood carries white blood cells and oxygen, and it makes the skin look red and feel warm—like a busy highway of defenders rushing to the scene.

- Swelling at or near the incision: When bacteria invade, tiny blood vessels leak fluid into the surrounding tissue as part of the alarm response. That pooling fluid causes the area to puff up, like a sponge soaking up water.
- Pain or tenderness that gets worse instead of better: Chemicals released by injured cells and immune cells—histamine, bradykinin, and prostaglandins (messengers that tell the body something is wrong)—irritate nerve endings. That sends sharper pain signals to the brain, making the wound hurt more as infection grows.
- Pus or cloudy fluid draining from the wound: White blood cells called neutrophils swarm to the site, eat bacteria, and die in the fight. Pus is the thick, creamy mixture of dead white blood cells, dead bacteria, and broken-down tissue. It's a sign the body is battling hard, but it also means infection is present.
- Fever (body temperature above 100.4°F or 38°C): Bacteria release toxins, and immune cells release chemicals called pyrogens—interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor tissue death (necrosis) factor-alpha (TNF-alpha). These pyrogens travel to the hypothalamus (the brain's thermostat in the center of your head) and tell it to crank up the body's temperature. A higher temp helps immune cells work faster and slows bacterial growth.
- Foul-smelling drainage from the incision: Certain bacteria—like *Clostridium*, *Bacteroides*, and some *Staphylococcus* strains—produce gases and waste chemicals as they break down tissue. These chemicals have a sour or rotten smell, signaling that infection is deep and destroying living tissue.

## Red flags — act now

---

- Fever above 101°F (38.3°C) with chills or shaking
- Redness spreading fast—tracking up an arm or leg or moving more than an inch away from the wound edge in a few hours
- Wound edges pulling apart or organs or tissue visible inside the opening
- Feeling confused, very dizzy, breathing fast (more than 22 breaths a minute), or heart racing (over 100 beats a minute at rest)
- Foul smell or dead-looking (black, gray, or green) tissue around the cut

## Treat it — in this order

---

1. Open the wound, drain all pus, cut away dead tissue (debridement), and pack it open or apply negative-pressure wound therapy (vacuum-assisted closure) → Removes the source—pus and dead tissue—where bacteria hide and multiply beyond the reach of antibiotics or blood flow. Oxygen and healing cells can now get in to clean up and rebuild.

2. Start IV antibiotics immediately: vancomycin (covers MRSA) plus piperacillin-tazobactam or a carbapenem (covers gut bacteria) if the wound is near the belly or groin, or if the patient is septic → Kills bacteria circulating in the blood and tissue before they multiply into sepsis or seed other organs like heart valves or bones. Once culture results return, narrow to the specific antibiotic that targets the identified germ.
3. Send wound tissue or fluid for culture and sensitivity testing before starting antibiotics (if possible) or as soon as you open the wound → Tells you the exact species of bacteria and which antibiotics kill it. This lets you switch from broad-spectrum antibiotics (which can harm good bacteria and cause resistance) to the single best drug.
4. If signs of deep infection (pain out of proportion, skin that looks purple or black, a crackling feel or sound (crepitus)—crackling under the skin from gas, or spreading beyond the wound in hours), take the patient to the operating room immediately for wide surgical washout → Stops necrotizing fasciitis (flesh-eating infection) from racing through tissue planes and destroying muscle, fascia, and blood vessels. Washout removes infected and dying tissue and stops toxins from poisoning the bloodstream.
5. Control blood sugar (goal under 180 mg/dL), optimize nutrition (protein at least 1.2–1.5 g/kg/day), stop smoking, and correct anemia or low albumin → Restores the body's own infection-fighting power. High blood sugar paralyzes neutrophils (the main germ-eating white cells). Low protein and oxygen starve the wound of building blocks and fuel for healing.

## Your top priorities at the bedside

---

1. Assess vital signs (temperature, heart rate, blood pressure, respiratory rate, oxygen saturation) and mental status immediately and every 4 hours — why: Fever over 100.4°F, heart rate over 90, respiratory rate over 20, low blood pressure, or confusion indicate sepsis (bloodstream infection causing organ dysfunction) requiring immediate IV fluid resuscitation and antibiotics within 1 hour
2. Inspect the surgical incision and measure the area of redness, swelling, and warmth in centimeters; photograph the wound if policy allows — why: Documenting the size and appearance of redness (erythema) (redness) and hardened tissue (induration) (firm swelling) allows nurses and doctors to track whether the infection is spreading or improving with treatment, and rapid expansion signals necrotizing infection
3. Notify the surgeon immediately if pus-filled (purulent) drainage is present, the wound has opened, or red streaks are visible — why: Surgical site infections require source control (opening the wound to drain pus and remove infected tissue) which only the surgeon can perform, and delaying drainage allows bacteria to multiply and spread into deeper tissue or bloodstream